



Politecnico
di Bari

Repository Istituzionale dei Prodotti della Ricerca del Politecnico di Bari

Equilibrium measures and equilibrium potentials in the Born-Infeld model

This is a post print of the following article

Original Citation:

Equilibrium measures and equilibrium potentials in the Born-Infeld model / Bonheure, Denis; D'Avenia, Pietro; Pomponio, Alessio; Reichel, Wolfgang. - In: JOURNAL DE MATHÉMATIQUES PURES ET APPLIQUÉES. - ISSN 0021-7824. - STAMPA. - 139:(2020), pp. 35-62. [10.1016/j.matpur.2020.05.001]

Availability:

This version is available at <http://hdl.handle.net/11589/147090> since: 2025-01-27

Published version

DOI:10.1016/j.matpur.2020.05.001

Publisher:

Terms of use:

(Article begins on next page)

**EQUILIBRIUM MEASURES AND EQUILIBRIUM POTENTIALS
IN THE BORN-INFELD MODEL**

DENIS BONHEURE, PIETRO D'AVENIA, ALESSIO POMPONIO, AND WOLFGANG REICHEL

ABSTRACT. In this paper, we consider the electrostatic Born-Infeld model

$$(\mathcal{BI}) \quad \begin{cases} -\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = \rho & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0 \end{cases}$$

where ρ is a charge distribution on the boundary of a bounded domain $\Omega \subset \mathbb{R}^N$, with $N \geq 3$. We are interested in its *equilibrium measures*, i.e. charge distributions which minimize the electrostatic energy of the corresponding potential among all possible distributions with fixed total charge. We prove existence of equilibrium measures and we show that the corresponding *equilibrium potential* is unique and constant in $\bar{\Omega}$. Furthermore, for smooth domains, we obtain the uniqueness of the equilibrium measure, we give its precise expression, and we verify that the equilibrium potential solves (\mathcal{BI}) . Finally we characterize balls in \mathbb{R}^N as the unique sets among all bounded $C^{2,\alpha}$ -domains Ω for which the equilibrium distribution is a constant multiple of the surface measure on $\partial\Omega$. The same results are obtained also for Taylor approximations of the electrostatic energy.

RÉSUMÉ. Dans cet article, nous considérons le modèle électrostatique de Born-Infeld

$$(\mathcal{BI}) \quad \begin{cases} -\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = \rho & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0 \end{cases}$$

où ρ est une distribution de charges sur le bord d'un domaine borné $\Omega \subset \mathbb{R}^N$, $N \geq 3$. Nous nous intéressons aux mesures d'équilibre, c'est-à-dire aux distributions de charges qui minimisent l'énergie électrostatique du potentiel correspondant, parmi toutes les distributions possibles dont la charge totale est fixée. Nous prouvons l'existence de mesures d'équilibre et nous montrons que le potentiel correspondant à l'équilibre est unique et constant dans $\bar{\Omega}$. De plus, pour les domaines réguliers, nous obtenons l'unicité de la mesure d'équilibre, nous donnons son expression précise et nous vérifions que le potentiel d'équilibre est une solution du problème aux limites (\mathcal{BI}) . Finalement, nous caractérisons les boules dans \mathbb{R}^N comme étant les uniques domaines bornés Ω de classe $C^{2,\alpha}$ pour lesquels la distribution d'équilibre est un multiple de la mesure de surface sur $\partial\Omega$. Les mêmes résultats sont obtenus aussi pour des approximations de Taylor de l'énergie électrostatique.

2010 *Mathematics Subject Classification.* 35J93, 35Q60, 78A30.

Key words and phrases. Born-Infeld model, nonlinear electromagnetism, equilibrium measure, equilibrium potential.

CONTENTS

1. Introduction and main results	2
2. Equilibrium measures and equilibrium potentials in electrostatic theories	4
3. Preliminary results	7
4. Equilibrium measures and equilibrium potentials: existence and properties	10
5. The approximated problem	18
6. Characterization of balls via equilibrium measures	22
Acknowledgements	24
References	24

1. INTRODUCTION AND MAIN RESULTS

In this paper we consider the electrostatic Born-Infeld model. Let ρ be a charge distribution on the boundary of a bounded domain $\Omega \subset \mathbb{R}^N$, with $N \geq 3$ ($N = 3$ being the physically relevant case), i.e., we consider ρ as an element of the set $P(\partial\Omega)$ of all positive Borel measures on \mathbb{R}^N supported on $\partial\Omega$ with $\int_{\partial\Omega} d\rho = 1$. We consider the *electrostatic potential* generated by ρ as the unique minimizer ϕ_ρ of the Born-Infeld electrostatic action

$$\mathcal{I}(\phi) = \int_{\mathbb{R}^N} \left(b^2 - b\sqrt{b^2 - |\nabla\phi|^2} \right) dx - \int_{\partial\Omega} \phi d\rho$$

where ϕ runs through the set \mathcal{X} of all Lipschitz functions with Lipschitz-constant less or equal $b > 0$ and $\int_{\mathbb{R}^N} |\nabla\phi|^2 dx < \infty$. One of the main results of [7] shows that \mathcal{I} has a unique minimizer in \mathcal{X} ; further uniqueness questions were addressed in [26]. For each electrostatic potential ϕ_ρ we can consider the Born-Infeld electrostatic energy \mathcal{E} (details of the definition are given in Section 2)

$$\mathcal{E}(\phi_\rho) := -\mathcal{I}(\phi_\rho).$$

Among all possible charge distributions $\rho \in P(\partial\Omega)$ one can search for those distributions, which create least-energy potentials. Such a distribution $\rho^* \in P(\partial\Omega)$ is called *equilibrium distribution* and is defined as follows:

$$\mathcal{E}(\phi_{\rho^*}) = \min_{\rho \in P(\partial\Omega)} \mathcal{E}(\phi_\rho).$$

The corresponding minimizer ϕ_{ρ^*} is called an *equilibrium potential*. The main purpose of this paper is the study of the existence and the properties of ρ^* .

Our first set of results is as follows. We prove that

- (i) equilibrium distributions exist, cf. Theorem 4.1;
- (ii) the equilibrium potential ϕ_{ρ^*} is unique and takes a constant value λ^* in $\bar{\Omega}$, cf. Theorem 4.4 and Corollary 4.5.

An open question in the electrostatic Born-Infeld theory is the following: is it true or not that electrostatic potentials, i.e., minimizers of \mathcal{I} on \mathcal{X} , are weak solutions of the electrostatic Born-Infeld equations

$$(BI) \quad \begin{cases} -\operatorname{div} \left(\frac{b\nabla\phi}{\sqrt{b^2 - |\nabla\phi|^2}} \right) = \rho & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

The answer in general seems not to be known. It is however true when Ω is a ball and ρ a constant multiple of the surface-measure on $\partial\Omega$, cf. [7]. In the case where $\partial\Omega \in C^{2,\alpha}$ we can say more on the equilibrium potential. This is our second set of results. We prove that

- (iii) ϕ_{ρ^*} is a weak solution of (\mathcal{BI}) which is smooth outside $\partial\Omega$, cf. Theorem 4.7;
- (iv) the (unique) value λ^* has a unique characterization given in Proposition 4.11: in particular

$$d\rho^* = -b \frac{\partial_\nu \phi_{\rho^*}}{\sqrt{b^2 - |\nabla \phi_{\rho^*}|^2}} d\sigma,$$

where $\partial_\nu \phi_{\rho^*}$ denotes the outer normal derivative of ϕ_{ρ^*} w.r.t. Ω ;

- (v) the equilibrium distribution ρ^* is unique, cf. Corollary 4.12.

As a third set of results we consider in Section 5 a regularized Born-Infeld model, where the action is given by

$$(1.1) \quad \mathcal{I}^n(\phi) = \sum_{h=1}^n \frac{\alpha_h}{2h} \|\nabla \phi\|_{2h}^{2h} - \oint_{\partial\Omega} \phi d\rho, \quad \alpha_1 = 1, \alpha_h = \frac{1 \cdot 3 \cdot 5 \cdots (2h-3)}{2^{h-1}(h-1)!} b^{2(1-h)}.$$

This arises from the Taylor expansion formula $b^2 - b\sqrt{b^2 - x^2} = \sum_{h=1}^{\infty} \frac{\alpha_h}{2h} x^{2h}$ which converges for $x \in [-b, b]$. As before the corresponding electrostatic energy is given by $\mathcal{E}^n(\phi) := -\mathcal{I}^n(\phi)$. One can define the *electrostatic potential* ϕ_ρ^n , the *equilibrium distribution* $\rho^{*,n}$ and the *equilibrium potential* $\phi_{\rho^{*,n}}^n$ within the approximated model as before based on the notion of action and energy. Our results for this approximated model are as follows:

- (vi) there exist equilibrium distributions $\rho^{*,n}$ and the associated equilibrium potential $\phi_{\rho^{*,n}}^n$ is unique and constant on $\bar{\Omega}$; if, moreover, $\partial\Omega$ is $C^{1,\alpha}$ then $\rho^{*,n}$ is unique, cf. Theorem 5.4;
- (vii) the weak limit $\bar{\rho}$ of $\rho^{*,n}$ as $n \rightarrow \infty$ is an equilibrium distribution for the Born-Infeld electrostatic model. In case $\partial\Omega \in C^{2,\alpha}$ it coincides with ρ^* , i.e., the unique equilibrium distribution of the Born-Infeld model, cf. Proposition 5.9.

In the final Section 6 we characterize balls in \mathbb{R}^N in the following way:

- (viii) for the Born-Infeld model, the ball is the unique set among all bounded $C^{2,\alpha}$ -domains Ω for which the equilibrium distribution ρ^* is a constant multiple of the surface measure $d\sigma$ on $\partial\Omega$;
- (ix) the same holds for the approximated model if $\rho^{*,n}$ is a constant multiple of $d\sigma$ on $\partial\Omega$.

A regularity result for the electrostatic Born-Infeld equation can be derived from the milestone paper of Bartnik and Simon [2]. The result was extended to more general data by Bonheure and Iacopetti [6] but a sharp result is not known. When ρ is a finite sum of point measures, we mention Kiessling's work [27, 28]. The electrostatic Born-Infeld problem has many similarities with the mean curvature problem for graphs in Minkowski space. A variational approach to this geometric problem is given in [3]. Essential properties of the constant mean curvature problem were already considered in fundamental work by Calabi [13], Cheng and Yau [15], Treibergs [41], Bartnik and Simon [2], Gerhardts [23], Ecker [17], Klyachin and Miklyukov [29], and Carley and Kiessling [14]. We refer to [32] for more insight on the geometry of Lorentz-Minkowski spaces and we also mention the recent contribution [5] concerning spacelike radial graphs.

To the best of our knowledge results on the Born-Infeld electrodynamic theory are relatively sparse in the mathematical literature. Well-posedness and stability issues are addressed in [40]. In [37] the interaction of two circularly polarized waves was shown to fulfill the Born-Infeld model in vacuum. In [43] one can find a discussion of the electro- and magnetostatic cases without sources. Special explicit solutions of the dynamical Born-Infeld model, e.g., propagating and counter-propagating plane waves, as well as connections to minimal surfaces can be found in [33]. We also mention [12, 38] in the context of conservation laws.

Finally we point out that the characterization of balls by their electrostatic properties in the linear Maxwell-Poisson context was achieved in [34], [35] for dimensions $N \geq 2$, in [16], [20] for $N = 2$, and in [21] for the p -Laplacian context.

2. EQUILIBRIUM MEASURES AND EQUILIBRIUM POTENTIALS IN ELECTROSTATIC THEORIES

Let us consider an electromagnetic field (\mathbf{E}, \mathbf{B}) in \mathbb{R}^3 described by a gauge potential (ϕ, \mathbf{A}) as follows:

$$\mathbf{E} = -\nabla\phi - \partial_t\mathbf{A}, \quad \mathbf{B} = \nabla \times \mathbf{A}.$$

In the Lagrangian formulation of electromagnetic theories like Maxwell, Born, Born-Infeld the evolution of the fields (\mathbf{E}, \mathbf{B}) is described by the principle of stationary action, i.e., the equations of motion are the Euler-Lagrange equations of the action functional

$$\int_{\mathbb{R}^4} \mathcal{L}(\phi, \mathbf{A}, \nabla\phi, \nabla\mathbf{A}, \partial_t\phi, \partial_t\mathbf{A}) d(x, t).$$

Here \mathcal{L} is called the Lagrangian density. Let us recall the following definition of the energy of an electromagnetic field (\mathbf{E}, \mathbf{B}) (see e.g. [19, 22]). In the following we write $\langle \cdot, \cdot \rangle$ to denote the dual pairing between a space and its dual space.

Definition 2.1. *Let $\mathbf{V} = (\phi, \mathbf{A})$ and suppose that $\mathcal{L} = \mathcal{L}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V})$ is the Lagrangian density of the electromagnetic field (\mathbf{E}, \mathbf{B}) . The Lagrangian at time t is defined by*

$$\mathcal{I}(\mathbf{V}) := \int_{\mathbb{R}^3} \mathcal{L}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V}) dx$$

and the energy at time t is defined as

$$\mathcal{E}(\mathbf{V}) := \int_{\mathbb{R}^3} T^{00}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V}) dx,$$

where

$$T^{00}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V}) = \frac{\partial\mathcal{L}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V})}{\partial(\partial_t\mathbf{V})} \cdot \partial_t\mathbf{V} - \mathcal{L}(\mathbf{V}, \nabla\mathbf{V}, \partial_t\mathbf{V}).$$

In the electrostatic case $\mathbf{V} = (\phi, 0)$ with ϕ independent of t , the Lagrangian becomes

$$\mathcal{I}(\phi) = \int_{\mathbb{R}^3} \mathcal{L}(\phi, \nabla\phi) dx$$

and the electrostatic energy is given by

$$(2.1) \quad \mathcal{E}(\phi) = -\mathcal{I}(\phi) = - \int_{\mathbb{R}^3} \mathcal{L}(\phi, \nabla\phi) dx.$$

In the classical Maxwell theory, the Lagrangian density is given by

$$\mathcal{L}_M = \frac{1}{2}(|\mathbf{E}|^2 - |\mathbf{B}|^2) - \rho\phi + \mathbf{J} \cdot \mathbf{A},$$

where ρ is a charge distribution and \mathbf{J} is a current density. According to (2.1) the electrostatic Maxwell energy reads

$$\mathcal{E}_M(\phi) = -\mathcal{I}_M(\phi) = -\frac{1}{2} \int_{\mathbb{R}^3} |\nabla\phi|^2 dx + \langle \rho, \phi \rangle.$$

The notation $\langle \rho, \phi \rangle$ stands for the duality bracket between a finite Borel measure ρ on \mathbb{R}^3 and a continuous function. The electrostatic potential generated by a given distribution ρ in absence of currents arises as the unique solution of the Euler-Lagrange equation corresponding to the Lagrangian \mathcal{I}_M , i.e. the unique solution of the Poisson equation

$$(2.2) \quad -\Delta\phi = \rho \quad \text{in } \mathbb{R}^3, \quad \lim_{|x| \rightarrow \infty} \phi(x) = 0.$$

For a positive Borel measure ρ with compact support and with finite total mass, the distributional solution to (2.2) is given by the (Newtonian) potential (see [42, Definition 2.1, Theorem 4.1])

$$(2.3) \quad \phi_\rho(x) = \frac{1}{4\pi} \int_{\text{supp } \rho} \frac{1}{|x-y|} d\rho(y).$$

In general, this is not a finite energy solution. For instance, it is well known that if $\rho = \delta_0$ (the Dirac-delta measure at 0), then $\phi_\rho(x) = \frac{1}{4\pi|x|}$ does not have finite energy. Notice that, whenever ϕ_ρ from (2.3) has finite energy, we have

$$\mathcal{E}_M(\phi_\rho) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla \phi_\rho|^2 dx = \frac{1}{8\pi} \int_{\text{supp } \rho} \int_{\text{supp } \rho} \frac{1}{|x-y|} d\rho(y) d\rho(x).$$

Given $\Omega \subset \mathbb{R}^3$, we denote by $P(\partial\Omega)$ the set of probability measures on $\partial\Omega$, i.e., the set of positive Borel measures ρ on \mathbb{R}^3 which are supported on $\partial\Omega$ and normalized by $\int_{\partial\Omega} d\rho = 1$. For every $\rho \in P(\partial\Omega)$, let ϕ_ρ be the corresponding solution of (2.2) given by (2.3). We can now define the notion of an *equilibrium measure* in the Maxwell potential theory.

Definition 2.2. *A probability measure $\rho^* \in P(\partial\Omega)$ is called an equilibrium measure or equilibrium distribution for \mathcal{E}_M if*

$$\mathcal{E}_M(\phi_{\rho^*}) = \inf_{\rho \in P(\partial\Omega)} \mathcal{E}_M(\phi_\rho) = \inf_{\rho \in P(\partial\Omega)} \frac{1}{8\pi} \int_{\partial\Omega} \int_{\partial\Omega} \frac{1}{|x-y|} d\rho(y) d\rho(x).$$

The corresponding solution ϕ_{ρ^} of (2.2) given by (2.3) is called equilibrium potential.*

Notice that the expressions for \mathcal{E}_M and the double integral may be infinite for some measures $\rho \in P(\partial\Omega)$. The infimum, however, is always attained by a unique equilibrium measure and the equilibrium potential is constant on $\bar{\Omega}$, except for a subset of capacity zero, cf. [42, Theorem 6.3, Theorem 7.1, and Theorem 9.1].

For a more recent literature about the traditional electrostatic equilibrium problem using the Coulomb potential of electrostatics we refer to [30, 36].

Let us now consider the Lagrangian density introduced by Born and the modified one by Born and Infeld, namely

$$\mathcal{L}_B = b^2 \left(1 - \sqrt{1 - \frac{|\mathbf{E}|^2 - |\mathbf{B}|^2}{b^2}} \right) - \rho\phi + \mathbf{J} \cdot \mathbf{A}$$

and

$$\mathcal{L}_{BI} = b^2 \left(1 - \sqrt{1 - \frac{|\mathbf{E}|^2 - |\mathbf{B}|^2}{b^2} - \frac{(\mathbf{E} \cdot \mathbf{B})^2}{b^4}} \right) - \rho\phi + \mathbf{J} \cdot \mathbf{A}$$

where $b > 0$ is the Born *field strength parameter* (see [8–11]). In the electrostatic case, Born and Born-Infeld Lagrangian densities lead by (2.1) to the same energy

$$\mathcal{E}(\phi) = -\mathcal{I}(\phi) = -b^2 \int_{\mathbb{R}^3} \left(1 - \sqrt{1 - |\nabla\phi|^2/b^2} \right) dx + \langle \rho, \phi \rangle.$$

In the sequel we set $b = 1$ since the exact value of b has no qualitative influence on our analysis.

One can generalize the functionals \mathcal{E}, \mathcal{I} to \mathbb{R}^N with $N \geq 3$ and $P(\partial\Omega)$ will denote Borel probability measures on \mathbb{R}^N supported on $\partial\Omega$ for bounded domains $\Omega \subset \mathbb{R}^N$. Then, at least formally, the Euler-Lagrange equation for the Lagrangian \mathcal{I} is given by

$$(BI) \quad \begin{cases} -\operatorname{div} \left(\frac{\nabla\phi}{\sqrt{1 - |\nabla\phi|^2}} \right) = \rho & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

As shown in [7] for a large class of measures, finite energy solutions of (\mathcal{BI}) can be found by minimizing the functional \mathcal{I} on the set

$$\mathcal{X} = D^{1,2}(\mathbb{R}^N) \cap \{\phi \in C^{0,1}(\mathbb{R}^N) : \|\nabla\phi\|_\infty \leq 1\},$$

which is a closed and convex subset of the space $D^{1,2}(\mathbb{R}^N)$.

In particular, following the definition of a weak solution [7, Definition 1.2], we have that a weak solution of (\mathcal{BI}) must minimize \mathcal{I} on \mathcal{X} , cf. [7, Proposition 2.6].

For any $\rho \in P(\partial\Omega)$, we have that $\rho \in \mathcal{X}^*$ and by [7, Theorem 1.3] there exists a unique $\phi_\rho \in \mathcal{X}$ which minimizes \mathcal{I} (some details on the properties of \mathcal{X}^* are given in Section 3 after Lemma 3.2). Notice that, by [7, Lemma 2.12], $\phi_\rho \geq 0$. Unfortunately, as far as we are aware, it is not known whether in general the minimizer ϕ_ρ is a solution of (\mathcal{BI}) .¹ Thus, dealing with the Born-Infeld Lagrangian, we have to define equilibrium measures through the minimizer ϕ_ρ of \mathcal{I} rather than through the solution of (\mathcal{BI}) .

Definition 2.3. *A probability measure $\rho^* \in P(\partial\Omega)$ is called an equilibrium measure or equilibrium distribution for \mathcal{E} if*

$$\mathcal{E}(\phi_{\rho^*}) = \inf_{\rho \in P(\partial\Omega)} \mathcal{E}(\phi_\rho),$$

where for every $\rho \in P(\partial\Omega)$ we denote by ϕ_ρ the unique minimizer of \mathcal{I} on \mathcal{X} . The corresponding minimizer ϕ_{ρ^*} of \mathcal{I} is called equilibrium potential.

Remark 2.4. *Another way to obtain an expression for the energy is to move from the Lagrangian (expressed in terms of $\nabla\phi$) to the Hamiltonian (expressed in terms of the dual variable ρ) via the Legendre transform. Formally, this yields in the Maxwell case*

$$\begin{aligned} \mathcal{H}_M(\rho) &= \sup_{\phi} \left(\langle \rho, \phi \rangle - \frac{1}{2} \int_{\mathbb{R}^3} |\nabla\phi|^2 dx \right) \\ &= \langle \rho, \phi_\rho \rangle - \frac{1}{2} \int_{\mathbb{R}^3} |\nabla\phi_\rho|^2 dx \\ &= \frac{1}{2} \int_{\mathbb{R}^3} |\nabla\phi_\rho|^2 dx \end{aligned}$$

since ϕ_ρ satisfies (2.2). Similarly, in the Born-Infeld case one finds

$$\begin{aligned} \mathcal{H}(\rho) &= \sup_{\phi} \left(\langle \rho, \phi \rangle - \frac{1}{2} \int_{\mathbb{R}^3} 1 - \sqrt{1 - |\nabla\phi|^2} dx \right) \\ &= \langle \rho, \phi_\rho \rangle - \frac{1}{2} \int_{\mathbb{R}^3} 1 - \sqrt{1 - |\nabla\phi_\rho|^2} dx \\ &= -\mathcal{I}(\phi_\rho). \end{aligned}$$

In both cases the Hamiltonian of a measure ρ is the same as the energy of ϕ_ρ . If ϕ_ρ satisfies (\mathcal{BI}) more can be said: as a consequence of the definition of a weak solution of (\mathcal{BI}) , see [7, Definition 1.2], we have, in particular, that a weak solution $\phi_\rho \in \mathcal{X}$ satisfies

$$(2.4) \quad \int_{\mathbb{R}^N} \frac{|\nabla\phi_\rho|^2}{\sqrt{1 - |\nabla\phi_\rho|^2}} dx = \langle \rho, \phi_\rho \rangle,$$

¹Known cases, where ϕ_ρ solves (\mathcal{BI}) , include $\rho \in L^\infty(\mathbb{R}^N) \cap \mathcal{X}^*$ and ρ being a radially distributed charge density or a finite sum of point charges, cf. [7] and [27, 28].

and so, in this case, the Hamiltonian of the measure ρ (and hence the energy of the solution ϕ_ρ) is

$$\begin{aligned}\mathcal{E}(\phi_\rho) &= \mathcal{H}(\rho) = \int_{\mathbb{R}^N} \left(-1 + \sqrt{1 - |\nabla\phi_\rho|^2} + \frac{|\nabla\phi_\rho|^2}{\sqrt{1 - |\nabla\phi_\rho|^2}} \right) dx \\ &= \int_{\mathbb{R}^N} \left(\frac{1}{\sqrt{1 - |\nabla\phi_\rho|^2}} - 1 \right) dx =: \mathcal{K}(\phi_\rho).\end{aligned}$$

In general however, we only know that a minimizer $\phi_\rho \in \mathcal{X}$ of \mathcal{I} satisfies (2.4) with “ \leq ” instead of “ $=$ ”, and thus we only have $\mathcal{E}(\phi_\rho) = \mathcal{H}(\rho) \geq \mathcal{K}(\phi_\rho)$.

3. PRELIMINARY RESULTS

Here we provide the functional analytic setting for the minimization of the Lagrangian. We prove in Lemma 3.7 that the minimizer ϕ_ρ depends continuously on the measure ρ . Thereafter we provide several tools which allow to deduce that ϕ_ρ attains its maximum on $\partial\Omega$, cf. Lemma 3.11.

Let $\rho \in P(\partial\Omega)$. In the following we add a subscript to any functional that depends on the choice of the measure ρ in its definition to emphasize this dependence and to keep the notation closer to [7]. For instance,

$$\mathcal{I}_\rho(\phi) := \int_{\mathbb{R}^N} \left(1 - \sqrt{1 - |\nabla\phi|^2} \right) dx - \langle \rho, \phi \rangle, \quad \phi \in \mathcal{X}.$$

We also define the functional $\mathcal{J} : \mathcal{X} \rightarrow \mathbb{R}$ that does not depend on the measure ρ by

$$\mathcal{J}(\phi) := \int_{\mathbb{R}^N} \left(1 - \sqrt{1 - |\nabla\phi|^2} \right) dx, \quad \phi \in \mathcal{X}.$$

Observe that, since $\|\nabla\phi\|_\infty \leq 1$, we can write \mathcal{J} as the infinite series

$$(3.1) \quad \mathcal{J}(\phi) = \sum_{h=1}^{+\infty} \frac{\alpha_h}{2h} \|\nabla\phi\|_{2h}^{2h}.$$

The exact values of the coefficients α_h are given in (1.1). Next we recall some fundamental properties of the set \mathcal{X} ([7, Lemma 2.1]).

Lemma 3.1. *The following assertions hold:*

- (i) \mathcal{X} is continuously embedded in $W^{1,p}(\mathbb{R}^N)$ for all $p \geq 2^* = 2N/(N-2)$;
- (ii) \mathcal{X} is continuously embedded in $L^\infty(\mathbb{R}^N)$;
- (iii) if $\phi \in \mathcal{X}$, then $\lim_{|x| \rightarrow \infty} \phi(x) = 0$;
- (iv) \mathcal{X} is weakly closed;
- (v) if $(\phi_n)_n \subset \mathcal{X}$ is bounded, there exists $\bar{\phi} \in \mathcal{X}$ such that, up to a subsequence, $\phi_n \rightharpoonup \bar{\phi}$ weakly in $D^{1,2}(\mathbb{R}^N)$ and uniformly on compact sets.

Next we give a more detailed version of (ii) from the previous lemma.

Lemma 3.2. *For any $t > \max\{N, \frac{2N}{N-2}\}$ there exists a constant $C(t)$ such that, for all $\phi \in \mathcal{X}$,*

$$\|\phi\|_\infty \leq C(t) \|\nabla\phi\|_2^{\frac{2}{t}} (1 + \|\nabla\phi\|_2^{\frac{2}{N}}).$$

Proof. With t as in the assumption, define $s = \frac{tN}{N+t}$. Then $s < N$ and $t = s^* = \frac{Ns}{N-s}$. Since $t > 2^* = \frac{2N}{N-2}$, we deduce that $s > 2$. Recall also that, by (i) and (ii) in Lemma 3.1, \mathcal{X} embeds into $W^{1,t}(\mathbb{R}^N)$ and therefore into $L^\infty(\mathbb{R}^N)$. Hence, for every $\phi \in \mathcal{X}$, using Sobolev inequality with $t = s^*$ and the fact that $|\nabla\phi| \leq 1$, we get

$$\|\phi\|_\infty \leq C(t) (\|\nabla\phi\|_t + \|\phi\|_t) \leq \bar{C}(t) (\|\nabla\phi\|_t + \|\nabla\phi\|_s) \leq \bar{C}(t) (\|\nabla\phi\|_2^{\frac{2}{t}} + \|\nabla\phi\|_2^{\frac{2}{s}})$$

since $s, t > 2$. This proves the claim because $\frac{1}{s} = \frac{1}{t} + \frac{1}{N}$. \square

As mentioned in the introduction, one of the main results of [7, Theorem 1.3] states that for every $\rho \in \mathcal{X}^*$ there exists a unique minimizer $\phi_\rho \in \mathcal{X}$ of \mathcal{I}_ρ . We denote by \mathcal{X}^* the dual space of $D^{1,2}(\mathbb{R}^N) \cap C^{0,1}(\mathbb{R}^N)$ endowed with the norm $\|\nabla \cdot\|_{L^2} + \|\nabla \cdot\|_{L^\infty}$. Note that $\mathcal{X}^* \supsetneq D^{1,2}(\mathbb{R}^N)^*$ since, e.g., finite Borel measures on \mathbb{R}^N are included in \mathcal{X}^* . Moreover, cf. [7, Theorem 1.5], if $\rho \in L^\infty(\mathbb{R}^N)$ then this unique minimizer is a *strictly spacelike, weak solution* in the sense given in the following two definitions.

Definition 3.3 (cf. [2]). *Let $\Omega \subset \mathbb{R}^N$ be open. A function $\phi \in C^{0,1}(\Omega)$ is called strictly spacelike if $\phi \in C^1(\Omega)$ and $|\nabla\phi(x)| < 1$ for all $x \in \Omega$.*

Definition 3.4 (cf. [7]). *Let $\rho \in \mathcal{X}^*$. A function $\phi \in \mathcal{X}$ is called a weak solution of (BT) if*

$$(3.2) \quad \int_{\mathbb{R}^N} \frac{\nabla\phi \cdot \nabla\psi}{\sqrt{1 - |\nabla\phi|^2}} dx = \langle \rho, \psi \rangle \text{ for all } \psi \in C_c^\infty(\mathbb{R}^N).$$

Remark 3.5. *Note that the requirement that (3.2) holds for all $\psi \in C_c^\infty(\mathbb{R}^N)$ is equivalent to asking that (3.2) holds for all $\psi \in \mathcal{X}$, which is the original definition given in [7].*

Lemma 3.6. *There exists a constant $C > 0$ such that*

$$\|\nabla\phi_\rho\|_2 \leq C$$

for every $\rho \in P(\partial\Omega)$.

Proof. Since $\mathcal{I}_\rho(0) = 0$ and $\frac{1}{2}\tau \leq 1 - \sqrt{1 - \tau}$, for $\tau \in [0, 1]$, we have

$$0 \geq \mathcal{I}_\rho(\phi_\rho) \geq \frac{1}{2}\|\nabla\phi_\rho\|_2^2 - \|\phi_\rho\|_{L^\infty(\partial\Omega)}$$

and so, by Lemma 3.2,

$$\|\nabla\phi_\rho\|_2^2 \leq 2C(t)\|\nabla\phi_\rho\|_2^{\frac{2}{t}}(1 + \|\nabla\phi_\rho\|_2^{\frac{2}{N}})$$

with $t > \max\{N, \frac{2N}{N-2}\}$. Thus, since $\frac{2}{t} < 1$ and $\frac{2}{t} + \frac{2}{N} < 1$, we get the claim. \square

Lemma 3.7 (Continuous dependence of ϕ_ρ on ρ). *If $\rho_k, \rho \in P(\partial\Omega)$ with $\rho_k \rightharpoonup \rho$ as $k \rightarrow \infty$ weakly in the sense of measures, then*

$$(3.3) \quad \lim_{k \rightarrow \infty} \mathcal{I}_{\rho_k}(\phi_{\rho_k}) = \mathcal{I}_\rho(\phi_\rho)$$

and $\phi_{\rho_k} \rightarrow \phi_\rho$ strongly in $D^{1,2}(\mathbb{R}^N)$ and locally uniformly on \mathbb{R}^N as $k \rightarrow \infty$.

Proof. The boundedness of $\{\phi_{\rho_k}\}_{k \in \mathbb{N}}$ (see Lemma 3.6) and Lemma 3.1 imply the existence of a $\phi \in \mathcal{X}$ such that $\phi_{\rho_k} \rightharpoonup \phi$ weakly in $D^{1,2}(\mathbb{R}^N)$ and uniformly on compact sets as $k \rightarrow \infty$. Then, since \mathcal{J} is convex and continuous on \mathcal{X} , it is weakly lower semicontinuous and so

$$\mathcal{J}(\phi) \leq \liminf_{k \rightarrow \infty} \mathcal{J}(\phi_{\rho_k}).$$

Moreover, as $k \rightarrow \infty$,

$$(3.4) \quad |\langle \rho_k - \rho, \phi_{\rho_k} \rangle| \leq 2\|\phi_{\rho_k} - \phi\|_{L^\infty(\partial\Omega)} + |\langle \rho_k - \rho, \phi \rangle| \rightarrow 0,$$

$$(3.5) \quad |\langle \rho_k, \phi_{\rho_k} \rangle - \langle \rho, \phi \rangle| \leq |\langle \rho_k - \rho, \phi \rangle| + \|\phi_{\rho_k} - \phi\|_{L^\infty(\partial\Omega)} \rightarrow 0.$$

Thus, using (3.4) and the fact that ϕ_{ρ_k} minimizes \mathcal{I}_{ρ_k} , we have

$$\begin{aligned} \mathcal{I}_\rho(\phi) &\leq \liminf_{k \rightarrow \infty} \mathcal{I}_\rho(\phi_{\rho_k}) = \liminf_{k \rightarrow \infty} [\mathcal{I}_{\rho_k}(\phi_{\rho_k}) + \langle \rho_k - \rho, \phi_{\rho_k} \rangle] = \liminf_{k \rightarrow \infty} \mathcal{I}_{\rho_k}(\phi_{\rho_k}) \\ &\leq \limsup_{k \rightarrow \infty} \mathcal{I}_{\rho_k}(\phi_{\rho_k}) \leq \limsup_{k \rightarrow \infty} \mathcal{I}_{\rho_k}(\phi_\rho) = \lim_{k \rightarrow \infty} \mathcal{I}_{\rho_k}(\phi_\rho) = \mathcal{I}_\rho(\phi_\rho) \end{aligned}$$

that implies, by the uniqueness of the minimum, $\phi = \phi_\rho$ and (3.3) holds. In particular, using (3.5), we have that

$$(3.6) \quad \lim_{k \rightarrow \infty} \mathcal{J}(\phi_{\rho_k}) = \mathcal{J}(\phi_\rho).$$

On the other hand, by Clarkson's inequality [25], we have that

$$\left\| \frac{\nabla \phi_{\rho_k} - \nabla \phi_\rho}{2} \right\|_{2h}^{2h} + \left\| \frac{\nabla \phi_{\rho_k} + \nabla \phi_\rho}{2} \right\|_{2h}^{2h} \leq \frac{1}{2} (\|\nabla \phi_{\rho_k}\|_{2h}^{2h} + \|\nabla \phi_\rho\|_{2h}^{2h}), \quad h \in \mathbb{N},$$

and so

$$\sum_{h=1}^{+\infty} \frac{\alpha_h}{2h} \left\| \frac{\nabla \phi_{\rho_k} - \nabla \phi_\rho}{2} \right\|_{2h}^{2h} + \sum_{h=1}^{+\infty} \frac{\alpha_h}{2h} \left\| \frac{\nabla \phi_{\rho_k} + \nabla \phi_\rho}{2} \right\|_{2h}^{2h} \leq \frac{1}{2} \sum_{h=1}^{+\infty} \frac{\alpha_h}{2h} (\|\nabla \phi_{\rho_k}\|_{2h}^{2h} + \|\nabla \phi_\rho\|_{2h}^{2h}),$$

namely

$$\mathcal{J} \left(\frac{\phi_{\rho_k} - \phi_\rho}{2} \right) + \mathcal{J} \left(\frac{\phi_{\rho_k} + \phi_\rho}{2} \right) \leq \frac{1}{2} (\mathcal{J}(\phi_{\rho_k}) + \mathcal{J}(\phi_\rho)).$$

By the weak lower semicontinuity of \mathcal{J} we see that

$$\mathcal{J}(\phi_\rho) \leq \liminf_{k \rightarrow \infty} \mathcal{J} \left(\frac{\phi_{\rho_k} + \phi_\rho}{2} \right).$$

Therefore, using (3.6), we obtain

$$\begin{aligned} 0 &\leq \liminf_{k \rightarrow \infty} \mathcal{J} \left(\frac{\phi_{\rho_k} - \phi_\rho}{2} \right) \leq \limsup_{k \rightarrow \infty} \mathcal{J} \left(\frac{\phi_{\rho_k} - \phi_\rho}{2} \right) \\ &\leq \limsup_{k \rightarrow \infty} \left[\frac{1}{2} (\mathcal{J}(\phi_{\rho_k}) + \mathcal{J}(\phi_\rho)) - \mathcal{J} \left(\frac{\phi_{\rho_k} + \phi_\rho}{2} \right) \right] \\ &= \mathcal{J}(\phi_\rho) - \liminf_{k \rightarrow \infty} \mathcal{J} \left(\frac{\phi_{\rho_k} + \phi_\rho}{2} \right) \leq 0 \end{aligned}$$

and thus

$$(3.7) \quad \lim_{k \rightarrow \infty} \mathcal{J} \left(\frac{\phi_{\rho_k} - \phi_\rho}{2} \right) = 0.$$

Hence (3.7) allows us to conclude since, by the inequality $\frac{1}{2}\tau \leq 1 - \sqrt{1 - \tau}$, for $\tau \in [0, 1]$, we have that for all $\phi \in \mathcal{X}$, $\|\nabla \phi\|_2^2 \leq 2\mathcal{J}(\phi)$. \square

Lemma 3.8. *Let $\rho \in L^\infty(\mathbb{R}^N)$ have compact support and let $\phi_\rho \in \mathcal{X}$ be the minimizer of \mathcal{I}_ρ . Suppose that $0 < \max_{\mathbb{R}^N} \phi_\rho = \phi_\rho(x_0)$ for some point $x_0 \in \mathbb{R}^N$. Then x_0 lies in $\text{supp } \rho$ or x_0 lies in a bounded connected component of $\mathbb{R}^N \setminus \text{supp } \rho$ where $\phi_\rho \equiv \phi_\rho(x_0)$. In any case*

$$(3.8) \quad \text{supp } \rho \cap \{x \in \mathbb{R}^N : \phi_\rho(x) = \max_{\mathbb{R}^N} \phi_\rho\} \neq \emptyset.$$

Proof. Since $\rho \in L^\infty(\mathbb{R}^N)$ has compact support, we see that $\rho \in \mathcal{X}^*$. Therefore [7, Theorem 1.5] implies that ϕ_ρ is a weak, strictly spacelike solution of (\mathcal{BI}) , i.e. $\phi_\rho \in C^1(\mathbb{R}^N) \cap \mathcal{X}$, $|\nabla \phi_\rho| < 1$ in \mathbb{R}^N and (3.2) holds.

Suppose $x_0 \notin \text{supp } \rho$. Then there exists an open ball B centered at x_0 such that $\bar{B} \cap \text{supp } \rho = \emptyset$ and

$$(3.9) \quad \int_B a(x) \nabla \phi_\rho \cdot \nabla \psi \, dx = 0 \text{ for all } \psi \in C_c^\infty(B)$$

where $a(x) = (1 - |\nabla \phi_\rho(x)|^2)^{-1/2} \geq \delta > 0$ in B . Since ϕ_ρ attains its maximum on \bar{B} at $x_0 \in B$, the strong maximum principle (see e.g. [24, Theorem 8.19]) applies. We then deduce that $\phi_\rho \equiv$

const. = $\phi_\rho(x_0)$ on \overline{B} and in fact on the entire connected component of $\mathbb{R}^N \setminus \text{supp } \rho$ containing x_0 . This connected component must be bounded since $\phi_\rho(x) \rightarrow 0$ as $|x| \rightarrow \infty$ by (iii) in Lemma 3.1. \square

To develop our theory further we next rely on the idea that one can “mollify” a measure as expressed in the following lemma. The proof is a straightforward consequence of the definition of the mollified measure.

Lemma 3.9. *Let $\mu \in P(\partial\Omega)$, $\varepsilon > 0$ and $\eta_\varepsilon : \mathbb{R}^N \rightarrow [0, \infty)$ be the standard mollifier supported in $\overline{B}_\varepsilon(0)$. Define the function $f_\varepsilon \in L^\infty(\mathbb{R}^N)$ by*

$$f_\varepsilon(x) := \oint_{\partial\Omega} \eta_\varepsilon(y-x) d\mu_y, \quad x \in \mathbb{R}^N,$$

and the measure $\mu_\varepsilon \in P(\mathbb{R}^N)$ by

$$d\mu_\varepsilon := f_\varepsilon dx.$$

Then $\mu_\varepsilon \rightarrow \mu$ as $\varepsilon \rightarrow 0$ in the sense of measures and $\text{supp } \mu_\varepsilon \subset \text{supp } \mu + \overline{B}_\varepsilon(0)$.

We also use the following lemma on the location of maxima of locally uniformly convergent sequences of continuous functions. A proof by contradiction is again straightforward.

Lemma 3.10. *Let $\{\phi_k\}_{k \in \mathbb{N}}$ be a sequence of continuous functions on \mathbb{R}^N that converges locally uniformly to ϕ and suppose that $T := \{x \in \mathbb{R}^N : \phi(x) = \max_{\mathbb{R}^N} \phi\}$ is compact and that for all $k \in \mathbb{N}$ the sets $T_k := \{x \in \mathbb{R}^N : \phi_k(x) = \max_{\mathbb{R}^N} \phi_k\}$ are contained in a fixed compact set K . For any given $\delta > 0$ and sufficiently large k the function ϕ_k attains its maximum only at points in $T + \overline{B}_\delta(0)$.*

As a consequence of the previous lemmas, we can now deduce that for any $\rho \in P(\partial\Omega)$, ϕ_ρ attains its maximum value on the boundary of Ω .

Lemma 3.11. *For any $\rho \in P(\partial\Omega)$, we have that $\max_{\mathbb{R}^n} \phi_\rho = \max_{\partial\Omega} \phi_\rho$.*

Proof. Suppose that $M := \max_{\mathbb{R}^N} \phi_\rho > \max_{\partial\Omega} \phi_\rho$ and define the compact set $T = \{x \in \mathbb{R}^N : \phi_\rho(x) = M\}$. Clearly $T \cap \partial\Omega = \emptyset$ and $\text{dist}(T, \partial\Omega) > 0$. By Lemma 3.9, let $\varepsilon_0, \delta > 0$ be so small that the mollified measure ρ_ε satisfies $\text{dist}(\text{supp } \rho_\varepsilon, T + \overline{B}_\delta(0)) > 0$ for all $\varepsilon \in (0, \varepsilon_0]$. Note that $\text{supp } \rho_\varepsilon \subset \partial\Omega + \overline{B}_\varepsilon(0) \subset \overline{B}_R(0)$ for all $\varepsilon \in (0, \varepsilon_0]$ and a suitable $R > 0$. Therefore, if a point x belongs to a bounded component of $\mathbb{R}^N \setminus \text{supp } \rho_\varepsilon$ then $|x| \leq R$. By Lemma 3.8, applied to ρ_ε and ϕ_{ρ_ε} , we find that ϕ_{ρ_ε} has its maximum points all within the set $\overline{B}_R(0)$ for all $\varepsilon \in (0, \varepsilon_0]$. Since $\phi_{\rho_\varepsilon} \rightarrow \phi_\rho$ as $\varepsilon \rightarrow 0$ locally uniformly on \mathbb{R}^N , by Lemma 3.10 we know that for small $\varepsilon > 0$ the function ϕ_{ρ_ε} attains its maximum over \mathbb{R}^N only at points $y_\varepsilon \in T + \overline{B}_\delta(0)$. However, since $T + \overline{B}_\delta(0)$ and $\text{supp } \rho_\varepsilon$ are disjoint we get a contradiction to (3.8) from Lemma 3.8. This contradiction finishes the proof. \square

4. EQUILIBRIUM MEASURES AND EQUILIBRIUM POTENTIALS: EXISTENCE AND PROPERTIES

In this section, we prove our main results. We begin with existence of equilibrium measures (Theorem 4.1), their properties (Theorem 4.4) and the uniqueness of equilibrium potentials (Corollary 4.5). Then, in case of $C^{2,\alpha}$ -smooth domains we show that the equilibrium potential solves (\mathcal{BI}) (Theorem 4.7), we give a characterization (Proposition 4.11) and address the uniqueness questions for the equilibrium measure (Corollary 4.12, Proposition 4.13).

Theorem 4.1. *There exists an equilibrium measure ρ^* for \mathcal{E} .*

Proof. Recall from [7, Proposition 2.3] that $\mathcal{I}_\rho(\phi_\rho) = -\mathcal{E}(\phi_\rho) < 0$. Hence we infer that

$$\inf_{\rho \in P(\partial\Omega)} \mathcal{E}(\phi_\rho) \geq 0.$$

Let $\{\rho_k\}_{k \in \mathbb{N}}$ be a minimizing sequence. Since $\partial\Omega$ is compact, the sequence $\{\rho_k\}_{k \in \mathbb{N}}$ is tight and hence there exists $\rho^* \in P(\partial\Omega)$ such that $\rho_k \rightarrow \rho^*$ (see [4]). By Lemma 3.7 we get that

$$\mathcal{E}(\phi_{\rho^*}) = \lim_{k \rightarrow \infty} \mathcal{E}(\phi_{\rho_k}) = \inf_{\rho \in P(\partial\Omega)} \mathcal{E}(\phi_\rho)$$

so that ρ^* is an equilibrium measure for \mathcal{E} . \square

From now on, we denote by ρ^* an equilibrium measure. It satisfies the following properties.

Proposition 4.2. *For every $\mu \in P(\partial\Omega)$ we have the inequality*

$$(4.1) \quad \lambda^* := \langle \rho^*, \phi_{\rho^*} \rangle \leq \langle \mu, \phi_{\rho^*} \rangle.$$

Proof. Let μ be an arbitrary measure in $P(\partial\Omega)$. For $t \in [0, 1]$, define the probability measure $\rho_t := (1-t)\rho^* + t\mu$. Then

$$\mathcal{E}(\phi_{\rho_t}) - \mathcal{E}(\phi_{\rho^*}) = \mathcal{I}_{\rho^*}(\phi_{\rho^*}) - \mathcal{I}_{\rho^*}(\phi_{\rho_t}) + t\langle \mu - \rho^*, \phi_{\rho_t} \rangle \leq t\langle \mu - \rho^*, \phi_{\rho_t} \rangle$$

and

$$\mathcal{E}(\phi_{\rho_t}) - \mathcal{E}(\phi_{\rho^*}) = -\mathcal{I}_{\rho_t}(\phi_{\rho_t}) + \mathcal{I}_{\rho_t}(\phi_{\rho^*}) + t\langle \mu - \rho^*, \phi_{\rho^*} \rangle \geq t\langle \mu - \rho^*, \phi_{\rho^*} \rangle,$$

for $t \in (0, 1]$ imply that

$$\langle \mu - \rho^*, \phi_{\rho^*} \rangle \leq \frac{\mathcal{E}(\phi_{\rho_t}) - \mathcal{E}(\phi_{\rho^*})}{t} \leq \langle \mu - \rho^*, \phi_{\rho_t} \rangle.$$

Passing to the limit as $t \rightarrow 0^+$ and using the Lemma 3.7 we get that

$$\lim_{t \rightarrow 0} \frac{\mathcal{E}(\phi_{\rho_t}) - \mathcal{E}(\phi_{\rho^*})}{t} \text{ exists and } = \langle \mu - \rho^*, \phi_{\rho^*} \rangle.$$

Finally, since $\mathcal{E}(\phi_{\rho_t}) - \mathcal{E}(\phi_{\rho^*}) \geq 0$, we obtain the claimed inequality (4.1). \square

Lemma 4.3. *For every $x \in \partial\Omega$ we have $\phi_{\rho^*}(x) \geq \lambda^*$. Moreover $\phi_{\rho^*} = \lambda^*$ a.e. with respect to ρ^* on $\partial\Omega$.*

Proof. If we take $\mu = \delta_x$ for $x \in \partial\Omega$ and insert this measure into (4.1), we get $\phi_{\rho^*}(x) \geq \lambda^*$. Moreover, by the definition of λ^* we have

$$0 = \int_{\partial\Omega} \underbrace{(\phi_{\rho^*}(x) - \lambda^*)}_{\geq 0} d\rho^*$$

which implies $\phi_{\rho^*} = \lambda^*$ a.e. with respect to the measure ρ^* on $\partial\Omega$. \square

Theorem 4.4. *For every $x \in \bar{\Omega}$ we have $\phi_{\rho^*}(x) = \lambda^*$.*

Proof. We divide the proof in two steps: we first show that the statement is true on $\partial\Omega$ and, then, also in Ω .

STEP 1: for every $x \in \partial\Omega$ we have $\phi_{\rho^*}(x) = \lambda^*$.

Suppose $M := \max_{\partial\Omega} \phi_{\rho^*} > \lambda^*$ and define

$$T = \{x \in \partial\Omega : \phi_{\rho^*}(x) = M\}.$$

Note that $\rho^*(T) = 0$ and, since $M > \lambda^*$, the continuity of ϕ_{ρ^*} implies that even $\rho^*(T + \bar{B}_\delta(0)) = 0$ for all sufficiently small $\delta \in (0, \delta_0]$ by Lemma 4.3.

Let us show that in particular $(T + \bar{B}_{\delta_0/2}(0)) \cap \text{supp } \rho^* = \emptyset$. Suppose for contradiction that $x \in (T + \bar{B}_{\delta_0/2}(0)) \cap \text{supp } \rho^*$. Then for every open neighbourhood N_x of x we have $\rho^*(N_x) > 0$,

and in particular for $N_x = B_{\delta_0/2}(x)$. But $B_{\delta_0/2}(x) \subset T + \overline{B}_{\delta_0}(0)$ and hence $\rho^*(N_x) = 0$ so that we have reached a contradiction. We now have that $\text{dist}(\text{supp } \rho^*, T + \overline{B}_{\delta_0/2}(0)) > 0$.

Let ε_0 be so small that the mollified measure ρ_ε^* (according to Lemma 3.9) satisfies $\text{dist}(\text{supp } \rho_\varepsilon^*, T + \overline{B}_{\delta_0/2}(0)) > 0$ for all $\varepsilon \in (0, \varepsilon_0]$. Arguing as in Lemma 3.11 we get that $\phi_{\rho_\varepsilon^*}$ attains its maximum only at $T + \overline{B}_{\delta_0/2}(0)$ for sufficiently small ε . However, since $T + \overline{B}_{\delta_0/2}(0)$ and $\text{supp } \rho_\varepsilon^*$ are disjoint we get, as before, a contradiction to (3.8) from Lemma 3.8. This finishes the first step.

STEP 2: for every $x \in \Omega$ we have $\phi_{\rho^*}(x) = \lambda^*$.

Let us define the function

$$\hat{\phi}_{\rho^*}(x) = \begin{cases} \phi_{\rho^*}(x), & x \in \mathbb{R}^N \setminus \Omega, \\ \lambda^*, & x \in \Omega. \end{cases}$$

Then $\hat{\phi}_{\rho^*} \in \mathcal{X}$, $\mathcal{J}(\hat{\phi}_{\rho^*}) \leq \mathcal{J}(\phi_{\rho^*})$ and by Step 1

$$\langle \rho^*, \hat{\phi}_{\rho^*} \rangle = \int_{\partial\Omega} \lambda^* d\rho^* = \langle \rho^*, \phi_{\rho^*} \rangle$$

so that finally $\mathcal{I}_{\rho^*}(\hat{\phi}_{\rho^*}) \leq \mathcal{I}_{\rho^*}(\phi_{\rho^*})$. The uniqueness of the minimizer implies $\hat{\phi}_{\rho^*} = \phi_{\rho^*}$ and this finishes the proof. \square

From Theorem 4.4 we obtain the following result, which shows in particular the interesting fact that although in general we do not know about the uniqueness of the *equilibrium measure*, the *equilibrium potential* is always unique.

Corollary 4.5. *For every $\mu \in P(\partial\Omega)$ we have the equality*

$$(4.2) \quad \lambda^* = \langle \rho^*, \phi_{\rho^*} \rangle = \langle \mu, \phi_{\rho^*} \rangle.$$

Moreover, the value λ^* and the equilibrium potential ϕ_{ρ^*} are unique in the following sense: whenever ρ^*, ρ^{**} are equilibrium measures then $\phi_{\rho^*} = \phi_{\rho^{**}}$ and $\lambda^* = \lambda^{**}$.

Proof. The equality (4.2) follows directly from Theorem 4.4. In particular, $\langle \rho^{**} - \rho^*, \phi_{\rho^{**}} \rangle = 0$. Therefore, we have

$$-\mathcal{E}(\phi_{\rho^*}) = \mathcal{I}_{\rho^*}(\phi_{\rho^*}) \leq \mathcal{I}_{\rho^*}(\phi_{\rho^{**}}) = \mathcal{I}_{\rho^{**}}(\phi_{\rho^{**}}) = -\mathcal{E}(\phi_{\rho^{**}}).$$

Since the two energies $\mathcal{E}(\phi_{\rho^*})$, $\mathcal{E}(\phi_{\rho^{**}})$ are equal, we obtain $\mathcal{I}_{\rho^*}(\phi_{\rho^*}) = \mathcal{I}_{\rho^*}(\phi_{\rho^{**}})$. By the uniqueness of the minimizer of \mathcal{I}_{ρ^*} we get $\phi_{\rho^*} = \phi_{\rho^{**}}$ which also implies $\lambda^* = \lambda^{**}$. \square

Lemma 4.6. *The value λ^* is strictly positive.*

Proof. By Theorem 4.4 and Lemma 3.11 we find $\lambda^* = \max_{\partial\Omega} \phi_{\rho^*} = \max_{\mathbb{R}^N} \phi_{\rho^*}$. Since moreover ϕ_{ρ^*} vanishes at infinity, we deduce that $\lambda^* \geq 0$. Suppose for contradiction that $\lambda^* = 0$. Then, by [7, Proposition 2.7],

$$\int_{\mathbb{R}^N} \frac{|\nabla \phi_{\rho^*}|^2}{\sqrt{1 - |\nabla \phi_{\rho^*}|^2}} dx \leq \langle \rho^*, \phi_{\rho^*} \rangle = \lambda^* = 0.$$

Hence $\phi_{\rho^*} = 0$ and so $\mathcal{I}_{\rho^*}(\phi_{\rho^*}) = 0$, in contradiction to $\mathcal{I}_{\rho^*}(\phi_{\rho^*}) < 0$, cf. [7, Proposition 2.3]. \square

For smooth domains, we can give a more precise description of the equilibrium measure and of its properties.

Theorem 4.7. *Suppose that $\partial\Omega \in C^{2,\alpha}$, for some $\alpha > 0$. Then ϕ_{ρ^*} is the unique weak solution of*

$$(4.3) \quad \begin{cases} -\text{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = \rho^* & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

Moreover, for every $r > 0$ sufficiently large, $\phi_{\rho^*} \in C^{2,\alpha}(\overline{B}_r(0) \setminus \Omega)$ and there exists $\theta = \theta(r) \in (0, 1)$ such that $|\nabla \phi_{\rho^*}| \leq 1 - \theta < 1$ in $\overline{B}_r(0)$.

Proof. By Theorem 4.4 we know that $\phi_{\rho^*} = \lambda^*$ in $\overline{\Omega}$. The proof will be divided into two steps. In the first step, we show local regularity and strict positivity. In the second step, boundary regularity is included.

STEP 1 (LOCAL REGULARITY): Let us show that $\phi_{\rho^*} \in C^\infty(\mathbb{R}^N \setminus \overline{\Omega})$, that it is strictly positive and strictly spacelike in $\mathbb{R}^N \setminus \overline{\Omega}$ and satisfies

$$(4.4) \quad \begin{cases} -\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = 0 & \text{in } \mathbb{R}^N \setminus \overline{\Omega}, \\ \phi = \lambda^* & \text{on } \partial\Omega, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

To prove this claim we adapt some ideas of [2, 7] to our case. Let \mathcal{O} be an arbitrary bounded domain with smooth boundary in $\mathbb{R}^N \setminus \overline{\Omega}$. We define the set

$$C_{\phi_{\rho^*}}^{0,1}(\mathcal{O}) = \{ \phi \in C^{0,1}(\mathcal{O}) \mid \phi|_{\partial\mathcal{O}} = \phi_{\rho^*}|_{\partial\mathcal{O}}, |\nabla \phi| \leq 1 \}$$

and the functional $I_{\mathcal{O}} : C_{\phi_{\rho^*}}^{0,1}(\mathcal{O}) \rightarrow \mathbb{R}$ by

$$I_{\mathcal{O}}(\phi) = \int_{\mathcal{O}} \left(1 - \sqrt{1 - |\nabla \phi|^2} \right) dx.$$

The set of all possible *light rays* in \mathcal{O} is

$$K = \{ \overline{xy} \subset \mathcal{O} \mid x, y \in \partial\mathcal{O}, x \neq y, |\phi_{\rho^*}(x) - \phi_{\rho^*}(y)| = |x - y| \}.$$

Since $\phi_{\rho^*}|_{\mathcal{O}}$ is the unique minimizer of $I_{\mathcal{O}}$ and we infer from [2, Corollary 4.2] that ϕ_{ρ^*} is strictly spacelike in $\mathcal{O} \setminus K$, that it satisfies

$$-\operatorname{div} \left(\frac{\nabla \phi_{\rho^*}}{\sqrt{1 - |\nabla \phi_{\rho^*}|^2}} \right) = 0 \quad \text{in } \mathcal{O} \setminus K,$$

and that

$$\phi_{\rho^*}(tx + (1-t)y) = t\phi_{\rho^*}(x) + (1-t)\phi_{\rho^*}(y), \quad 0 < t < 1,$$

holds for every $x, y \in \partial\mathcal{O}$ such that $|\phi_{\rho^*}(x) - \phi_{\rho^*}(y)| = |x - y|$ and $\overline{xy} \subset \mathcal{O}$. Assume now by contradiction that $K \neq \emptyset$. Then there exist $x, y \in \partial\mathcal{O}$ such that $x \neq y$, $|\phi_{\rho^*}(x) - \phi_{\rho^*}(y)| = |x - y|$, $\overline{xy} \subset \mathcal{O}$. Without loss of generality we can assume that $\phi_{\rho^*}(x) > \phi_{\rho^*}(y)$. It follows that for all $t \in (0, 1)$,

$$(4.5) \quad \phi_{\rho^*}(tx + (1-t)y) = t\phi_{\rho^*}(x) + (1-t)\phi_{\rho^*}(y) = \phi_{\rho^*}(y) + t|x - y|.$$

Since, for any $R > 0$ such that $\mathcal{O} \subset B_R$, the function $\phi_{\rho^*}|_{B_R \setminus \overline{\Omega}}$ is a minimizer of $I_{B_R \setminus \overline{\Omega}}$ on the set $C_{\phi_{\rho^*}}^{0,1}(B_R \setminus \overline{\Omega})$, then, by [2, Theorem 3.2], we have that (4.5) holds for all $t \in \mathbb{R}$ such that $tx + (1-t)y \in B_R \setminus \overline{\Omega}$. Therefore we can stretch the *light ray* until one of these two possibilities occur: either both end-points belong to $\partial\Omega$, or, at least in one direction, the light ray is unbounded. The first case is clearly impossible since $\phi_{\rho^*} = \lambda^*$ on $\partial\Omega$ whereas the second case contradicts the boundedness of ϕ_{ρ^*} . This shows that $K = \emptyset$ and therefore the C^∞ -regularity of ϕ_{ρ^*} on the open set $\mathbb{R}^N \setminus \overline{\Omega}$ follows from [2, Remarks at page 148].

Finally, suppose for contradiction that $\phi_{\rho^*} > 0$ in \mathbb{R}^N fails. Since $\phi_{\rho^*}(x) \rightarrow 0$ as $x \rightarrow \infty$ this implies that ϕ_{ρ^*} attains its non-positive minimum at a point x_0 in \mathbb{R}^N . By Theorem 4.4 and Lemma 4.6 we find that $x_0 \notin \overline{\Omega}$, i.e., x_0 belongs to a connected component Z of $\mathbb{R}^N \setminus \overline{\Omega}$. Since ϕ_{ρ^*}

is a classical solution of (4.4) and since the equation in (4.4) is locally uniformly elliptic in $\mathbb{R}^N \setminus \bar{\Omega}$, we find by the classical maximum-principle that $\phi_{\rho^*} \equiv \phi_{\rho^*}(x_0) \leq 0$ on Z . But $\partial Z \cap \partial\Omega \neq \emptyset$ and $\phi_{\rho^*}|_{\partial\Omega} = \lambda^* > 0$. This contradiction shows that $\phi_{\rho^*} > 0$ in \mathbb{R}^N .

STEP 2 (BOUNDARY REGULARITY): Now we show that for every $r > 0$ sufficiently large, $\phi_{\rho^*} \in C^{2,\alpha}(\bar{B}_r(0) \setminus \Omega)$ and there exists $\theta = \theta(r) \in (0, 1)$ such that $|\nabla\phi_{\rho^*}| \leq 1 - \theta < 1$ in $\bar{B}_r(0)$.

Note that the claim implies that ϕ_{ρ^*} is not only the unique minimizer of \mathcal{I}_{ρ^*} but actually a weak solution of (4.3). The reason is that the strictly spacelike function ϕ_{ρ^*} lies in the interior of \mathcal{X} , and hence variations in all directions can be computed for the minimizer ϕ_{ρ^*} of \mathcal{I}_{ρ^*} which then turns out to be a weak solution of (\mathcal{BI}) . The uniqueness follows from [7, Proposition 2.6] and this ends the proof of the theorem.

The proof relies on an application of [2, Theorem 3.6]. It uses the construction of a strictly spacelike function $\bar{\psi}$ which has the same boundary values as ϕ_{ρ^*} on suitable sets exhausting $\mathbb{R}^N \setminus \Omega$. We begin with the construction of $\bar{\psi}$. For any $\varepsilon > 0$, let us define $\Phi_\varepsilon = \{x \in \mathbb{R}^N \mid \phi_{\rho^*}(x) < \varepsilon\}$, Σ_ε the complementary set in \mathbb{R}^N of the unbounded connected component of Φ_ε and $\Gamma_\varepsilon = \partial\Sigma_\varepsilon$. We set, moreover,

$$R = \max_{x \in \partial\Omega} |x| \quad \text{and} \quad R_\varepsilon = \min_{x \in \Gamma_\varepsilon} |x|.$$

We want to show that

$$(4.6) \quad \exists \bar{\varepsilon} > 0 \text{ such that } \lambda^* + R < R_{\bar{\varepsilon}}.$$

Suppose by contradiction that this does not hold. Then $R_{\frac{1}{n}} \leq \lambda^* + R$ for any $n \geq 1$. Hence, for any $n \geq 1$, there exists $x_n \in \mathbb{R}^N$ such that $|x_n| = R_{\frac{1}{n}} \leq \lambda^* + R$ and $\phi_{\rho^*}(x_n) = \frac{1}{n}$. Since $\{x_n\}_{n \in \mathbb{N}} \subset \bar{B}_{\lambda^*+R}$, there exists $\bar{x} \in \bar{B}_{\lambda^*+R}$ such that, up to a subsequence, $x_n \rightarrow \bar{x}$. Therefore $\phi_{\rho^*}(\bar{x}) = 0$ contradicting the fact that $\phi_{\rho^*} > 0$ in the whole \mathbb{R}^N .

By Sard's Lemma Γ_ε is of class C^∞ for almost all $\varepsilon \in (0, \lambda^*)$. Since moreover $\varepsilon \mapsto R_\varepsilon$ is decreasing, we can find a suitable $\bar{\varepsilon} > 0$ such that $\Gamma_{\bar{\varepsilon}}$ is of class C^∞ and simultaneously (4.6) holds. Let us define $\psi : \mathbb{R}^N \rightarrow \mathbb{R}$, as follows

$$\psi(x) = \begin{cases} \lambda^* & x \in B_R, \\ \frac{\bar{\varepsilon} - \lambda^*}{R_{\bar{\varepsilon}} - R} (|x| - R) + \lambda^* & x \in B_{R_{\bar{\varepsilon}}} \setminus B_R, \\ \bar{\varepsilon} & x \in \mathbb{R}^N \setminus B_{R_{\bar{\varepsilon}}}. \end{cases}$$

Observe that, in $B_{R_{\bar{\varepsilon}}} \setminus B_R$, we have that $|\nabla\psi| = \frac{\lambda^* - \bar{\varepsilon}}{R_{\bar{\varepsilon}} - R} < \frac{\lambda^*}{R_{\bar{\varepsilon}} - R} < 1$, by (4.6). Note that ψ is a Lipschitz function on \mathbb{R}^N . By taking R' slightly larger than R and $R'_{\bar{\varepsilon}}$ slightly smaller than $R_{\bar{\varepsilon}}$ we can construct a corresponding function $\psi' : \mathbb{R}^N \rightarrow \mathbb{R}$ with gradient still bounded away from 1. Mollifying ψ' we obtain a C^∞ -function $\bar{\psi} : \mathbb{R}^N \rightarrow \mathbb{R}$ such that $\bar{\psi} = \lambda^*$ on $\partial\Omega$, $\bar{\psi} = \bar{\varepsilon}$ on $\Gamma_{\bar{\varepsilon}}$ and $|\nabla\bar{\psi}| \leq 1 - \theta_0$ in \mathbb{R}^N for some $\theta_0 > 0$. The function $\bar{\psi}$ is therefore a strictly spacelike extension of the boundary values of ϕ_{ρ^*} to the entire set $\Sigma_{\bar{\varepsilon}} \setminus \bar{\Omega}$. We can therefore apply [2, Theorem 3.6] to conclude that there exists $\theta_{\bar{\varepsilon}} > 0$ such that $|\nabla\phi_{\rho^*}| \leq 1 - \theta_{\bar{\varepsilon}} < 1$ in $\Sigma_{\bar{\varepsilon}} \setminus \Omega$. The claim now follows since Sard's Lemma again implies that there exists a sequence of $\bar{\varepsilon}_n \searrow 0$ such that $\Gamma_{\bar{\varepsilon}_n}$ is C^∞ , (4.6) holds and $\Sigma_{\bar{\varepsilon}_n} \setminus \Omega$ exhausts $\mathbb{R}^N \setminus \Omega$ as $n \rightarrow \infty$. \square

At least for convex bounded sets Ω the function ϕ_ρ , $\rho \in P(\partial\Omega)$, solves pointwise in $\mathbb{R}^N \setminus \bar{\Omega}$ a homogeneous Born-Infeld equation. This is achieved by adapting the proof of Theorem 4.7.

Proposition 4.8. *Let $\Omega \subset \mathbb{R}^N$ bounded and convex. Then, for any $\rho \in P(\partial\Omega)$, ϕ_ρ is a strictly spacelike in $\mathbb{R}^N \setminus \overline{\Omega}$ and satisfies pointwise*

$$\begin{cases} -\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = 0 & \text{in } \mathbb{R}^N \setminus \overline{\Omega}, \\ \phi = \phi_\rho & \text{on } \partial\Omega, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

Proof. Replacing the constant boundary values λ^* by $\phi_\rho|_{\partial\Omega}$ one can follow the arguments of Step 1 of Theorem 4.7 until one reaches the two possibilities for light rays. The second kind of light ray extending to infinity leads to a contradiction as before. However, the first kind of light ray would only extend to two points on $\partial\Omega$, where ϕ_ρ takes certain values (not necessarily equal to the constant λ^* like for ϕ_{ρ^*}). Therefore, no contradiction can be reached in general, and thus these kind of light rays must be a-priori excluded by assuming the convexity of Ω . \square

The next two propositions lead to a unique characterization of the value λ^* , and hence, to the uniqueness of the equilibrium measure (Corollary 4.12). We denote by ν the outer normal to $\partial\Omega$ and write ∂_ν for the normal derivative.

Proposition 4.9. *Let $\Omega \subset \mathbb{R}^N$ be $C^{2,\alpha}$ and $\lambda \in (0, +\infty)$. Then there exists a unique classical solution $\phi_\lambda \in \mathcal{X}$ of*

$$(4.7) \quad \begin{cases} -\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = 0 & \text{in } \mathbb{R}^N \setminus \overline{\Omega}, \\ \phi = \lambda & \text{in } \overline{\Omega}, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

Additionally, $\phi_\lambda \equiv \lambda$ in every bounded connected component of $\mathbb{R}^N \setminus \overline{\Omega}$, and if Z_0 denotes the unbounded connected component of $\mathbb{R}^N \setminus \overline{\Omega}$ then $0 < \phi_\lambda < \lambda$ in Z_0 and $\partial_\nu \phi_\lambda < 0$ on ∂Z_0 . Moreover ϕ_λ is the unique solution of (BT) with $\rho = \rho_\lambda$, where

$$(4.8) \quad d\rho_\lambda := -\frac{\partial_\nu \phi_\lambda(x)}{\sqrt{1 - |\partial_\nu \phi_\lambda(x)|^2}} d\sigma, \quad x \in \partial\Omega,$$

is a positive and bounded measure supported on $\partial\Omega$.

Remark 4.10. *Note that ρ_λ defined in (4.8) is a positive and bounded measure on $\partial\Omega$ with finite total mass, but not necessarily a probability measure.*

Proof of Proposition 4.9. We argue as in the proof of Theorem 4.7. Replacing λ^* by λ in Step 1 we obtain the existence of a unique classical solution ϕ_λ of (4.7), which is strictly spacelike and positive in $\mathbb{R}^N \setminus \overline{\Omega}$. Step 2 then establishes, that ϕ_λ is strictly spacelike and regular up to $\partial\Omega$ and satisfies $\phi_\lambda \in C^{2,\alpha}(\overline{B}_r(0) \setminus \Omega)$ for sufficiently large $r > 0$. In particular $\partial_\nu \phi_\lambda$ is well defined on $\partial\Omega$. Since (4.7) is locally uniformly elliptic we can apply the classical maximum principle to show that $0 < \phi_\lambda < \lambda$ in the unbounded connected component Z_0 of $\mathbb{R}^N \setminus \overline{\Omega}$ and $\phi_\lambda \equiv \lambda$ in every bounded connected component of $\mathbb{R}^N \setminus \overline{\Omega}$. This implies $\partial_\nu \phi_\lambda(x) \leq 0$ for all $x \in \partial\Omega$, and the Hopf Lemma even allows to conclude $\partial_\nu \phi_\lambda < 0$ on $\partial Z_0 \subset \partial\Omega$, where ν is the exterior unit normal on $\partial\Omega$. Finally, let us show that ϕ_λ solves (BT) with right-hand side $\rho = \rho_\lambda$ given by (4.8). Since ϕ_λ is

constant in $\bar{\Omega}$, for any $\varphi \in C_c^\infty(\mathbb{R}^N)$ we have

$$\begin{aligned} \int_{\mathbb{R}^N} \frac{\nabla\phi_\lambda \cdot \nabla\varphi}{\sqrt{1 - |\nabla\phi_\lambda|^2}} &= \int_{\mathbb{R}^N \setminus \bar{\Omega}} \frac{\nabla\phi_\lambda \cdot \nabla\varphi}{\sqrt{1 - |\nabla\phi_\lambda|^2}} \\ &= \int_{\mathbb{R}^N \setminus \bar{\Omega}} -\operatorname{div} \left(\frac{\nabla\phi_\lambda}{\sqrt{1 - |\nabla\phi_\lambda|^2}} \right) \varphi - \int_{\partial\Omega} \frac{\partial_\nu\phi_\lambda}{\sqrt{1 - |\partial_\nu\phi_\lambda|^2}} \varphi \, d\sigma \\ &= - \int_{\partial\Omega} \frac{\partial_\nu\phi_\lambda}{\sqrt{1 - |\partial_\nu\phi_\lambda|^2}} \varphi \, d\sigma \end{aligned}$$

which shows that ϕ_λ is the (unique) weak solution of (\mathcal{BI}) with ρ_λ given by (4.8). \square

The next proposition shows that λ^* is the unique value λ such that ρ_λ is actually a probability measure.

Proposition 4.11. *Let $\Omega \subset \mathbb{R}^N$ be of class $C^{2,\alpha}$ and for all $\lambda > 0$ let ϕ_λ be the unique solution of (4.7). Then the value λ^* from Proposition 4.2 is the unique value of $\lambda \in (0, +\infty)$ such that the measure*

$$d\rho_\lambda = - \frac{\partial_\nu\phi_\lambda}{\sqrt{1 - |\partial_\nu\phi_\lambda|^2}} d\sigma, \quad x \in \partial\Omega,$$

is a probability measure on $\partial\Omega$.

Proof. Step 1 in Theorem 4.7 shows that $\phi_{\rho^*} = \phi_{\lambda^*}$. Moreover, since Theorem 4.7 and Proposition 4.9 also imply that ϕ_{ρ^*} and ϕ_{λ^*} are solutions of (\mathcal{BI}) with right-hand sides ρ^* and ρ_{λ^*} respectively, we can conclude that $\rho^* = \rho_{\lambda^*}$. Thus the existence of a value λ such that ρ_λ is a probability measure on $\partial\Omega$ is proven and we just need to show its uniqueness. We can conclude if we prove that the map

$$\Upsilon : \lambda \mapsto \int_{\partial\Omega} d\rho_\lambda \quad \text{for } \lambda \in (0, +\infty)$$

is strictly increasing. For this purpose let $\lambda_1, \lambda_2 \in (0, +\infty)$ with $\lambda_1 < \lambda_2$ and, for $i = 1, 2$, $\phi_i := \phi_{\lambda_i}$ be the corresponding solutions of (4.7) and $\rho_i := \rho_{\lambda_i}$. Let us observe that $\tilde{\phi}_1 := \phi_1 + \lambda_2 - \lambda_1$ and ϕ_2 satisfy both the following problem

$$\begin{cases} -\operatorname{div} \left(\frac{\nabla\phi}{\sqrt{1 - |\nabla\phi|^2}} \right) = 0 & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ \phi = \lambda_2 & \text{in } \bar{\Omega}, \end{cases}$$

while

$$\lim_{|x| \rightarrow \infty} \tilde{\phi}_1(x) = \lambda_2 - \lambda_1 > 0 = \lim_{|x| \rightarrow \infty} \phi_2(x).$$

Next we apply a comparison argument to $\tilde{\phi}_1$ and ϕ_2 , cf. [24, Theorem 10.1]. For this purpose, let $F(\xi) = (1 - |\xi|^2)^{-1/2}$, $\xi \in \mathbb{R}^N$ with $|\xi| < 1$. Notice that for $\psi := \tilde{\phi}_1 - \phi_2$ and $\chi_t := t\tilde{\phi}_1 + (1-t)\phi_2$, $t \in [0, 1]$ we have

$$\frac{\nabla\tilde{\phi}_1}{\sqrt{1 - |\nabla\tilde{\phi}_1|^2}} - \frac{\nabla\phi_2}{\sqrt{1 - |\nabla\phi_2|^2}} = F(\nabla\tilde{\phi}_1)\nabla\tilde{\phi}_1 - F(\nabla\phi_2)\nabla\phi_2 = \int_0^1 \frac{d}{dt} (F(\nabla\chi_t)\nabla\chi_t) dt = a(x)\nabla\psi,$$

where

$$a(x) := \int_0^1 (F(\nabla\chi_t(x)) + F'(\nabla\chi_t(x)) \cdot \nabla\chi_t(x)) dt = \int_0^1 (1 - |\nabla\chi_t(x)|^2)^{-3/2} dt.$$

Arguing as in Theorem 4.7 the function $a(x) > 0$ is bounded in bounded subsets of $\mathbb{R}^N \setminus \bar{\Omega}$. Hence, ψ satisfies the locally uniformly elliptic equation

$$\begin{cases} -\operatorname{div}(a(x)\nabla\psi) = 0 & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ \psi = 0 & \text{in } \bar{\Omega}, \\ \lim_{|x| \rightarrow \infty} \psi(x) = \lambda_2 - \lambda_1 > 0. \end{cases}$$

Therefore, the maximum principle applies and states that ψ attains its zero minimum on $\partial\Omega$ and its positive maximum at infinity. On every bounded connected component Z of $\mathbb{R}^N \setminus \bar{\Omega}$ we get that $\psi \equiv 0$ since $\psi = 0$ on ∂Z . On the unbounded component Z_0 of $\mathbb{R}^N \setminus \bar{\Omega}$ we get $0 < \psi < \lambda_2 - \lambda_1$ and the Hopf Lemma implies that $\partial_\nu \psi > 0$ on $\partial Z_0 \subset \partial\Omega$, where ν is the exterior unit normal on $\partial\Omega$. Thus, $\partial_\nu \phi_1 = \partial_\nu \tilde{\phi}_1 > \partial_\nu \phi_2$ on ∂Z_0 which implies that $\Upsilon(\lambda_1) < \Upsilon(\lambda_2)$ as claimed. \square

An immediate consequence of Proposition 4.11 is the following uniqueness result.

Corollary 4.12. *If $\Omega \subset \mathbb{R}^N$ is of class $C^{2,\alpha}$ then the equilibrium measure is unique.*

Without regularity assumptions on the domain we are able to show only some partial result about the uniqueness of the equilibrium measure. This result is strictly related to the conjecture that any minimizer of the functional \mathcal{I}_ρ is also a (weak) solution of the corresponding PDE.

Proposition 4.13. *Let $\Omega \subset \mathbb{R}^N$ be bounded with no further regularity of $\partial\Omega$. Suppose that the equilibrium potential weakly solves (\mathcal{BI}) . Then the equilibrium measure is unique.*

Proof. Let ρ^* be an equilibrium measure for which ϕ_{ρ^*} weakly solves (\mathcal{BI}) with right-hand side ρ^* . Let ρ^{**} be any other equilibrium measure. From Corollary 4.5 we know that the equilibrium potentials ϕ_{ρ^*} and $\phi_{\rho^{**}}$ coincide. Since ϕ_{ρ^*} solves (\mathcal{BI}) , we have in particular

$$(4.9) \quad \int_{\mathbb{R}^N} \frac{|\nabla \phi_{\rho^*}|^2}{\sqrt{1 - |\nabla \phi_{\rho^*}|^2}} dx = \langle \rho^*, \phi_{\rho^*} \rangle.$$

But since $\langle \rho^*, \phi_{\rho^*} \rangle = \langle \rho^{**}, \phi_{\rho^{**}} \rangle$ we get that (4.9) also holds for $\phi_{\rho^{**}}$. By [7, Remark 2.8] this implies that $\phi_{\rho^{**}}$ also solves (\mathcal{BI}) with right-hand side ρ^{**} . Hence $\phi_{\rho^*}, \phi_{\rho^{**}}$ are (identical) weak solutions of (\mathcal{BI}) with right-hand sides ρ^*, ρ^{**} , and therefore these measures coincide. \square

We conclude this section with the following observation: the functionals \mathcal{E} and \mathcal{K} (cf. Remark 2.4) coincide on solutions of (\mathcal{BI}) but they can differ at a minimizer ϕ_ρ of \mathcal{I}_ρ in case it is not a solution. However, since \mathcal{K} is well defined on each minimizer of the functional \mathcal{I}_ρ for all $\rho \in \mathcal{X}^*$ we can also study the minimization of $\mathcal{K}(\phi_\rho)$ with respect to $\rho \in P(\partial\Omega)$. The result is given next.

Proposition 4.14. *There exists a measure $\tilde{\rho}$ such that*

$$(4.10) \quad \mathcal{K}(\phi_{\tilde{\rho}}) = \inf_{\rho \in P(\partial\Omega)} \mathcal{K}(\phi_\rho).$$

If $\phi_{\tilde{\rho}}$ and ϕ_{ρ^} are both solutions of (\mathcal{BI}) with the respective measures, then $\tilde{\rho} = \rho^*$.*

Proof. By Remark 2.4 we know that $\mathcal{K}(\phi_\rho) \leq \mathcal{E}(\phi_\rho)$ for every $\rho \in \mathcal{X}^*$. Moreover, if we take a minimizing sequence $\{\rho_k\}_{k \in \mathbb{N}} \subset P(\partial\Omega)$ for \mathcal{K} , there exists $\tilde{\rho} \in P(\partial\Omega)$ such that, up to a subsequence, $\rho_k \rightharpoonup \tilde{\rho}$ weakly in the sense of measures and, by Lemma 3.7, $\phi_{\rho_k} \rightarrow \phi_{\tilde{\rho}}$ strongly in $D^{1,2}(\mathbb{R}^N)$. Since \mathcal{K} is lower-semicontinuous, cf. Remark 4.15, one finds

$$\mathcal{K}(\phi_{\tilde{\rho}}) \leq \liminf_k \mathcal{K}(\phi_{\rho_k}) = \inf_{\rho \in P(\partial\Omega)} \mathcal{K}(\phi_\rho),$$

and so we get (4.10). Finally, recalling that \mathcal{E} and \mathcal{K} coincide on solutions of (\mathcal{BI}) , we observe that

$$\inf_{\rho \in P(\partial\Omega)} \mathcal{E}(\phi_\rho) = \mathcal{E}(\phi_{\rho^*}) = \mathcal{K}(\phi_{\rho^*}) \geq \mathcal{K}(\phi_{\tilde{\rho}}) = \mathcal{E}(\phi_{\tilde{\rho}}) \geq \mathcal{E}(\phi_{\rho^*})$$

and hence equality holds. Therefore we conclude by Proposition 4.13. \square

Remark 4.15. *The functional $\mathcal{K} : \mathcal{X} \rightarrow [0, \infty]$ is a proper, convex and lower semicontinuous functional with domain*

$$\text{dom}(\mathcal{K}) = \{\phi \in \mathcal{X} : \mathcal{K}(\phi) < \infty\}.$$

Moreover, \mathcal{K} is continuous on $\text{dom}(\mathcal{K})^\circ$. Lower-semicontinuity follows from Fatou's lemma. The continuity on $\text{dom}(\mathcal{K})^\circ$ can be found in [18, Corollary 2.5].

5. THE APPROXIMATED PROBLEM

Using (1.1) and (3.1), we can approximate, at least formally, the equation (BT) by

$$(5.1) \quad \begin{cases} -\sum_{h=1}^n \alpha_h \Delta_{2h} \phi = \rho & \text{in } \mathbb{R}^N, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0 \end{cases}$$

where for $h \in \mathbb{N}$ the operator $\Delta_{2h}(\cdot) = \text{div}(|\nabla \cdot|^{2h-2} \nabla \cdot)$ is the $2h$ -Laplacian. Weak solutions of (5.1) can be found as critical points of the Lagrangian functional

$$\mathcal{I}_\rho^n(\phi) = \sum_{h=1}^n \frac{\alpha_h}{2h} \|\nabla \phi\|_{2h}^{2h} - \langle \rho, \phi \rangle_n$$

on the space \mathcal{X}_{2n} , which is defined as the completion of $C_c^\infty(\mathbb{R}^N)$ with respect to the norm $\|\nabla \cdot\|_2 + \|\nabla \cdot\|_{2n}$. A similar construction can be found in [27]. Naturally, we assume that $\rho \in \mathcal{X}_{2n}^*$. The symbol $\langle \cdot, \cdot \rangle_n$ denotes the duality bracket between \mathcal{X}_{2n} and \mathcal{X}_{2n}^* . Because of the continuity, convexity and coercivity of the functional \mathcal{I}_ρ^n there exists a unique minimizer ϕ_ρ^n on \mathcal{X}_{2n} which is also a weak solution of (5.1) and $\mathcal{I}_\rho^n(\phi_\rho^n) \leq 0$. Observe that for all $m \leq n$ we have that $\mathcal{X} \subset \mathcal{X}_{2n} \subset \mathcal{X}_{2m}$.

As in the Maxwell and Born-Infeld case, cf. Section 2, we define the electrostatic energy as $\mathcal{E}^n(\phi) := -\mathcal{I}_\rho^n(\phi)$. For weak solutions ϕ_ρ^n of (5.1) the energy coincides with the expression

$$\mathcal{E}^n(\phi_\rho^n) = -\mathcal{I}_\rho^n(\phi_\rho^n) = \mathcal{K}^n(\phi_\rho^n) \quad \text{where} \quad \mathcal{K}^n(\phi) := \sum_{h=1}^n \frac{2h-1}{2h} \alpha_h \|\nabla \phi\|_{2h}^{2h}.$$

For this reason, in the discussion of equilibrium measures we will consider only the following minimization problem

$$\min_{\rho \in P(\partial\Omega)} \mathcal{K}^n(\phi_\rho^n).$$

The next result shows that, for n large, the functional \mathcal{I}_ρ^n is well defined in \mathcal{X}_{2n} for all $\rho \in P(\partial\Omega)$ and so, the existence of the unique minimizer ϕ_ρ^n is guaranteed.

Lemma 5.1. *If $n > N/2$ then $\mathcal{X}_{2n} \subset C_b(\mathbb{R}^N)$ and hence $P(\partial\Omega) \subset \mathcal{X}_{2n}^*$. Moreover, \mathcal{X}_{2n} embeds compactly into $C_b(D)$ for every bounded set $D \subset \mathbb{R}^N$.*

Proof. If $\phi \in \mathcal{X}_{2n}$, then $|\nabla \phi| \in L^2(\mathbb{R}^N)$ and therefore $\phi \in L^{2^*}(\mathbb{R}^N)$. We will show that there exists $C > 0$ such that

$$\|\phi\|_\infty \leq C(\|\nabla \phi\|_2 + \|\nabla \phi\|_{2n}).$$

We argue similarly as in [19, Proposition 8]. Let us consider a family of N -dimensional cubes Q_k such that $|Q_k| = 1$ and $\bigcup_k Q_k = \mathbb{R}^N$ and $\varphi \in C_c^\infty(\mathbb{R}^N)$. For every $x, y \in \mathbb{R}^N$ and $t \in [0, 1]$ we have

$$\varphi(x) - \varphi(y) = \int_0^1 \nabla \varphi(tx + (1-t)y) \cdot (x-y) dt,$$

and averaging with respect to y on an arbitrary Q_k ,

$$\varphi(x) - \int_{Q_k} \varphi(y) dy = \int_{Q_k} \left(\int_0^1 \nabla \varphi(tx + (1-t)y) \cdot (x-y) dt \right) dy.$$

Then, if $x \in Q_k$,

$$\begin{aligned} |\varphi(x)| &\leq \left| \int_{Q_k} \varphi(y) dy \right| + \int_{Q_k} \left(\int_0^1 |\nabla \varphi(tx + (1-t)y) \cdot (x-y)| dt \right) dy \\ &\leq \|\varphi\|_{L^{2^*}(Q_k)} + \sum_{i=1}^N \int_{Q_k} \left(\int_0^1 |\partial_i \varphi(tx + (1-t)y)| dt \right) dy \\ &\leq C \|\nabla \varphi\|_2 + \sum_{i=1}^N \int_0^1 \left[\frac{1}{(1-t)^N} \left(\int_{(1-t)Q_k+tx} |\partial_i \varphi(y)|^{2n} dy \right)^{\frac{1}{2n}} (1-t)^{\frac{N(2n-1)}{2n}} \right] dt \\ &\leq C \|\nabla \varphi\|_2 + N \int_0^1 \left[(1-t)^{-\frac{N}{2n}} \|\nabla \varphi\|_{L^{2n}((1-t)Q_k+tx)} \right] dt \\ &\leq C \|\nabla \varphi\|_2 + \frac{2Nn}{2n-N} \|\nabla \varphi\|_{2n}. \end{aligned}$$

The conclusion $\mathcal{X}_{2n} \subset C_b(\mathbb{R}^N)$ follows now from a density argument. Since the L^∞ -estimate implies that $\mathcal{X}_{2n} \subset W^{1,2n}(B)$ for every ball $B \subset \mathbb{R}^N$ and since we have $2n > N$ we get the compact embedding of \mathcal{X}_{2n} into $C_b(B)$. \square

Lemma 5.2. *Let $n > N/2$. Then there exists a constant $C_n > 0$ such that for every $\rho \in P(\partial\Omega)$*

$$\|\nabla \phi_\rho^n\|_2 + \|\nabla \phi_\rho^n\|_{2n} \leq C_n.$$

The constant C_n is uniformly bounded if n is bounded away from $N/2$.

Proof. The proof of Lemma 5.1 shows that there exists $C_n > 0$ with

$$\|\phi_\rho^n\|_\infty \leq C_n (\|\nabla \phi_\rho^n\|_2 + \|\nabla \phi_\rho^n\|_{2n}),$$

and that C_n is uniformly bounded if n is bounded away from $N/2$. Then

$$0 \geq \mathcal{I}_\rho^n(\phi_\rho^n) \geq \frac{\alpha_n}{2n} \|\nabla \phi_\rho^n\|_{2n}^2 + \frac{\alpha_1}{2} \|\nabla \phi_\rho^n\|_2^2 - C_n (\|\nabla \phi_\rho^n\|_2 + \|\nabla \phi_\rho^n\|_{2n})$$

and we conclude. \square

Next we give the counterpart of Lemma 3.7. The proof, based on Clarkson's inequalities, is almost the same as the one of Lemma 3.7 with Lemma 5.2 replacing Lemma 3.6. We omit the details.

Lemma 5.3 (Continuous dependence of ϕ_ρ^n on ρ). *Let $n > N/2$. If $\rho_k, \rho \in P(\partial\Omega)$ with $\rho_k \rightharpoonup \rho$ as $k \rightarrow \infty$ weakly in the sense of measures then $\phi_{\rho_k}^n \rightarrow \phi_\rho^n$ strongly in \mathcal{X}_{2n} and locally uniformly on \mathbb{R}^N , as $k \rightarrow \infty$.*

We now establish existence and uniqueness of the equilibrium measure $\rho^{*,n}$ for \mathcal{K}^n and show that $\phi_{\rho^{*,n}}^n$ is constant on $\bar{\Omega}$. First, we assume as before only boundedness of Ω and no further regularity. We need to adapt the proof of Lemma 3.8. Since ϕ_ρ^n is a weak solution of (5.1) on \mathbb{R}^N we get that ϕ_ρ^n weakly solves

$$-\sum_{h=1}^n \alpha_h \Delta_{2h} \phi_\rho^n = 0 \quad \text{on } B$$

where B is an open ball with $\bar{B} \cap \text{supp } \rho = \emptyset$. By [31, Lemma 1] we obtain that ϕ_ρ^n is C^1 on $\mathbb{R}^N \setminus \text{supp } \rho$ and hence it satisfies (3.9) on B with $a(x) = \sum_{h=1}^n \alpha_h |\nabla \phi_\rho^n(x)|^{2h-2}$ being a continuous function which is locally bounded on $\mathbb{R}^N \setminus \text{supp } \rho$ and bounded away from zero. Therefore, [24, Theorem 8.19] applies and we can finish the proof as in Lemma 3.8. The remaining results of Section 3 stay valid for ϕ_ρ^n without any change. Likewise, the results of Section 4 from Theorem 4.1 up to and including Lemma 4.6 stay true without change. This all works under the only assumption that Ω is bounded. In particular, we have that the equilibrium potential is always unique.

It remains to establish uniqueness of the equilibrium measure, and here we need (as before) more regularity of $\partial\Omega$. We do not need any analogue for Theorem 4.7 since we already have that $\phi_{\rho^{*,n}}^n$ is a weak solution of (5.1) with right-hand side $\rho^{*,n}$. Furthermore $\phi_{\rho^{*,n}}^n$ weakly solves

$$(5.2) \quad \begin{cases} -\sum_{h=1}^n \alpha_h \Delta_{2h} \phi = 0 & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ \phi = \lambda & \text{in } \bar{\Omega}, \\ \lim_{|x| \rightarrow \infty} \phi(x) = 0. \end{cases}$$

with $\lambda = \lambda^{*,n} = \langle \rho^{*,n}, \phi_{\rho^{*,n}}^n \rangle$. The existence of a weak solution ϕ_λ of (5.2) for any $\lambda > 0$ follows from minimization of a suitable functional in a space similar to \mathcal{X}_{2n} . We leave these details to the reader. The $C^{1,\alpha}$ -regularity of this ϕ_λ on $\bar{B}_R(0) \setminus \Omega$ for every $R > 0$ follows by assuming $\partial\Omega \in C^{1,\alpha}$ and combining Theorem 1 and Lemma 1 from [31]. This leads to the definition of the positive and bounded measure

$$(5.3) \quad d\rho_\lambda^n := - \left(\sum_{h=1}^n \alpha_h |\partial_\nu \phi_\lambda^n|^{2h-2} \right) \partial_\nu \phi_\lambda^n d\sigma$$

as in Proposition 4.9. Note that due to the uniform ellipticity of (5.2) in a neighbourhood of the boundary, the Hopf Lemma shows that $d\rho_\lambda^n$ is strictly positive on the boundary of the unbounded connected component of $\mathbb{R}^N \setminus \bar{\Omega}$. The characterization of the value $\lambda^{*,n}$ as the unique value such that $d\rho_\lambda^n$ is a probability measure on $\partial\Omega$ is then established in the same way as in Proposition 4.11 with the direct consequence of uniqueness of the equilibrium measure $\rho^{*,n}$. Notice that this works under the assumption $\partial\Omega \in C^{1,\alpha}$. The following statement summarizes these results.

Theorem 5.4. *Let $n > N/2$ and assume $\partial\Omega \in C^{1,\alpha}$. There exists a unique equilibrium measure $\rho^{*,n} \in P(\partial\Omega)$ for \mathcal{K}^n . Moreover, $\phi_{\rho^{*,n}}^n = \lambda^{*,n}$ on $\bar{\Omega}$ and for every $\mu \in P(\partial\Omega)$*

$$(5.4) \quad \lambda^{*,n} := \langle \rho^{*,n}, \phi_{\rho^{*,n}}^n \rangle = \langle \mu, \phi_{\rho^{*,n}}^n \rangle.$$

Remark 5.5. *Based on the local $C^{1,\alpha}$ regularity of $\phi_{\rho^{*,n}}^n$ we can also improve the regularity. In case $\partial\Omega \in C^{2,\alpha}$ we can interpret the differential equation $-\sum_{h=1}^n \alpha_h \Delta_h \phi_{\rho^{*,n}}^n = 0$ on $\mathbb{R}^N \setminus \Omega$ as a linear differential equation for $\phi_{\rho^{*,n}}^n$ with locally α -Hölder continuous and locally uniformly elliptic coefficients. This leads to local $C^{2,\alpha}$ regularity up to the boundary and in particular that $\phi_{\rho^{*,n}}^n$ is a classical solution on $\mathbb{R}^N \setminus \Omega$.*

The remaining part of this section is devoted to show that the sequence $\rho^{*,n}$ weakly converges to a limit measure $\bar{\rho}$ (Lemma 5.8) which is in fact an equilibrium measure for the Born-Infeld energy \mathcal{E} (Proposition 5.9).

Lemma 5.6. *Let $N/2 < m < n$ and $\rho \in P(\partial\Omega)$. Then*

$$\mathcal{I}_\rho^m(\phi_\rho^m) < \mathcal{I}_\rho^n(\phi_\rho^n)$$

and

$$\mathcal{K}^n(\phi_\rho^n) < \mathcal{K}^m(\phi_\rho^m).$$

Proof. The conclusion follows easily observing that

$$\mathcal{I}_\rho^m(\phi_\rho^m) \leq \mathcal{I}_\rho^m(\phi_\rho^n) < \mathcal{I}_\rho^n(\phi_\rho^n)$$

and, since ϕ_ρ^n, ϕ_ρ^m are weak solutions,

$$\mathcal{K}^n(\phi_\rho^n) = -\mathcal{I}_\rho^n(\phi_\rho^n) < -\mathcal{I}_\rho^m(\phi_\rho^m) = \mathcal{K}^m(\phi_\rho^m).$$

□

For $n \in \mathbb{N}$, let us consider $\mathfrak{k}^n := \min_{\rho \in P(\partial\Omega)} \mathcal{K}^n(\phi_\rho) = \mathcal{K}^n(\phi_{\rho^{*,n}}^n)$ and $\mathcal{I}^n := \min_{\psi \in \mathcal{X}_{2n}} \mathcal{I}_{\rho^{*,n}}^n(\psi) = \mathcal{I}_{\rho^{*,n}}^n(\phi_{\rho^{*,n}}^n)$. As an immediate consequence of Lemma 5.6 and of the fact that $\mathcal{I}^n = -\mathfrak{k}^n$, we have the following monotonicity result.

Lemma 5.7. *The sequence $\{\mathfrak{k}^n\}_{n>N/2}$ is strictly decreasing and the sequence $\{\mathcal{I}^n\}_{n>N/2}$ is strictly increasing.*

Lemma 5.8. *There exists $\bar{\rho} \in P(\partial\Omega)$ such that $\rho^{*,n} \rightharpoonup \bar{\rho}$ weakly in the sense of measures, $\phi_{\rho^{*,n}}^n \rightharpoonup \phi_{\bar{\rho}}$ weakly in \mathcal{X}_{2m} for all $m > N/2$ and uniformly on compact subsets of \mathbb{R}^N as $n \rightarrow \infty$. Here $\phi_{\bar{\rho}} \in \mathcal{X}$ is the unique minimizer of $\mathcal{I}_{\bar{\rho}} : \mathcal{X} \rightarrow \mathbb{R}$. Moreover*

$$(5.5) \quad \mathcal{K}^n(\phi_{\bar{\rho}}) \leq \liminf_n \mathcal{K}^n(\phi_{\rho^{*,n}}^n).$$

Proof. Let $\bar{\rho} \in P(\partial\Omega)$ such that, up to subsequences, $\rho^{*,n} \rightharpoonup \bar{\rho}$ weakly in the sense of measures, as $n \rightarrow \infty$. Let $m, n \in \mathbb{N}$ with $N/2 < m \leq n$. Since

$$\mathcal{K}^m(\phi_{\rho^{*,n}}^n) \leq \mathcal{K}(\phi_{\rho^{*,n}}^n) = \mathfrak{k}^n \leq \mathfrak{k}^1,$$

we see that $\|\nabla \phi_{\rho^{*,n}}^n\|_2 + \|\nabla \phi_{\rho^{*,n}}^n\|_{2m} \leq C(m)$. Therefore, up to a subsequence, $\{\phi_{\rho^{*,n}}^n\}_{n \in \mathbb{N}}$ is weakly convergent in \mathcal{X}_{2m} and, by Lemma 5.1, locally uniformly in \mathbb{R}^N . By a diagonalization argument, we deduce that the weak limit $\phi \in \bigcap_{m \geq 1} \mathcal{X}_{2m}$. The fact that $\phi \in \mathcal{X}$ (i.e., $\|\nabla \phi\|_\infty \leq 1$) follows as in [7, Theorem 5.2]. Let $\mathcal{I} := \lim_{n \rightarrow \infty} \mathcal{I}^n$.

Let us show that $\mathcal{I} = \mathcal{I}_{\bar{\rho}}(\phi)$. First note that

$$\mathcal{I}^n \leq \mathcal{I}_{\rho^{*,n}}^n(\phi) \leq \mathcal{I}_{\rho^{*,n}}(\phi) = \mathcal{I}_{\bar{\rho}}(\phi) + \underbrace{\langle \bar{\rho} - \rho^{*,n}, \phi \rangle}_{\rightarrow 0 \text{ as } n \rightarrow \infty}$$

which implies that $\mathcal{I} \leq \mathcal{I}_{\bar{\rho}}(\phi)$. Now let us show the reverse inequality. Next note that since $\mathcal{I}_{\rho^{*,m}}^m(\phi_{\rho^{*,n}}^n) \leq \mathcal{I}_{\rho^{*,m}}^m(\phi_{\rho^{*,n}}^n)$, for all $m \leq n$, and by weak lower semicontinuity

$$\begin{aligned} \mathcal{I}_{\rho^{*,m}}^m(\phi) &\leq \liminf_n \mathcal{I}_{\rho^{*,m}}^m(\phi_{\rho^{*,n}}^n) \leq \liminf_n \underbrace{\mathcal{I}_{\rho^{*,n}}^n(\phi_{\rho^{*,n}}^n)}_{=\mathcal{I}^n} + \lim_{n \rightarrow \infty} \langle \rho^{*,n} - \rho^{*,m}, \phi_{\rho^{*,n}}^n \rangle \\ &= \mathcal{I} + \langle \bar{\rho} - \rho^{*,m}, \phi \rangle \end{aligned}$$

where we have used that $\phi_{\rho^{*,n}}^n \rightarrow \phi$ uniformly on $\partial\Omega$ as $n \rightarrow \infty$. Subtracting $\langle \bar{\rho} - \rho^{*,m}, \phi \rangle$ from both sides, we get

$$\mathcal{I}_{\bar{\rho}}^m(\phi) \leq \mathcal{I}.$$

Passing to the limit on m we get $\mathcal{I}_{\bar{\rho}}(\phi) \leq \mathcal{I}$. Altogether we have obtained the claim $\mathcal{I}_{\bar{\rho}}(\phi) = \mathcal{I}$.

To see that $\phi = \phi_{\bar{\rho}}$ observe first that we have the inequalities

$$\mathcal{I}^n \leq \mathcal{I}_{\rho^{*,n}}^n(\psi) \leq \mathcal{I}_{\bar{\rho}}(\psi) + \langle \bar{\rho} - \rho^{*,n}, \psi \rangle \quad \text{for all } \psi \in \mathcal{X}.$$

This implies

$$\mathcal{I}_{\bar{\rho}}(\phi) = \mathcal{I} = \lim_{n \rightarrow \infty} \mathcal{I}^n \leq \mathcal{I}_{\bar{\rho}}(\psi) \quad \text{for all } \psi \in \mathcal{X}$$

so that $\phi = \phi_{\bar{\rho}}$.

Finally, it remains to show that (5.5) holds. By the monotonicity of \mathfrak{k}^n from Lemma 5.7 and $\mathfrak{k}^n \geq 0$ we infer that $\mathfrak{k}^\infty := \lim_{n \rightarrow \infty} \mathfrak{k}^n$ exists. Notice that the lower semicontinuity and the fact that $\mathcal{K}^m(\phi_{\rho^{*,n}}^n) \leq \mathcal{K}^n(\phi_{\rho^{*,n}}^n)$ for $m \leq n$, imply that, for every $m \in \mathbb{N}$, we have

$$\mathcal{K}^m(\phi_{\bar{\rho}}) \leq \liminf_n \mathcal{K}^m(\phi_{\rho^{*,n}}^n) \leq \liminf_n \mathcal{K}^n(\phi_{\rho^{*,n}}^n) = \mathfrak{k}^\infty.$$

Hence (5.5) holds. \square

Now we can finally show that the weak limit $\bar{\rho}$ of the sequence $\rho^{*,n}$ is indeed an equilibrium measure for the Born-Infeld energy \mathcal{E} .

Proposition 5.9. *The measure $\bar{\rho}$ is an equilibrium measure for \mathcal{E} considered in the Section 4. In particular we have $\phi_{\rho^*} = \phi_{\bar{\rho}}$. If $\partial\Omega$ is $C^{2,\alpha}$ for some $\alpha > 0$ then $\bar{\rho}$ is the unique equilibrium measure for \mathcal{E} .*

Proof. By (5.4) and Lemma 5.8, we have that for every $\mu \in P(\partial\Omega)$

$$\bar{\lambda} := \langle \bar{\rho}, \phi_{\bar{\rho}} \rangle = \langle \mu, \phi_{\bar{\rho}} \rangle.$$

Hence

$$\mathcal{I}_\mu(\phi_\mu) \leq \mathcal{I}_\mu(\phi_{\bar{\rho}}) = \mathcal{I}_{\bar{\rho}}(\phi_{\bar{\rho}}) + \langle \bar{\rho} - \mu, \phi_{\bar{\rho}} \rangle = \mathcal{I}_{\bar{\rho}}(\phi_{\bar{\rho}}),$$

which implies that $\bar{\rho}$ is an equilibrium measure for \mathcal{E} . Since by Corollary 4.5 we know that the equilibrium potential for \mathcal{E} is unique, this implies $\phi_{\rho^*} = \phi_{\bar{\rho}}$. Uniqueness of the equilibrium measure under the assumption $\partial\Omega \in C^{2,\alpha}$ is already contained in Corollary 4.12. \square

6. CHARACTERIZATION OF BALLS VIA EQUILIBRIUM MEASURES

Theorem 6.1. *Let $\Omega \subset \mathbb{R}^N$ be of class $C^{2,\alpha}$ and let ϕ_{ρ^*} be the equilibrium potential for the Born-Infeld electrostatic model. Then Ω is a ball if and only if the equilibrium distribution ρ^* is a constant multiple of the surface measure $d\sigma$ on $\partial\Omega$. The same characterization is true for the approximated model.*

Proof. One direction of the theorem is trivial: if Ω is a ball, then the uniqueness of the equilibrium potentials (see Corollary 4.5 and Theorem 5.4) implies their radial symmetry. In particular, $\partial_\nu \phi_{\rho^*}$ and $\partial_\nu \phi_{\rho^{*,n}}^n$ are constant on the boundary of the ball. Therefore, using Proposition 4.11 for ϕ_{ρ^*} and its counterpart, formula (5.3), for $\phi_{\rho^{*,n}}^n$, the corresponding equilibrium measures given by

$$(6.1) \quad d\rho^* = -\frac{\partial_\nu \phi_{\rho^*}}{\sqrt{1 - |\partial_\nu \phi_{\rho^*}|^2}} d\sigma$$

and

$$(6.2) \quad d\rho^{*,n} = -\left(\sum_{h=1}^n \alpha_h |\partial_\nu \phi_{\rho^{*,n}}^n|^{2h-2} \right) \partial_\nu \phi_{\rho^{*,n}}^n d\sigma$$

are both constant multiples of $d\sigma$ on the boundary of the ball.

Now we consider the nontrivial part of the theorem. It will follow from an application and slight modification of [35, Theorem 1], which itself is based on the moving plane method of Alexandrov-Serrin [1, 39]. Define

$$g(s) = \frac{1}{\sqrt{1-s^2}} \text{ for } s \in [0, 1) \text{ and } g_n(s) = \sum_{h=1}^n \alpha_h s^{2h-2} \text{ for } s \in [0, \infty)$$

with α_h as given in (1.1). Notice that

$$g \in C^2[0, 1), \text{ and } g(s) > 0, (g(s)s)' > 0 \text{ for all } s \in [0, 1)$$

and

$$g_n \in C^2[0, \infty), \text{ and } g_n(s) > 0, (g_n(s)s)' > 0 \text{ for all } s \in [0, \infty).$$

Observe that the function g_n directly satisfies condition (I)₁ of [35, Theorem 1], whereas g satisfies (I)₁ not on $[0, \infty)$ but only on $[0, 1)$. As we will see later this is still sufficient since $[0, 1)$ covers the range of $|\nabla\phi_{\rho^*}|$.

Next we recall the properties of ϕ_{ρ^*} and $\phi_{\rho^{*,n}}^n$. Since we are assuming that ρ^* and $\rho^{*,n}$ are constant multiples of the surface measure $d\sigma$ on $\partial\Omega$, and by (6.1) and (6.2), we get that the functions $g(\partial_\nu\phi_{\rho^*})\partial_\nu\phi_{\rho^*}$ and $g_n(\partial_\nu\phi_{\rho^{*,n}}^n)\partial_\nu\phi_{\rho^{*,n}}^n$ are both constant on $\partial\Omega$. Since $g(s)s$ and $g_n(s)s$ are strictly increasing this implies that $\partial_\nu\phi_{\rho^*}$ and $\partial_\nu\phi_{\rho^{*,n}}^n$ are both constant on $\partial\Omega$. Proposition 4.9 implies for ϕ_{ρ^*} that this constant is strictly negative. The same is true for $\phi_{\rho^{*,n}}^n$ by the counterpart of Proposition 4.9 explained after (5.3). The fact that $\partial_\nu\phi_{\rho^*}$ is strictly negative on $\partial\Omega$ implies again by Proposition 4.9 that $\mathbb{R}^N \setminus \bar{\Omega}$ only consists of one connected (unbounded) component and $0 < \phi_{\rho^*} < \lambda^*$ in $\mathbb{R}^N \setminus \bar{\Omega}$. The same holds for $\phi_{\rho^{*,n}}^n$.

We can now list the properties of the equilibrium potentials. Using Theorem 4.7, Proposition 4.9 and Proposition 4.11, for every ball $B_r(0)$, with $r > 0$ large enough, the potential ϕ_{ρ^*} is a $C^{2,\alpha}(\overline{B_r(0)} \setminus \Omega)$ -solution of

$$(6.3) \quad \begin{cases} -\operatorname{div}(g(|\nabla\phi_{\rho^*}|)\nabla\phi_{\rho^*}) = 0, & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ 0 < \phi_{\rho^*} < \lambda^*, & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ \phi_{\rho^*} = \lambda^* & \text{on } \partial\Omega, \\ \partial_\nu\phi_{\rho^*} = \text{const.} < 0 & \text{on } \partial\Omega, \\ \lim_{|x| \rightarrow \infty} \phi_{\rho^*}(x) = 0 & \end{cases}$$

with $\sup_{\overline{B_r(0)}} |\nabla\phi_{\rho^*}| = 1 - \theta(r) < 1$ for some $\theta(r) \in (0, 1)$.

Similarly, as stated in Remark 5.5, for every ball $B_r(0)$, with $r > 0$ large enough, the potential $\phi_{\rho^{*,n}}^n$ is a $C^{2,\alpha}(\overline{B_r(0)} \setminus \Omega)$ -solution of

$$(6.4) \quad \begin{cases} -\operatorname{div}(g_n(|\nabla\phi_{\rho^{*,n}}^n|)\nabla\phi_{\rho^{*,n}}^n) = 0, & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ 0 < \phi_{\rho^{*,n}}^n < \lambda^* & \text{in } \mathbb{R}^N \setminus \bar{\Omega}, \\ \phi_{\rho^{*,n}}^n = \lambda^{*,n} & \text{on } \partial\Omega, \\ \partial_\nu\phi_{\rho^{*,n}}^n = \text{const.} < 0 & \text{on } \partial\Omega, \\ \lim_{|x| \rightarrow \infty} \phi_{\rho^{*,n}}^n(x) = 0. & \end{cases}$$

To finish the proof, let us first consider the approximated model (6.4). Here the function g_n and the problem (6.4) directly fulfill the assumptions of case (I) of [35, Theorem 1] which yields radial symmetry of $\phi_{\rho^{*,n}}^n$ and that Ω is a ball.

Now let us consider the Born-Infeld model (6.3). Here the function g is not defined on $[0, \infty)$ as required by [35, Theorem 1], but only on $[0, 1)$, where it also satisfies the requirement (I)₁, namely $g \in C^2[0, 1)$, $g(s) > 0$, $(g(s)s)' > 0$ on $[0, 1)$. If one investigates the proof of [35, Theorem 1] then it is clear that (I)₁ is required for the linearized equation fulfilled by $w(x) = \phi_{\rho^*}(x^\lambda) - \phi_{\rho^*}(x)$. Here x^λ stands for the reflection of x at a hyperplane $T_\lambda := \{x \in \mathbb{R}^N : x_1 = \lambda\}$ and λ denotes the geometrically critical position of the hyperplane. This linearized equation is determined as

$$(6.5) \quad \operatorname{div}(A(x)\nabla w) = 0 \quad \text{on } \{x \in \mathbb{R}^N : x_1 > \lambda \text{ and } x^\lambda \notin \bar{\Omega}\},$$

where

$$A(x)\nabla w := \int_0^1 \frac{d}{dt} [g(|\nabla w_t|)\nabla w_t] dt = \left[\int_0^1 \left(\frac{1}{\sqrt{1 - |\nabla w_t|^2}} \operatorname{Id} + \frac{\nabla w_t \otimes \nabla w_t}{(1 - |\nabla w_t|^2)^{3/2}} \right) dt \right] \nabla w$$

and $w_t(x) := t\phi_{\rho^*}(x^\lambda) + (1-t)\phi_{\rho^*}(x)$. Clearly, when x ranges in a compact subset K of the domain of definition of w then $|\nabla w_t(x)|$ takes values in a compact subset of $[0, 1)$ and $A(x)$ is uniformly positive definite on K . As a consequence, (6.5) is uniformly elliptic in K . This is sufficient for the proof of [35, Theorem 1] when applied to problem (6.3). In fact, this shows that it is enough to satisfy the requirement $(I)_1$ on $\text{range}(|\nabla\phi_{\rho^*}|)$ instead of $[0, \infty)$. As a result, we get that ϕ_{ρ^*} is radially symmetric and that Ω is a ball. This completes the proof of Theorem 6.1. \square

ACKNOWLEDGEMENTS

D. Bonheure is supported by MIS F.4508.14 (FNRS), PDR T.1110.14F (FNRS) & ARC AUWB-2012-12/17-ULB1- IAPAS. P. d'Avenia and A. Pomponio are partially supported by the group GNAMPA of INdAM, by FRA2016 of Politecnico di Bari and by PRIN 2017JPCAPN *Qualitative and quantitative aspects of nonlinear PDEs*. W. Reichel gratefully acknowledges funding by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation) Project-ID 258734477 SFB 1173.

REFERENCES

- [1] A.D. Alexandrov, *A characteristic property of spheres*, Ann. Mat. Pura Appl. (4) **58** (1962), 303–315. 22
- [2] R. Bartnik, L. Simon, *Spacelike hypersurfaces with prescribed boundary values and mean curvature*, Comm. Math. Phys. **87** (1982), 131–152. 3, 8, 13, 14
- [3] C. Bereanu, P. Jebelean, J. Mawhin, *The Dirichlet problem with mean curvature operator in Minkowski space – a variational approach*, Adv. Nonlinear Stud. **14** (2014), 315–326. 3
- [4] P. Billingsley, *Convergence of probability measures*, Wiley Series in Probability and Statistics: Probability and Statistics, John Wiley & Sons, Inc., New York, 1999. 11
- [5] D. Bonheure, A. Iacopetti, *Spacelike radial graphs of prescribed mean curvature in the Lorentz-Minkowski space*, Analysis & PDE **12** (2019), 1805–1842. 3
- [6] D. Bonheure, A. Iacopetti, *On the regularity of the minimizer of the electrostatic Born-Infeld energy*, Arch. Ration. Mech. Anal. **232** (2019), 697–725. 3
- [7] D. Bonheure, P. d'Avenia, A. Pomponio, *On the electrostatic Born-Infeld equation with extended charges*, Comm. Math. Phys. **346** (2016), 877–906. 2, 6, 7, 8, 9, 11, 12, 13, 14, 17, 21
- [8] M. Born, *Modified field equations with a finite radius of the electron*, Nature **132** (1933), 282. 5
- [9] M. Born, *On the quantum theory of the electromagnetic field*, Proc. Roy. Soc. London Ser. A **143** (1934), 410–437. 5
- [10] M. Born, L. Infeld, *Foundations of the new field theory*, Nature **132** (1933), 1004. 5
- [11] M. Born, L. Infeld, *Foundations of the new field theory*, Proc. Roy. Soc. London Ser. A **144** (1934), 425–451. 5
- [12] Y. Brenier, *Hydrodynamic Structure of the Augmented Born-Infeld Equations*, Arch. Rational Mech. Anal. **172** (2004), 65–91. 3
- [13] E. Calabi, *Examples of Bernstein problems for some nonlinear equations*, 1970 Global Analysis (Proc. Sympos. Pure Math., Vol. XV, Berkeley, Calif., 1968) pp. 223–230 Amer. Math. Soc., Providence, R.I. 3
- [14] H. Carley, M.K.-H. Kiessling, *Constructing graphs over \mathbb{R}^n with small prescribed mean-curvature*, Math. Phys. Anal. Geom. **18** (2015), no. 1, Art. 11. 3
- [15] S.Y. Cheng, S.T. Yau, *Maximal space-like hypersurfaces in the Lorentz-Minkowski spaces*, Ann. of Math. **104** (1976), 407–419. 3
- [16] P. Ebenfelt, D. Khavinson, H. Shapiro, *A free boundary problem related to single-layer potentials*, Ann. Acad. Sci. Fenn. Math. **27** (2002), no. 1, 21–46. 3
- [17] K. Ecker, *Area maximizing hypersurfaces in Minkowski space having an isolated singularity*, Manuscr. Math. **56** (1986), 375–397. 3
- [18] I. Ekeland, R. Témam, *Convex analysis and variational problems*, SIAM, Philadelphia (1999). 18
- [19] D. Fortunato, L. Orsina, L. Pisani, *Born-Infeld type equations for electrostatic fields*, J. Math. Phys. **43** (2002), 5698–5706. 4, 18
- [20] S. Gardiner, *An equilibrium measure characterization of circles*, Forum Math. **14** (2002), 953–954. 3
- [21] N. Garofalo, E. Sartori, *Symmetry in exterior boundary value problems for quasilinear elliptic equations via blow-up and a priori estimates*, Adv. Differential Equations **4** (1999), 137–161. 3
- [22] I.M. Gelfand, S.V. Fomin, *Calculus of variations*, Prentice-Hall, Inc., Englewood Cliffs, N.J. 1963. 4
- [23] C. Gerhardt, *H-Surfaces in Lorentzian Manifolds*, Commun. Math. Phys. **89** (1983), 523–553. 3

- [24] D. Gilbarg, N.S. Trudinger, *Elliptic partial differential equations of second order*, Springer-Verlag, Berlin (2001). 9, 16, 20
- [25] E. Hewitt, K. Stromberg, *Real and abstract analysis. A modern treatment of the theory of functions of a real variable*, Third printing. Graduate Texts in Mathematics, No. 25. Springer-Verlag, New York-Heidelberg, 1975. x+476 pp. 9
- [26] M.K.-H. Kiessling, *Some uniqueness results for stationary solutions to the Maxwell-Born-Infeld field equations and their physical consequences*, Phys. Lett. A **375** (2011), 3925–3930. 2
- [27] M.K.-H. Kiessling, *On the quasi-linear elliptic PDE $-\nabla \cdot (\nabla u / \sqrt{1 - |\nabla u|^2}) = 4\pi \sum_k a_k \delta_{s_k}$ in physics and geometry*, Comm. Math. Phys. **314** (2012), 509–523. 3, 6, 18
- [28] M.K.-H. Kiessling, *Correction to: On the quasi-linear elliptic PDE $-\nabla \cdot (\nabla u / \sqrt{1 - |\nabla u|^2}) = 4\pi \sum_k a_k \delta_{s_k}$ in physics and geometry*, Comm. Math. Phys. **364** (2018), 825–833. 3, 6
- [29] A.A. Klyachin, V.M. Miklyukov, *Existence of solutions with singularities for the maximal surface equation in Minkowski space*, Russ. Acad. Sci. Sb. Math. **80** (1995), 87–104. 3
- [30] N.S. Landkof, *Foundations of modern potential theory*, Springer Verlag, Berlin, 1972. 5
- [31] G.M. Lieberman, *Boundary regularity for solutions of degenerate elliptic equations*, Nonlinear Anal. **12** (1988), 1203–1219. 20
- [32] R. López, *Differential Geometry of curves and surfaces in Lorentz-Minkowski space*, International Electronic Journal of Geometry **7** (2014), 44–107. 3
- [33] K. Mallory, R. Van Gorder, K. Vajravelu, *Several classes of exact solutions to the 1+1 Born-Infeld equation*, Commun. Nonlinear Sci. Numer. Simul. **19** (2014), 1669–1674. 3
- [34] O. Mendez, W. Reichel, *Electrostatic characterization of spheres*, Forum Math. **12** (2000), 223–245. 3
- [35] W. Reichel, *Radially symmetry for an electrostatic, a capillarity and some fully nonlinear overdetermined problems on exterior domains*, Z. Anal. Anwendungen **15** (1996), 619–635. 3, 22, 23, 24
- [36] E.B. Saff, V. Totik, *Logarithmic potentials with external fields*, Springer Verlag, Berlin, 1997. 5
- [37] E. Schrödinger, *A new exact solution in non-linear optics. (Two-wave-system)*, Proc. Roy. Irish Acad. Sect. A. **49**, (1943), 59–66. 3
- [38] D. Serre, *Hyperbolicity of the Nonlinear Models of Maxwell's Equations*, Arch. Rational Mech. Anal. **172** (2004), 309–331. 3
- [39] J. Serrin, *A symmetry problem in potential theory*, Arch. Rational Mech. Anal. **43** (1971), 304–318. 22
- [40] J. Speck, *The nonlinear stability of the trivial solution to the Maxwell-Born-Infeld system*, J. Math. Phys. **53** (2012), 083703, 83 pp. 3
- [41] A. Treibergs, *Entire Spacelike Hypersurfaces of Constant Mean Curvature in Minkowski Space*, Invent. Math. **66** (1982), 39–56. 3
- [42] J. Wermer, *Potential theory*, Second edition, Lecture Notes in Mathematics **408**, Springer Verlag, Berlin, 1981. 5
- [43] Y. Yang, *Classical solutions in the Born-Infeld theory*, R. Soc. Lond. Proc. Ser. A Math. Phys. Eng. Sci. **456** (2000), 615–640. 3

(D. Bonheure)

DÉPARTEMENT DE MATHÉMATIQUE, UNIVERSITÉ LIBRE DE BRUXELLES,
CP 214, BOULEVARD DU TRIOMPHE, B-1050 BRUXELLES, BELGIUM
E-mail address: denis.bonheure@ulb.ac.be

(P. d'Avenia and A. Pomponio)

DIPARTIMENTO DI MECCANICA, MATEMATICA E MANAGEMENT, POLITECNICO DI BARI
VIA ORABONA 4, 70125 BARI, ITALY
E-mail address: pietro.davenia@poliba.it
E-mail address, Corresponding Author: alessio.pomponio@poliba.it

(W. Reichel)

INSTITUT FUER ANALYSIS, KARLSRUHER INSTITUT FUER TECHNOLOGIE
ENGLERSTRASSE 2, 76131 KARLSRUHE, GERMANY
E-mail address: wolfgang.reichel@kit.edu