



# Convergence of the Rosenau-Korteweg-de Vries Equation to the Korteweg-de Vries One

Giuseppe Maria Coclite<sup>1\*</sup>, Lorenzo di Ruvo<sup>2</sup>

<sup>1</sup>Department of Mechanics, Mathematics and Management, Polytechnic University of Bari, Via E. Orabona 4, 70125 Bari, Italy

<sup>2</sup>Department of Mathematics, University of Bari, via E. Orabona 4, 70125 Bari, Italy

Email: giuseppemaria.coclite@poliba.it

---

**Abstract:** The Rosenau-Korteweg-de Vries equation describes the wave-wave and wave-wall interactions. In this paper, we prove that, as the diffusion parameter is near zero, it coincides with the Korteweg-de Vries equation. The proof relies on deriving suitable a priori estimates together with an application of the Aubin-Lions Lemma.

**Keywords:** existence, uniqueness, stability, rosenau-korteweg-de vries-equation, korteweg-de vries equation

---

## 1. Introduction

The study of nonlinear wave phenomena is an important area of scientific research. In the last years several mathematical models describing wave behavior have been proposed. Some of them are the Korteweg-de Vries (KdV), the regularized long wave, the Rosenau, Rosenau-Kawahara, the Rosenau-KdV and the Rosenau-KdV-RLW equations (see [1-6]).

The KdV equation [7]

$$\partial_t u + au\partial_x u + \alpha\partial_x^3 u = 0, \quad a, \alpha \in \mathbb{R}, \quad (1)$$

has a wide range of applications, such as magnetic fluid waves, ion sound waves, and longitudinal astigmatic waves.

From a mathematical point of view, in [8-11], the Cauchy problem for (1) is studied, while in [12], the author reviewed the travelling wave solutions for (1). Moreover, in [13-15], the convergence of the solution of (1) to the unique entropy one of the Burgers equation is proven.

The KdV equation cannot describe the wave-wave and wave-wall interactions that appear in compact discrete systems. To overcome the shortcoming of KdV equation, Rosenau proposed the equation (see [3, 16-18])

$$\partial_t u + au\partial_x u + \alpha\partial_x^3 u + \beta^2\partial_t\partial_x^4 u = 0, \quad a, \alpha, \beta \in \mathbb{R}. \quad (2)$$

From a mathematical point of view, the well-posedness of the classical solution of the Cauchy problem of (2) is studied in [19-20]. In particular, in [19], the well-posedness of the solution of the (2) is proven, under the assumption:

$$u_0(x) \in H^2(\mathbb{R}). \quad (3)$$

In [21], Zuo discussed the solitary wave solutions of (2). In [3], a conservative linear finite difference scheme for the numerical solution of an initial-boundary value problem for (2) was considered. Esfahani in [16] and Razborova, Triki, Biswas in [22] studied the solitary solutions for (2) with the solitary ansatz method, and also gave two invariants for (2). In particular, in [22], the two types of soliton solutions were analyzed, one is a solitary wave and the other is a singular soliton. In [23], Zheng and Zhou proposed an average linear finite difference scheme for the numerical approximation of the solutions of the initial-boundary value problem for (2). Finally, in [13, 24], the convergence of the solution of (2) to the unique entropy one of the Burgers equation is proven.

Observe that, if we send  $\beta \rightarrow 0$  in (2), we obtain (1). Therefore, if  $\beta$  is near 0, the wave-wave and wave-wall interactions are described by (1).

The aim of this paper is to prove that, when  $\beta$  goes to 0, the solution of (2) converges to the unique one of (1). In other to do this, first, we augment (2) with the initial datum  $u_0(x)$ , on which assume (3). After, following [24] and fixed two small numbers  $0 < \varepsilon, \beta < 1$ , we consider the following Cauchy problem:

$$\begin{cases} \partial_t u_{\varepsilon, \beta} + \alpha u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} + \alpha \partial_x^3 u_{\varepsilon, \beta} + \beta^2 \partial_t \partial_x^4 u_{\varepsilon, \beta} = \varepsilon \partial_x^2 u_{\varepsilon, \beta}, & t > 0, x \in \mathbb{R}, \\ u_{\varepsilon, \beta}(0, x) = u_{\varepsilon, \beta, 0}(x), & x \in \mathbb{R}, \end{cases} \quad (4)$$

where  $u_{\varepsilon, \beta, 0}$  is a  $C^\infty$  approximation of  $u_0$  such that

$$\begin{aligned} \|u_{\varepsilon, \beta, 0}\|_{H^2(\mathbb{R})} &\leq \|u_0\|_{H^2(\mathbb{R})}, \quad \beta \|\partial_x^2 u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} \leq C_0, \\ \beta \|\partial_x^3 u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} + \varepsilon \beta \|\partial_x u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} + \varepsilon \beta \|\partial_x^2 u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} &\leq C_0, \\ \beta \|\partial_x^4 u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} + \varepsilon \|\partial_x u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} + \varepsilon \beta^{\frac{1}{2}} \|\partial_x^2 u_{\varepsilon, \beta, 0}\|_{L^2(\mathbb{R})} &\leq C_0, \end{aligned} \quad (5)$$

where  $C_0$  is a positive constant, independent on  $\varepsilon$  and  $\beta$ . In what follows we denote with  $C_0$  the constants which depend only on the initial data, and with  $C(T)$  the constants which depend also on  $T$ .

Equation (4) is known as the Rosenau-Korteweg-de Vries-Burgers equation (see [25-26]). The well-posedness of (4) can be proven as in [19]. In [25], the initial-boundary value problem for (4) is studied, while, in [26], the well-posedness of the periodic solutions for (4) is proven.

The main result of this paper is the following theorem.

**Theorem 1.1** Fix  $T > 0$  and consider (4). Assume (3) and (5). If

$$\beta = \mathcal{O}(\varepsilon^4), \quad (6)$$

then there exist two sequences  $\{\varepsilon_n\}_{n \in \mathbb{N}}, \{\beta_n\}_{n \in \mathbb{N}}$ , with  $\varepsilon_n, \beta_n \rightarrow 0$ , and a limit function

$$u \in L^\infty(0, T; H^2(\mathbb{R})), \quad (7)$$

which is the unique solution of (1). Moreover, if  $u_1$  and  $u_2$  are two solutions of (1), we have that

$$\|u_1(t, \cdot) - u_2(t, \cdot)\|_{L^2(\mathbb{R})} \leq e^{C(T)t} \|u_{1,0} - u_{2,0}\|_{L^2(\mathbb{R})}, \quad (8)$$

for some suitable  $C(T) > 0$ , and every  $0 \leq t \leq T$ .

The assumption (6) consists in

$$\beta \leq D^2 \varepsilon^4, \quad (9)$$

where  $D$  is a positive constant such that

$$D < \min \left\{ D_0, \frac{1}{|\alpha| \sqrt{2C_0}} \right\}, \quad (10)$$

where the positive constant  $D_0$  is the unique zero of the function  $F(X)$ , defined in (33). Note that in (6) we have  $\varepsilon^4$  and not  $\varepsilon^2$  because we have  $\beta^2$  and not  $\beta$  in (2).

From a physical point of view, Theorem 1.1, whose proof is based on the Aubin-Lions Lemma (see [27-30]), says that,

when  $\beta$  is near 0, Equations (2) or (4) coincides with (1). From a mathematical point of view, compared to [9], Theorem 1.1 gives a new method, to prove well-posedness of the Cauchy problem of (1).

The paper is organized as follows. In Section 2, we prove several a priori estimates on (4). Those play a key role in the proof of our main result, that is given in Section 3.

## 2. A priori estimates

This section is devoted to some a priori estimates on  $u_{\varepsilon, \beta}$ . We begin by proving the following result.

**Lemma 2.1** For each  $t \geq 0$ ,

$$\|u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \beta^2 \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \varepsilon \int_0^t \|\partial_x u_{\varepsilon, \beta}(s, \cdot)\|_{L^2(\mathbb{R})}^2 ds \leq C_0. \quad (11)$$

In particular, we have

$$\beta^2 \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})} \leq C_0. \quad (12)$$

Moreover,

$$\|u_{\varepsilon, \beta}(t, \cdot)\|_{L^\infty(\mathbb{R})} \leq C_0 \beta^{-\frac{1}{4}} \quad (13)$$

$$\|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^\infty(\mathbb{R})} \leq C_0 \beta^{-\frac{3}{4}}. \quad (14)$$

**Proof.** Arguing as in [24, Lemma 2.1] or [13, Lemma 2.2], by (5), we have (11).

We prove (12). Observe that

$$\beta \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 = \beta \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dx = -\beta \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} dx.$$

Therefore, by (11) and the Young inequality,

$$\begin{aligned} \beta \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 &\leq \int_{\mathbb{R}} |u_{\varepsilon, \beta}| \beta |\partial_x^2 u_{\varepsilon, \beta}| dx \\ &\leq \frac{1}{2} \|u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{\beta^2}{2} \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \leq C_0, \end{aligned}$$

which gives (12).

We prove (13). Thanks to the Hölder inequality,

$$\begin{aligned} u_{\varepsilon, \beta}^2(t, x) &= 2 \int_{-\infty}^x u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dy \leq 2 \int_{\mathbb{R}} |u_{\varepsilon, \beta}| |\partial_x u_{\varepsilon, \beta}| dx \\ &\leq 2 \|u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})} \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}. \end{aligned} \quad (15)$$

Consequently, by (11) and (12), we have that

$$\|u_{\varepsilon,\beta}(t,\cdot)\|_{L^\infty(\mathbb{R})}^2 \leq C_0 \beta^{-\frac{1}{2}},$$

which gives (13).

Finally, we prove (14). Again by the Hölder inequality,

$$\begin{aligned} (\partial_x u_{\varepsilon,\beta}(t,x))^2 &= 2 \int_{-\infty}^x \partial_x u_{\varepsilon,\beta} \partial_x^2 u_{\varepsilon,\beta} dy \leq 2 \int_{\mathbb{R}} |\partial_x u_{\varepsilon,\beta}| |\partial_x^2 u_{\varepsilon,\beta}| dx \\ &\leq 2 \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}. \end{aligned} \tag{16}$$

It follows from (11) and (12) that

$$\|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^\infty(\mathbb{R})}^2 \leq C_0 \beta^{-\frac{3}{2}},$$

which gives (14).

Following [24, Lemma 3.2], we prove the following result.

**Lemma 2.2** Fix  $T > 0$ . There exists a constant  $C_0 > 0$ , independent on  $\beta$  and  $\varepsilon$ , such that

$$\|u_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})} \leq C_0. \tag{17}$$

In particular, we have

$$\begin{aligned} \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} &\leq C_0, \\ \beta \|\partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} &\leq C_0, \\ \varepsilon \beta \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} &\leq C_0, \\ \varepsilon \beta \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})} &\leq C_0, \\ \varepsilon \int_0^t \|\partial_x^2 u_{\varepsilon,\beta}(s,\cdot)\|_{L^2(\mathbb{R})}^2 ds &\leq C_0, \\ \varepsilon \beta \int_0^t \|\partial_t u_{\varepsilon,\beta}(s,\cdot)\|_{L^2(\mathbb{R})}^2 ds &\leq C_0, \\ \varepsilon \beta^2 \int_0^t \|\partial_t \partial_x u_{\varepsilon,\beta}(s,\cdot)\|_{L^2(\mathbb{R})}^2 ds &\leq C_0, \\ \varepsilon \beta^3 \int_0^t \|\partial_t \partial_x^2 u_{\varepsilon,\beta}(s,\cdot)\|_{L^2(\mathbb{R})}^2 ds &\leq C_0, \\ \varepsilon \beta^4 \int_0^t \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(s,\cdot)\|_{L^2(\mathbb{R})}^2 ds &\leq C_0, \end{aligned} \tag{18}$$

for every  $0 \leq t \leq T$ . Moreover,

$$\|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^\infty(\mathbb{R})} \leq C_0 \beta^{-1}, \tag{19}$$

for every  $0 \leq t \leq T$ .

**Proof.** Let  $0 \leq t \leq T$ . Consider three real constants A, B and C, which will be specified. Multiplying (2) by

$$-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta},$$

we have that

$$\begin{aligned} & (-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta}) \partial_t u_{\varepsilon} \\ & + a(-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta}) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \\ & + \alpha(-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta}) \partial_x^3 u_{\varepsilon, \beta} \\ & + \beta^2(-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta}) \partial_t \partial_x^4 u_{\varepsilon} \\ & = \varepsilon(-2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta}) \partial_x^2 u_{\varepsilon}. \end{aligned} \tag{20}$$

Since

$$\begin{aligned} & \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon} \right) \partial_t u_{\varepsilon} dx \\ & = \frac{d}{dt} \left( \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{A}{3} \int_{\mathbb{R}} u_{\varepsilon, \beta}^3 dx \right) + B^2 \varepsilon \beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \\ & a \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta} \right) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dx \\ & = -2a \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} dx + B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t \partial_x u_{\varepsilon, \beta} dx \\ & + B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_{tx}^2 u_{\varepsilon, \beta} dx + C^2 a \varepsilon \beta \int_{\mathbb{R}} u_{\varepsilon} \partial_x u_{\varepsilon} \partial_t u_{\varepsilon} dx, \\ & \alpha \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta} \right) \partial_x^3 u_{\varepsilon, \beta} dx \\ & = -2A\alpha \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} dx + B^2 a \varepsilon \beta^2 dx + B^2 \alpha \varepsilon \beta^2 \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx - C^2 \alpha \varepsilon \beta \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx, \\ & \beta^2 \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon} \right) \partial_t \partial_x^4 u_{\varepsilon, \beta} dx \\ & = \beta^2 \frac{d}{dt} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - 2A\beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx + B^2 \beta^4 \varepsilon \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + C^2 \varepsilon \beta^3 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & = \beta^2 \frac{d}{dt} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - 4A\beta^2 \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon} \partial_t \partial_x u_{\varepsilon} dx \end{aligned}$$

$$\begin{aligned}
& +2A\beta^2 \int_{\mathbb{R}} u_{\varepsilon} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx + B^2 \beta^4 \varepsilon \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta^3 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \\
& \varepsilon \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta} \right) \partial_x^2 u_{\varepsilon, \beta} \\
& = -2\varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - 2A\varepsilon \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^2 dx - \frac{B^2 \varepsilon^2 \beta^2}{2} \frac{d}{dt} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& \varepsilon \int_{\mathbb{R}} \left( -2\partial_x^2 u_{\varepsilon, \beta} + Au_{\varepsilon, \beta}^2 - B^2 \varepsilon \beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} + C^2 \varepsilon \beta \partial_t u_{\varepsilon, \beta} \right) \partial_x^2 u_{\varepsilon, \beta} \\
& = -2\varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - 2A\varepsilon \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^2 dx - \frac{B^2 \varepsilon^2 \beta^2}{2} \frac{d}{dt} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& - \frac{C^2 \varepsilon^2 \beta}{2} \frac{d}{dt} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

an integration on  $\mathbb{R}$  of (20) gives

$$\begin{aligned}
& \frac{d}{dt} \left( \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{A}{3} \int_{\mathbb{R}} u_{\varepsilon, \beta}^3 dx + \beta^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& + \frac{d}{dt} \left( \frac{C^2 \varepsilon^2 \beta}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& + 2\varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& + B^2 \varepsilon \beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta^3 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + B^2 \varepsilon \beta^4 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& = 2(a + A\alpha) \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} dx - B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t \partial_x u_{\varepsilon, \beta} dx \\
& - B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_{tx}^2 u_{\varepsilon, \beta} dx - C^2 a \varepsilon \beta \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx \\
& - B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx - C^2 a \varepsilon \beta \int_{\mathbb{R}} \partial_x^3 u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx \\
& + 4A\beta^2 \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx - 2A\beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx - 2A\varepsilon \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^2 dx.
\end{aligned}$$

Taking  $A = -\frac{a}{\alpha}$ , we have that

$$\frac{d}{dt} \left( \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{a}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta}^3 dx + \beta^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right)$$

$$\begin{aligned}
& + \frac{d}{dt} \left( \frac{C^2 \varepsilon^2 \beta}{2} \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \right) \\
& + 2\varepsilon \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta \|\partial_t u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& + B^2 \varepsilon \beta^2 \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + C^2 \varepsilon \beta^3 \|\partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + B^2 \varepsilon \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& = -B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t \partial_x u_{\varepsilon, \beta} dx - B^2 a \varepsilon \beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_{tx}^2 u_{\varepsilon, \beta} dx \\
& - C^2 a \varepsilon \beta \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx - B^2 \alpha \varepsilon \beta^2 \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx \\
& + C^2 \alpha \varepsilon \beta \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx - \frac{4a\beta^2}{\alpha} \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx \\
& + \frac{2a\beta^2}{\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx + \frac{2a\varepsilon}{\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^2 dx. \tag{21}
\end{aligned}$$

Since  $0 < \varepsilon, \beta < 1$ , due to (10), (11), (13), (14) and the Young inequality,

$$\begin{aligned}
& B^2 |a| \varepsilon \beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 |\partial_t \partial_x u_{\varepsilon, \beta}| dx \\
& \leq \frac{B^2 a^2 \varepsilon \beta^2}{2} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^4 dx + \frac{B^2 \varepsilon \beta^2}{2} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq \frac{B^2 \varepsilon \beta^2}{2} \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^\infty(\mathbb{R})}^2 \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{2} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq \frac{B^2 \varepsilon \beta^2}{2} \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{2} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq C_0 B^2 D \varepsilon^3 \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{2} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq C_0 B^2 D \varepsilon \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{2} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2, \\
& B^2 |a| \varepsilon \beta^2 \int_{\mathbb{R}} |u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}| |\partial_{tx}^2 u_{\varepsilon, \beta}| dx \\
& \leq \frac{3B^2 a^2 \varepsilon \beta^2}{4} \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 (\partial_x^2 u_{\varepsilon, \beta})^2 dx + \frac{B^2 \varepsilon \beta^2}{3} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq \frac{3B^2 a^2 \varepsilon \beta^2}{4} \|u_{\varepsilon, \beta}(t, \cdot)\|_{L^\infty(\mathbb{R})}^2 \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{3} \|\partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2
\end{aligned}$$

$$\leq C_0 B^2 \varepsilon \beta^{\frac{3}{2}} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{3} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq C_0 B^2 D^3 \varepsilon^3 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{3} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq C_0 B^2 D^3 \varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{3} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$C^2 |a| \varepsilon \beta \int_{\mathbb{R}} |u_{\varepsilon} \partial_x u_{\varepsilon}| |\partial_t u_{\varepsilon}| dx$$

$$\leq \frac{C^2 a^2 \varepsilon \beta}{2} \int_{\mathbb{R}} u_{\varepsilon}^2 (\partial_x u_{\varepsilon})^2 dx + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{C^2 a^2 \varepsilon \beta}{2} \|u_{\varepsilon}(t, \cdot)\|_{L^{\infty}(\mathbb{R})}^2 \left\| \partial_x u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq C_0 C \varepsilon \beta^{\frac{1}{2}} \left\| \partial_x u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq C_0 C^2 D \varepsilon^2 \left\| \partial_x u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq C_0 C^2 D \varepsilon \left\| \partial_x u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t u_{\varepsilon}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$B^2 |\alpha| \varepsilon \beta^2 \int_{\mathbb{R}} |\partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x^3 u_{\varepsilon, \beta}| dx \leq \frac{B^2 \alpha^2 \varepsilon}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^4}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$C^2 |\alpha| \varepsilon \beta \int_{\mathbb{R}} |\partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x u_{\varepsilon, \beta}| dx \leq \frac{7C^4 \alpha^2 \varepsilon}{4B^2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{7} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$4 \left| \frac{a\beta^2}{\alpha} \right| \int_{\mathbb{R}} |\partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon}| |\partial_t \partial_x u_{\varepsilon, \beta}| dx \leq \frac{43a^2 \beta^2}{B^2 \alpha^2 \varepsilon} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 (\partial_x^2 u_{\varepsilon, \beta})^2 dx + \frac{B^2 \varepsilon \beta^2}{43} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{43a^2 \beta^2}{B^2 \alpha^2 \varepsilon} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^{\infty}(\mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{43} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{C_0 \beta^{\frac{1}{2}}}{B^2 \varepsilon} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{43} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{C_0 D \varepsilon}{B^2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{43} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$2 \left| \frac{a\beta^2}{\alpha} \right| \int_{\mathbb{R}} |u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x^2 u_{\varepsilon, \beta}| dx \leq \frac{2a^2 \beta}{\alpha^2 C^2 \varepsilon} \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 (\partial_x^2 u_{\varepsilon, \beta})^2 dx + \frac{C^2 \varepsilon \beta^3}{2} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$



$$\begin{aligned}
&\leq \frac{2a^2\beta}{\alpha^2 C^2 \varepsilon} \|u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta^3}{2} \|\partial_t \partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\
&\leq \frac{2a^2\beta^2}{\alpha^2 C^2 \varepsilon} \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta^3}{2} \|\partial_t \partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\
&\leq \frac{2a^2 D \varepsilon}{\alpha^2 C^2} \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta^3}{2} \|\partial_t \partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2.
\end{aligned}$$

Therefore, by (21),

$$\begin{aligned}
&\frac{d}{dt} \left( \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 - \frac{a}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon,\beta}^3 dx + \beta^2 \|\partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \right) \\
&+ \frac{d}{dt} \left( \frac{C^2 \varepsilon^2 \beta}{2} \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \right) \\
&+ \left( 2 - C_0 B^2 D^3 - \frac{B^2 \alpha^2}{2} - \frac{7C^4 \alpha^2}{4B^2} - \frac{C_0 D}{B^2} - \frac{2a^2 D}{\alpha^2 C^2} \right) \varepsilon \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\
&+ \frac{C^2 \varepsilon \beta}{2} \|\partial_t u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^2}{1806} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\
&+ \frac{C^2 \varepsilon \beta^3}{2} \|\partial_t \partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \beta^4 \varepsilon}{2} \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\
&\leq C_0 D (B^2 + C^2) \varepsilon \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + 2\varepsilon \left| \frac{a}{\alpha} \int_{\mathbb{R}} |u_{\varepsilon,\beta}| (\partial_x u_{\varepsilon,\beta})^2 dx \right. \\
&\leq C_0 \left( D (B^2 + C^2) + \|u_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})} \right) \varepsilon \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2. \tag{22}
\end{aligned}$$

We search  $B, C, D$  such that

$$2 - C_0 B^2 D^3 - \frac{B^2 \alpha^2}{2} - \frac{7C^4 \alpha^2}{4B^2} - \frac{C_0 D}{B^2} - \frac{2a^2 D}{\alpha^2 C^2} > 0, \tag{23}$$

that is

$$C_0 B^2 D^3 + \frac{\alpha^2 C_0 C^2 + 2a^2 B^2}{\alpha^2 B^2 C^2} D + \frac{B^2 \alpha^2}{2} + \frac{7C^4 \alpha^2}{4B^2} - 2 < 0. \tag{24}$$

We search  $B, C$  such that

$$\frac{B^2 \alpha^2}{2} + \frac{7C^4 \alpha^2}{4B^2} = 1, \tag{25}$$

that is

$$\frac{2\alpha^2 B^4 - 4B^2 + 7\alpha^2 C^4}{4B^2} = 0. \quad (26)$$

(26) is verified when

$$2\alpha^2 B^4 - 4B^2 + 7\alpha^2 C^4 = 0. \quad (27)$$

$B$  does exist if and only if

$$2 - 7\alpha^2 C^4 > 0. \quad (28)$$

Choosing

$$C^2 = \frac{1}{\sqrt{7}\alpha^2}, \quad (29)$$

(28) is verified. Moreover, thanks to (28) and (29), by (27), we have that

$$B^2 = \frac{2 - \sqrt{2}}{2\alpha^2} \text{ or } B^2 = \frac{2 + \sqrt{2}}{2\alpha^2}. \quad (30)$$

Thanks to (29) and (30), we can define the following positive constant:

$$k_1^2 := C_0 B^2, \quad k_2^2 := \frac{\alpha^2 C_0 C^2 + 2a^2 B^2}{\alpha^2 B^2 C^2}. \quad (31)$$

It follows from (24), (25) and (31) that

$$k_1^2 D^3 + k_2^2 D - 1 < 0. \quad (32)$$

Let consider the following function:

$$F(X) := k_1^2 X^3 + k_2^2 X - 1, \quad X \geq 0. \quad (33)$$

Observe that

$$F(0) = -1, \quad \lim_{X \rightarrow \infty} F(X) = \infty. \quad (34)$$

Moreover,

$$F'(X) = 3k_1^2 X^2 + k_2^2 > 0. \quad (35)$$

Then, it follows from (34) and (35) that the function  $F$  has an only zero  $D_0 > 0$ . Therefore, the following inequality

$$k_1^2 X^3 + k_2^2 X - 1 < 0,$$

is verified when

$$0 < X < D_0.$$

Taking  $X = D$ , we have (10).

Consequently, by (10), (22), (23), (29) and (30),

$$\begin{aligned} & \frac{d}{dt} \left( \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{a}{3\alpha} u_{\varepsilon, \beta}^3 dx + \beta^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + \frac{d}{dt} \left( \frac{C^2 \varepsilon^2 \beta}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + K_1^2 \varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta}{2} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{B^2 \varepsilon \beta^2}{1806} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon \beta^3}{2} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon \beta^4}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & \leq C_0 \left( K_2^2 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right) \varepsilon \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

for some constants  $K_1^2, K_2^2$ .

(5), (11) and an integration on  $(0, t)$  give

$$\begin{aligned} & \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{a}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta}^3 dx + \beta^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{C^2 \varepsilon^2 \beta}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + K_1^2 \varepsilon \int_0^t \left\| \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{C^2 \varepsilon \beta}{2} \int_0^t \left\| \partial_t u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\ & + \frac{B^2 \varepsilon \beta^2}{1806} \int_0^t \left\| \partial_t \partial_x u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{C^2 \varepsilon \beta^3}{2} \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\ & + \frac{B^2 \varepsilon \beta^4}{2} \int_0^t \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\ & \leq C_0 \left( K_2^2 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right) \varepsilon \int_0^t \left\| \partial_x u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\ & \leq C_0 \left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right). \end{aligned}$$

Therefore, by (11),

$$\begin{aligned} & \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{C^2 \varepsilon^2 \beta}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{B^2 \varepsilon^2 \beta^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + K_1^2 \varepsilon \int_0^t \left\| \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{C^2 \varepsilon \beta}{2} \int_0^t \left\| \partial_t u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{B^2 \varepsilon \beta^2}{1806} \int_0^t \left\| \partial_t \partial_x u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \end{aligned}$$

$$\begin{aligned}
& + \frac{C^2 \varepsilon \beta^3}{2} \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{B^2 \varepsilon \beta^4}{2} \int_0^t \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& \leq C_0 \left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right) + \frac{a}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta}^3 dx \\
& \leq C_0 \left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right) + \left| \frac{a}{3\alpha} \right| \int_{\mathbb{R}} |u_{\varepsilon, \beta}|^3 dx \\
& \leq C_0 \left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right) + \left| \frac{a}{3\alpha} \right| \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \|u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq C_0 \left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right). \tag{36}
\end{aligned}$$

We prove (17). Thanks to (11), (15) and (36).

$$u_{\varepsilon, \beta}^2(t, x) \leq C_0 \sqrt{\left( 1 + \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \right)}.$$

Therefore,

$$\|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})}^4 - C_0 \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} - C_0 \leq 0.$$

Arguing as in [31, Lemma 2.4], we have (17).

(18) follows from (17) and (36).

Finally, we prove (19). Due to (11), (18) and the Hölder inequality,

$$\begin{aligned}
(\partial_x^2 u_{\varepsilon, \beta}(t, x))^2 & = 2 \int_{-\infty}^x \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dy \leq 2 \int_{\mathbb{R}} |\partial_x^2 u_{\varepsilon, \beta}| |\partial_x^3 u_{\varepsilon, \beta}| dx \\
& \leq 2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \leq C_0 \beta^{-2}.
\end{aligned}$$

Hence,

$$\left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^\infty(\mathbb{R})}^2 \leq C_0 \beta^{-2},$$

which gives (19)

**Lemma 2.3** Fix  $T > 0$ . There exists a constant  $C(T) > 0$ , independent on  $\beta$  and  $\varepsilon$ , such that

$$\left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})} \leq C(T). \tag{37}$$

In particular, we have that

$$\left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \leq C(T),$$

$$\begin{aligned}
& \beta \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \leq C(T), \\
& \varepsilon \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \leq C(T), \\
& \varepsilon \beta^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})} \leq C(T), \\
& \varepsilon \int_0^t \left\| \partial_t u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \\
& \varepsilon \beta \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \\
& \varepsilon \beta^2 \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \\
& \varepsilon \beta^3 \int_0^t \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \\
& \varepsilon \int_0^t \left\| \partial_x^3 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \leq C(T), \tag{38}
\end{aligned}$$

for every  $0 \leq t \leq T$ .

**Proof.** Let  $0 \leq t \leq T$ . Consider four real constants  $E, F, G$  and  $H$ , which will be specified later. Multiplying (2) by

$$2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} + \varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta},$$

we have

$$\begin{aligned}
& (2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}) \partial_t u_{\varepsilon, \beta} \\
& + (\varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}) \partial_t u_{\varepsilon, \beta} \\
& + \alpha (2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \\
& + \alpha (\varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \\
& + \alpha (2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}) \partial_x^3 u_{\varepsilon, \beta} \\
& + \alpha (\varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}) \partial_x^3 u_{\varepsilon, \beta} \\
& + \beta^2 (2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}) \partial_t \partial_x^4 u_{\varepsilon, \beta} \\
& + \beta^2 (\varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}) \partial_t \partial_x^4 u_{\varepsilon, \beta} \\
& = \varepsilon (2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + Fu_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}) \partial_x^2 u_{\varepsilon, \beta}
\end{aligned}$$

$$+ \varepsilon(\varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}) \partial_x^2 u_{\varepsilon, \beta}. \quad (39)$$

Since

$$\begin{aligned} & \int_{\mathbb{R}} \left( 2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + F u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \right) \partial_t u_{\varepsilon, \beta} dx \\ &= \frac{d}{dt} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + E \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t u_{\varepsilon, \beta} dx + F \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx, \\ & \int_{\mathbb{R}} \left( \varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \right) \partial_t u_{\varepsilon, \beta} dx \\ &= \varepsilon G^2 \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \\ & a \int_{\mathbb{R}} \left( 2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + F u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \right) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dx \\ &= -2a \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_x^3 u_{\varepsilon, \beta} - 2a \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx + a(E - F) \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx \\ &= 5a \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} (\partial_x^2 u_{\varepsilon, \beta})^2 dx + a(E - F) \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx, \\ & a \int_{\mathbb{R}} \left( \varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \right) u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dx \\ &= a \varepsilon G^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx - a \varepsilon \beta H^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx, \\ & \alpha \int_{\mathbb{R}} \left( 2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + F u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \right) \partial_x^3 u_{\varepsilon, \beta} dx \\ &= -\alpha \left( 2E + \frac{F}{2} \right) \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx, \\ & \alpha \int_{\mathbb{R}} \left( \varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \right) \partial_x^3 u_{\varepsilon, \beta} dx \\ &= \varepsilon \alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx - \varepsilon \beta \alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\ & \beta^2 \int_{\mathbb{R}} \left( 2\partial_x^4 u_{\varepsilon, \beta} + E(\partial_x u_{\varepsilon, \beta})^2 + F u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \right) \partial_t \partial_x^4 u_{\varepsilon, \beta} dx \\ &= \beta^2 \frac{d}{dt} \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \beta^2 (2E + F) \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx \\ & - \beta^2 F \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx, \\ & \beta^2 \int_{\mathbb{R}} \left( \varepsilon G^2 \partial_t u_{\varepsilon, \beta} - \varepsilon \beta H^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \right) \partial_t \partial_x^4 u_{\varepsilon, \beta} dx \end{aligned}$$

$$\begin{aligned}
&= \varepsilon\beta^2 G^2 \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon\beta^3 H^2 \left\| \partial_t \partial_x^3 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \\
&= \varepsilon \int_{\mathbb{R}} \left( 2\partial_x^4 u_{\varepsilon,\beta} + E(\partial_x u_{\varepsilon,\beta})^2 + F u_{\varepsilon,\beta} \partial_x^2 u_{\varepsilon,\beta} \right) \partial_x^2 u_{\varepsilon,\beta} dx \\
&= -2\varepsilon \left\| \partial_x^3 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon F \int_{\mathbb{R}} u_{\varepsilon,\beta} (\partial_x^2 u_{\varepsilon,\beta})^2 dx, \\
&\varepsilon \int_{\mathbb{R}} \left( \varepsilon G^2 \partial_t u_{\varepsilon,\beta} - \varepsilon\beta H^2 \partial_t \partial_x^2 u_{\varepsilon,\beta} \right) \partial_x^2 u_{\varepsilon,\beta} dx \\
&= -\frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right),
\end{aligned}$$

an integration on  $\mathbb{R}$  of (39) gives

$$\begin{aligned}
&\frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_x^4 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&+ E \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 \partial_t u_{\varepsilon,\beta} dx + F \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x^2 u_{\varepsilon,\beta} \partial_t u_{\varepsilon,\beta} dx \\
&+ \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&+ \varepsilon G^2 \left\| \partial_t u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon\beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&+ \varepsilon\beta^2 G^2 \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon\beta^3 H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\varepsilon \left\| \partial_x^3 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&= \left( 2\alpha E + \frac{\alpha F}{2} - 5a \right) \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} (\partial_x^2 u_{\varepsilon,\beta})^2 dx - a(E - F) \int_{\mathbb{R}} u_{\varepsilon,\beta} (\partial_x u_{\varepsilon,\beta})^3 dx \\
&- a\varepsilon G^2 \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x u_{\varepsilon,\beta} \partial_t u_{\varepsilon,\beta} dx + a\varepsilon\beta H^2 \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x u_{\varepsilon,\beta} \partial_t \partial_x^2 u_{\varepsilon,\beta} dx \\
&- \varepsilon\alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon,\beta} \partial_x^3 u_{\varepsilon,\beta} dx - \varepsilon\beta\alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon,\beta} \partial_x^3 u_{\varepsilon,\beta} dx \\
&+ \beta^2 (2E + F) \int_{\mathbb{R}} \partial_x u_{\varepsilon,\beta} \partial_x^2 u_{\varepsilon,\beta} \partial_t \partial_x^3 u_{\varepsilon,\beta} dx + \beta^2 F \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x^3 u_{\varepsilon,\beta} \partial_t \partial_x^3 u_{\varepsilon,\beta} dx \\
&+ \varepsilon F \int_{\mathbb{R}} u_{\varepsilon,\beta} (\partial_x^2 u_{\varepsilon,\beta})^2 dx. \tag{40}
\end{aligned}$$

Observe that

$$\begin{aligned}
&E \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 \partial_t u_{\varepsilon,\beta} dx + F \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x^2 u_{\varepsilon,\beta} \partial_t u_{\varepsilon,\beta} dx \\
&= (E - F) \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 \partial_t u_{\varepsilon,\beta} dx - F \int_{\mathbb{R}} u_{\varepsilon,\beta} \partial_x u_{\varepsilon,\beta} \partial_t \partial_x u_{\varepsilon,\beta} dx
\end{aligned}$$

$$= (E - F) \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t u_{\varepsilon, \beta} dx - \frac{F}{2} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_t ((\partial_x u_{\varepsilon, \beta})^2) dx.$$

Consequently, by (40),

$$\begin{aligned} & \frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + (E - F) \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t u_{\varepsilon, \beta} dx - \frac{F}{2} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_t ((\partial_x u_{\varepsilon, \beta})^2) dx \\ & + \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + \varepsilon G^2 \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \varepsilon \beta^2 G^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta^3 H^2 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + 2\varepsilon \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & = \left( 2\alpha E + \frac{\alpha F}{2} - 5a \right) \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} (\partial_x^2 u_{\varepsilon, \beta})^2 dx - a(E - F) \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx \\ & + a\varepsilon G^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx + a\varepsilon \beta H^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx \\ & - \varepsilon \alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx - \varepsilon \beta \alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\ & - \varepsilon \alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx - \varepsilon \beta \alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\ & + \beta^2 (2E + F) \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx + \beta^2 F \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx \\ & + \varepsilon F \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x^2 u_{\varepsilon, \beta})^2 dx. \end{aligned} \tag{41}$$

We search  $E, F$  such that

$$E - F = -\frac{F}{2}, \quad 2\alpha E + \frac{\alpha F}{2} - 5a = 0. \tag{42}$$

Since

$$(E, F) = \left( \frac{5a}{3\alpha}, \frac{10a}{3\alpha} \right)$$

is the unique solution of (42), it follows from (41) that



$$\begin{aligned}
& \frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& - \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t u_{\varepsilon, \beta} dx - \frac{5a}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_t ((\partial_x u_{\varepsilon, \beta})^2) dx \\
& + \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& + \varepsilon G^2 \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& + \varepsilon \beta^2 G^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta^3 H^2 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& + 2\varepsilon \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& = \frac{5a^2}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx - a\varepsilon G^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx \\
& + a\varepsilon \beta H^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx - \varepsilon \alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\
& + \varepsilon \beta \alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx - \varepsilon \alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\
& - \varepsilon \beta \alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx + \frac{20a\beta^2}{3\alpha} \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx \\
& + \frac{10a\beta^2}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx + \frac{10a\varepsilon}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x^2 u_{\varepsilon, \beta})^2 dx,
\end{aligned}$$

that is

$$\begin{aligned}
& \frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 u_{\varepsilon, \beta} dx + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& + \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
& + \varepsilon G^2 \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& + \varepsilon \beta^2 G^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta^3 H^2 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\varepsilon \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& = \frac{5a^2}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x u_{\varepsilon, \beta})^3 dx - a\varepsilon G^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx
\end{aligned}$$

$$\begin{aligned}
& +a\varepsilon\beta H^2 \int_{\mathbb{R}} u_{\varepsilon} \partial_x u_{\varepsilon} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx - \varepsilon\alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\
& +\varepsilon\beta\alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx - \varepsilon\alpha G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \\
& -\varepsilon\beta\alpha H^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx + \frac{20a\beta^2}{3\alpha} \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx \\
& + \frac{10a\beta^2}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx + \frac{10a\varepsilon}{3\alpha} \int_{\mathbb{R}} u_{\varepsilon, \beta} (\partial_x^2 u_{\varepsilon, \beta})^2 dx. \tag{43}
\end{aligned}$$

Since  $0 < \varepsilon, \beta < 1$ , due to (17), (18), (9) and the Young inequality,

$$\begin{aligned}
\left| \frac{5a^2}{3\alpha} \int_{\mathbb{R}} |u_{\varepsilon, \beta}| |\partial_x u_{\varepsilon, \beta}|^3 dx \right| & \leq \left| \frac{5a^2}{3\alpha} \right| \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \int_{\mathbb{R}} |\partial_x u_{\varepsilon, \beta}|^3 dx \\
& \leq C_0 \|\partial_x u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq C_0 \|\partial_x u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \\
& \leq C_0 + \frac{1}{2} \|\partial_x u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})}^2 \\
& \leq C_0 \left( 1 + \|\partial_x u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})}^2 \right),
\end{aligned}$$

$$\begin{aligned}
|a| \varepsilon G^2 \int_{\mathbb{R}} |u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta}| |\partial_t u_{\varepsilon, \beta}| dx & = 2\varepsilon G^2 \int_{\mathbb{R}} \left| \frac{\sqrt{3} a u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta}}{2} \right| \left| \frac{\partial_t u_{\varepsilon, \beta}}{\sqrt{3}} \right| dx \\
& \leq \frac{3\varepsilon a^2 G^2}{4} \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 (\partial_x u_{\varepsilon, \beta})^2 dx + \frac{\varepsilon G^2}{3} \|\partial_t u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq \frac{3\varepsilon a^2 G^2}{4} \|u_{\varepsilon}\|_{L^\infty((0, T) \times \mathbb{R})}^2 \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon G^2}{3} \|\partial_t u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq C_0 G^2 + \frac{\varepsilon G^2}{3} \|\partial_t u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

$$\begin{aligned}
|a| \varepsilon\beta H^2 \int_{\mathbb{R}} |u_{\varepsilon} \partial_x u_{\varepsilon}| |\partial_t \partial_x^2 u_{\varepsilon, \beta}| dx & = 2\varepsilon H^2 \int_{\mathbb{R}} \left| \frac{\sqrt{3} a u_{\varepsilon} \partial_x u_{\varepsilon}}{2} \right| \left| \frac{\beta_t \partial_x^2 u_{\varepsilon, \beta}}{\sqrt{3}} \right| \\
& \leq \frac{\varepsilon 3a^2 H^2}{4} \int_{\mathbb{R}} u_{\varepsilon}^2 (\partial_x u_{\varepsilon})^2 dx + \frac{\varepsilon\beta^2 H^2}{3} \|\partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \\
& \leq \frac{\varepsilon 3a^2 H^2}{4} \|u_{\varepsilon}\|_{L^\infty((0, T) \times \mathbb{R})}^2 \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon\beta^2 H^2}{3} \|\partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2
\end{aligned}$$

$$\leq C_0 H^2 + \frac{\varepsilon \beta^2 H^2}{3} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$\varepsilon |\alpha| G^2 \int_{\mathbb{R}} \partial_t u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta} dx \leq \frac{\varepsilon G^2}{2} \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \alpha^2 G^2}{2} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$\begin{aligned} \varepsilon \beta |\alpha| H^2 \int_{\mathbb{R}} |\partial_t \partial_x^2 u_{\varepsilon, \beta}| |\partial_x^3 u_{\varepsilon, \beta}| dx &= \varepsilon H^2 \int_{\mathbb{R}} \left| \beta \partial_t \partial_x^2 u_{\varepsilon, \beta} \right| \left| \alpha \partial_x^3 u_{\varepsilon, \beta} \right| dx \\ &\leq \frac{\varepsilon \beta^2 H^2}{2} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \alpha^2 H^2}{2} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

$$\left| \frac{20a\beta^2}{3\alpha} \int_{\mathbb{R}} |\partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x^3 u_{\varepsilon, \beta}| dx \right|$$

$$= \int_{\mathbb{R}} \left| \frac{20a\beta^2 \frac{1}{3\alpha H \sqrt{\varepsilon}} \partial_x u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}}{\beta^2 \sqrt{\varepsilon} H \partial_t \partial_x^3 u_{\varepsilon, \beta}} \right| dx$$

$$\leq \frac{200a^2 \beta}{9\alpha^2 H^2 \varepsilon} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 (\partial_x^2 u_{\varepsilon, \beta})^2 dx + \frac{\varepsilon \beta^3 H^2}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{200a^2 \beta}{9\alpha^2 H^2 \varepsilon} \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{200a^2 D^2 \varepsilon^4}{9\alpha^2 H^2 \varepsilon} \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{200a^2 D^2 \varepsilon^3}{9\alpha^2 H^2} \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{200a^2 D^2 \varepsilon}{9\alpha^2 H^2} \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{2} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,$$

$$\left| \frac{10a\beta^2}{3\alpha} \int_{\mathbb{R}} |u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta}| |\partial_t \partial_x^3 u_{\varepsilon, \beta}| dx \right| = 2 \int_{\mathbb{R}} \left| \frac{5\sqrt{3}a\beta^2 u_{\varepsilon, \beta} \partial_x^3 u_{\varepsilon, \beta}}{3\alpha H \sqrt{\varepsilon}} \right| \left| \frac{\beta^2 \sqrt{\varepsilon} H \partial_t \partial_x^3 u_{\varepsilon, \beta}}{\sqrt{3}} \right| dx$$

$$\leq \frac{75a^2 \beta}{9\alpha^2 H^2 \varepsilon} \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 (\partial_x^3 u_{\varepsilon, \beta})^2 dx + \frac{\varepsilon \beta^3 H^2}{3} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\leq \frac{75a^2 \beta}{9\alpha^2 H^2 \varepsilon} \left\| u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{3} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2$$

$$\begin{aligned}
&\leq \frac{C_0 D^2 \varepsilon^4}{H^2 \varepsilon} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{3} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\leq \frac{C_0 D^2 \varepsilon^3}{H^2} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{3} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\leq \frac{C_0 D^2 \varepsilon}{H^2} \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{3} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2,
\end{aligned}$$

$$\begin{aligned}
\left| \frac{10a}{3\alpha} \right| \varepsilon \int_{\mathbb{R}} |u_{\varepsilon, \beta}| (\partial_x^2 u_{\varepsilon, \beta})^2 dx &\leq \left| \frac{10a}{3\alpha} \right| \varepsilon \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\leq C_0 \varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2.
\end{aligned}$$

It follows from (43) that

$$\begin{aligned}
&\frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 u_{\varepsilon, \beta} dx + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&+ \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\
&+ \frac{\varepsilon G^2}{6} \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&+ \frac{\varepsilon \beta^2 G^2}{6} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{6} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&+ \left( 2 - \frac{\alpha^2 G^2}{2} - \frac{\alpha^2 H^2}{2} - \frac{C_0 D^2}{H^2} \right) \varepsilon \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
&\leq C_0 \left( 1 + \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \right) + C_0 G^2 + C_0 H^2 \\
&+ \frac{200a^2 D^2 \varepsilon}{9\alpha^2 H^2} \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + C_0 \varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2. \tag{44}
\end{aligned}$$

We search  $G, H$  such that

$$2 - \frac{\alpha^2 G^2}{2} - \frac{\alpha^2 H^2}{2} - \frac{C_0 D^2}{H^2} > 0. \tag{45}$$

Choosing

$$G^2 = \frac{2}{\alpha^2} \tag{46}$$

(45) reads

$$1 - \frac{\alpha^2 H^2}{2} - \frac{C_0 D^2}{H^2} > 0.$$

Therefore, we search  $H$  such that

$$\alpha^2 H^4 - 2H^2 + 2C_0 D^2 < 0. \tag{47}$$

Thanks to (10), there exists  $0 < H_1^2 < H_2^2$ , such that choosing

$$H_1^2 < H^2 < H_2^2, \tag{48}$$

(47) is verified.

Therefore, by (10), (44), (46), (47) and (48), we have

$$\begin{aligned} & \frac{d}{dt} \left( \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 u_{\varepsilon, \beta} dx + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + \frac{d}{dt} \left( \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \right) \\ & + \frac{\varepsilon G^2}{6} \left\| \partial_t u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \varepsilon \beta H^2 \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{\varepsilon \beta^2 G^2}{6} \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon \beta^3 H^2}{6} \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + K_3^2 \varepsilon \left\| \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & \leq C_0 \left( 1 + \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \right) + C_0 \varepsilon \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{200a^2 D^2 \varepsilon}{9\alpha^2 H^2} \left\| \partial_x u_\varepsilon \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \end{aligned} \tag{49}$$

where  $K_3^2$  is an appropriate positive constant.

It follows from (5), (18) and an integration on  $(0, t)$  that

$$\begin{aligned} & \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 - \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 u_{\varepsilon, \beta} dx + \beta^2 \left\| \partial_x^4 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & + \frac{\varepsilon G^2}{6} \int_0^t \left\| \partial_t u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \varepsilon \beta H^2 \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon, \beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \end{aligned}$$

$$\begin{aligned}
& + \frac{\varepsilon\beta^2 G^2}{6} \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{\varepsilon\beta^3 H^2}{6} \int_0^t \left\| \partial_t \partial_x^3 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& + K_3^2 \varepsilon \int_0^t \left\| \partial_x^3 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& \leq C_0 + C_0 \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right) t + C_0 \varepsilon \int_0^t \left\| \partial_x^2 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& + \frac{200a^2 D^2 \varepsilon}{9\alpha^2 H^2} \left\| \partial_x u_\varepsilon \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \int_0^t \left\| \partial_x^2 u_\varepsilon(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& \leq C(T) \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right).
\end{aligned}$$

Consequently, by (17) and (18), we get

$$\begin{aligned}
& \left\| \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_x^4 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 G^2}{2} \left\| \partial_x u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \frac{\varepsilon^2 \beta H^2}{2} \left\| \partial_x^2 u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& + \frac{\varepsilon G^2}{6} \int_0^t \left\| \partial_t u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \varepsilon \beta H^2 \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + \frac{\varepsilon \beta^2 G^2}{6} \int_0^t \left\| \partial_t \partial_x^2 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& + \frac{\varepsilon \beta^3 H^2}{6} \int_0^t \left\| \partial_t \partial_x^3 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds + K_3^2 \varepsilon \int_0^t \left\| \partial_x^3 u_{\varepsilon,\beta}(s, \cdot) \right\|_{L^2(\mathbb{R})}^2 ds \\
& \leq C(T) \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right) + \frac{5a}{3\alpha} \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 u_{\varepsilon,\beta} dx \\
& \leq C(T) \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right) + \left| \frac{5a}{3\alpha} \right| \int_{\mathbb{R}} (\partial_x u_{\varepsilon,\beta})^2 |u_{\varepsilon,\beta}| dx \\
& \leq C(T) \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right) + \left| \frac{5a}{3\alpha} \right| \left\| u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})} \left\| \partial_x u_{\varepsilon,\beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\
& \leq C(T) \left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right). \tag{50}
\end{aligned}$$

We prove (37). Thanks to (16), (18) and (50),

$$(\partial_x u_{\varepsilon,\beta}(t, x))^2 \leq C(T) \sqrt{\left( 1 + \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 \right)}.$$

Hence, we have

$$\left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^4 - C(T) \left\| \partial_x u_{\varepsilon,\beta} \right\|_{L^\infty((0,T) \times \mathbb{R})}^2 - C(T) \leq 0,$$

which gives (37).

Finally, (37) and (50) give (38).

**Lemma 2.4** Fix  $T > 0$ . There exists a constant  $C(T) > 0$ , independent on  $\beta$  and  $\varepsilon$ , such that

$$\frac{\beta^2}{6} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^4 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \leq C(T), \quad (51)$$

for every  $0 \leq t \leq T$ .

**Proof.** Let  $0 \leq t \leq T$ . Multiplying (2) by  $-2\beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta}$ , we have that

$$\begin{aligned} & -2\beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} - 2\beta^4 \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} \\ & = 2\beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} - 2\beta^2 \alpha \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} - 2\varepsilon \beta^2 \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta}. \end{aligned} \quad (52)$$

Observe that

$$\begin{aligned} & -2\beta^2 \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_t u_{\varepsilon, \beta} dx - 2\beta^4 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx = 2\beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\beta^4 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \\ & 2\beta^2 \int_{\mathbb{R}} \partial_t \partial_x^2 u_{\varepsilon, \beta} u_{\varepsilon, \beta} \partial_x u_{\varepsilon, \beta} dx = -2\beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t \partial_x u_{\varepsilon, \beta} dx - 2\beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx, \\ & -2\beta^2 \alpha \int_{\mathbb{R}} \partial_x^3 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx = 2\beta^2 \alpha \int_{\mathbb{R}} \partial_x^4 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx, \\ & -2\varepsilon \beta^2 \int_{\mathbb{R}} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x^2 u_{\varepsilon, \beta} dx = 2\varepsilon \beta^2 \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx. \end{aligned} \quad (53)$$

Consequently, an integration on  $\mathbb{R}$  of (52) gives

$$\begin{aligned} & 2\beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + 2\beta^4 \left\| \partial_t \partial_x^3 u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & - 2\beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 \partial_t \partial_x u_{\varepsilon, \beta} dx - 2\beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx \\ & + 2\beta^2 \alpha \int_{\mathbb{R}} \partial_x^4 u_{\varepsilon, \beta} \partial_t \partial_x u_{\varepsilon, \beta} dx + 2\varepsilon \beta^2 \int_{\mathbb{R}} \partial_x u_{\varepsilon, \beta} \partial_t \partial_x^3 u_{\varepsilon, \beta} dx. \end{aligned} \quad (54)$$

Since  $0 < \varepsilon, \beta < 1$ , thanks to (11), (17), (18), (37), (38) and the Young inequality,

$$\begin{aligned} & 2\beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^2 |\partial_t \partial_x u_{\varepsilon, \beta}| dx \leq \beta^2 \int_{\mathbb{R}} (\partial_x u_{\varepsilon, \beta})^4 dx + \beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & \leq \left\| \partial_x u_{\varepsilon, \beta} \right\|_{L^\infty((0, T) \times \mathbb{R})}^2 \left\| \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 + \beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \\ & \leq C(T) + \beta^2 \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

$$\begin{aligned} & 2\beta^2 \int_{\mathbb{R}} |u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x u_{\varepsilon, \beta}| dx = \beta^2 \int_{\mathbb{R}} |2u_{\varepsilon, \beta} \partial_x^2 u_{\varepsilon, \beta}| |\partial_t \partial_x u_{\varepsilon, \beta}| dx \\ & \leq 2\beta^2 \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 (\partial_x^2 u_{\varepsilon, \beta})^2 dx + \frac{\beta^2}{2} \left\| \partial_t \partial_x u_{\varepsilon, \beta}(t, \cdot) \right\|_{L^2(\mathbb{R})}^2 \end{aligned}$$

$$\begin{aligned} &\leq 2\beta^2 \|u_{\varepsilon,\beta}\|_{L^\infty((0,T)\times\mathbb{R})}^2 \|\partial_x^2 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{\beta^2}{2} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\ &\leq C_0 + \frac{\beta^2}{2} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

$$\begin{aligned} 2\beta^2 |\alpha| \int_{\mathbb{R}} |\partial_x^4 u_{\varepsilon,\beta}| |\partial_t \partial_x u_{\varepsilon,\beta}| dx &= 2\beta^2 \int_{\mathbb{R}} \left| \sqrt{3}\alpha \partial_x^4 u_{\varepsilon,\beta} \right| \left| \frac{\partial_t \partial_x u_{\varepsilon,\beta}}{\sqrt{3}} \right| dx \\ &\leq 3\alpha^2 \beta^2 \|\partial_x^4 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \frac{\beta^2}{3} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\ &\leq C(T) + \frac{\beta^2}{3} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2, \end{aligned}$$

$$\begin{aligned} 2\varepsilon\beta^2 \int_{\mathbb{R}} |\partial_x u_{\varepsilon,\beta}| |\partial_t \partial_x^3 u_{\varepsilon,\beta}| dx &\leq \varepsilon^2 \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\ &\leq \|\partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \\ &\leq C_0 + \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2. \end{aligned}$$

It follows from (54) that

$$\frac{\beta^2}{6} \|\partial_t \partial_x u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 + \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon,\beta}(t,\cdot)\|_{L^2(\mathbb{R})}^2 \leq C(T),$$

which gives (51).

### 3. Proof of theorem 1.1

In this section, we prove Theorem 1.1. We begin by proving the following result.

**Lemma 3.1** Fix  $T > 0$  and assume (6),

the sequence  $\{u_{\varepsilon,\beta}\}_{\varepsilon,\beta>0}$  is compact in  $L_{loc}^2((0,\infty)\times\mathbb{R})$ . (55)

Consequently, there exists a subsequence  $\{u_{\varepsilon_k,\beta_k}\}_{k\in\mathbb{N}}$  of  $\{u_{\varepsilon,\beta}\}_{\varepsilon,\beta>0}$  and  $u \in L_{loc}^2((0,\infty)\times\mathbb{R})$  such that, for each compact subset  $K$  of  $(0,\infty)\times\mathbb{R}$ ,

$$u_{\varepsilon_k,\beta_k} \rightarrow u \text{ in } L^2(K) \text{ and a.e.}, \tag{56}$$

Moreover,  $u$  is a solution of (1) and (7) holds.

**Proof.** We begin by proving (55). To prove (55), we rely on the Aubin-Lions Lemma (see [27-30]). We recall that

$$H_{loc}^1(\mathbb{R}) \hookrightarrow L_{loc}^2(\mathbb{R}) \hookrightarrow H_{loc}^{-1}(\mathbb{R}),$$

where the first inclusion is compact and the second is continuous. Owing to the Aubin-Lions Lemma [30], to prove (55), it suffices to show that



$$\{u_{\varepsilon, \beta}\}_{\varepsilon, \beta > 0} \text{ is uniformly bounded in } L^2(0, T; H_{loc}^1(\mathbb{R})), \quad (57)$$

$$\{\partial_t u_{\varepsilon, \beta}\}_{\varepsilon, \beta > 0} \text{ is uniformly bounded in } L^2(0, T; H_{loc}^{-1}(\mathbb{R})). \quad (58)$$

We prove (57). Thanks to (6) and Lemmas 2.1, 2.2 and 2.3,

$$\|u_{\varepsilon, \beta}(t, \cdot)\|_{H^2(\mathbb{R})}^2 = \|u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \|\partial_x u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 + \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2(\mathbb{R})}^2 \leq C(T).$$

Therefore,

$$\{u_{\varepsilon, \beta}\}_{\varepsilon, \beta > 0} \text{ is uniformly bounded in } L^\infty(0, T; H^2(\mathbb{R})),$$

which gives (57).

We prove (58). We begin by observing that, by (4),

$$\partial_t u_{\varepsilon, \beta} = \partial_x \left( -\frac{a}{2} u_{\varepsilon, \beta}^2 - \alpha \partial_x^2 u_{\varepsilon, \beta} - \beta^2 \partial_t \partial_x^3 u_{\varepsilon, \beta} + \varepsilon \partial_x u_{\varepsilon, \beta} \right). \quad (59)$$

We have that

$$\frac{a^2}{4} \|u_{\varepsilon, \beta}^2\|_{L^2((0, T) \times \mathbb{R})}^2 \leq C(T). \quad (60)$$

Thanks to (6) and Lemmas 2.1 and 2.2,

$$\begin{aligned} \frac{a^2}{4} \int_0^T \int_{\mathbb{R}} u_{\varepsilon, \beta}^4 dt dx &\leq \frac{a^2}{4} \|u_{\varepsilon, \beta}\|_{L^\infty((0, T) \times \mathbb{R})}^2 \int_0^T \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 dt dx \\ &\leq C(T) \int_0^T \int_{\mathbb{R}} u_{\varepsilon, \beta}^2 dt dx \leq C(T). \end{aligned}$$

Observe that, since  $0 < \varepsilon < 1$ , thanks to (6) and Lemmas 2.1, 2.3 and 2.41

$$\varepsilon^2 \|\partial_x u_{\varepsilon, \beta}\|_{L^2((0, T) \times \mathbb{R})}^2, \alpha^2 \|\partial_x^2 u_{\varepsilon, \beta}(t, \cdot)\|_{L^2((0, T) \times \mathbb{R})}^2, \beta^4 \|\partial_t \partial_x^3 u_{\varepsilon, \beta}\|_{L^2((0, T) \times \mathbb{R})}^2 \leq C(T). \quad (61)$$

Therefore, by (60) and (61),

$$\left\{ \partial_x \left( -\frac{a}{2} u_{\varepsilon, \beta}^2 - \alpha \partial_x^2 u_{\varepsilon, \beta} - \beta^2 \partial_t \partial_x^3 u_{\varepsilon, \beta} + \varepsilon \partial_x u_{\varepsilon, \beta} \right) \right\}_{\varepsilon, \beta > 0} \text{ is bounded in } H^1((0, T) \times \mathbb{R}).$$

Thanks to the Aubin-Lions Lemma, (55) and (56) hold.

Consequently,  $u$  is solution of (1) and (7) holds.

Following [32, Theorem 1.1], we prove Theorem 1.1.

**Proof of Theorem 1.1** Lemma 3.1 gives the existence of a solution of (1) satisfying (7).

Let  $u_1$  and  $u_2$  be two solutions of (1), which verify (7), that is

$$\begin{cases} \partial_t u_1 + a u_1 \partial_x u_1 + \alpha \partial_x^3 u_1 = 0, & t > 0, x \in \mathbb{R}, \\ u_1(0, x) = u_{1,0}(x), & x \in \mathbb{R}, \end{cases}$$

$$\begin{cases} \partial_t u_2 + a u_2 \partial_x u_2 + \alpha \partial_x^3 u_2 = 0, & t > 0, x \in \mathbb{R}, \\ u_2(0, x) = u_{2,0}(x), & x \in \mathbb{R}. \end{cases}$$

Then, the function

$$\omega = u_1 - u_2 \tag{62}$$

is the solution of the following Cauchy problem:

$$\begin{cases} \partial_t \omega + a(u_1 \partial_x u_1 - u_2 \partial_x u_2) + \alpha \partial_x^3 \omega = 0, & t > 0, x \in \mathbb{R}, \\ \omega(0, x) = u_{1,0}(x) - u_{2,0}(x), & x \in \mathbb{R}. \end{cases} \tag{63}$$

Observe that, thanks to (62)

$$\begin{aligned} u_1 \partial_x u_1 - u_2 \partial_x u_2 &= u_1 \partial_x u_1 - u_1 \partial_x u_2 + u_1 \partial_x u_2 - u_2 \partial_x u_2 \\ &= u_1 \partial_x \omega + \partial_x u_2 \omega, \end{aligned} \tag{64}$$

$$\partial_t \omega = -a u_1 \partial_x \omega - a \partial_x u_2 \omega - \alpha \partial_x^3 \omega. \tag{65}$$

Multiplying (65) by  $2\omega$ , an integration on  $\mathbb{R}$  gives

$$\begin{aligned} \frac{d}{dt} \|\omega(t, \cdot)\|_{L^2(\mathbb{R})}^2 &= 2 \int_{\mathbb{R}} \omega \partial_t \omega dx \\ &= -2a \int_{\mathbb{R}} u_1 \omega \partial_x \omega dx - 2a \int_{\mathbb{R}} \partial_x u_2 \omega^2 dx - 2\alpha \int_{\mathbb{R}} \omega \partial_x^3 \omega dx \\ &= a \int_{\mathbb{R}} \partial_x u_1 \omega^2 dx - 2a \int_{\mathbb{R}} \partial_x u_2 \omega^2 dx + 2\alpha \int_{\mathbb{R}} \partial_x \omega \partial_x^2 \omega dx \\ &\leq a \|\partial_x u_1\|_{L^\infty((0,T) \times \mathbb{R})} \|\omega(t, \cdot)\|_{L^2(\mathbb{R})}^2 + 2|a| \|\partial_x u_2\|_{L^\infty((0,T) \times \mathbb{R})} \|\omega(t, \cdot)\|_{L^2(\mathbb{R})}^2. \end{aligned} \tag{66}$$

Fix  $T > 0$ . Observe that, since  $u_1, u_2 \in H^2(\mathbb{R})$ , for every  $0 \leq t \leq T$ , we have

$$\|\partial_x u_1\|_{L^\infty((0,T) \times \mathbb{R})}, \|\partial_x u_2\|_{L^\infty((0,T) \times \mathbb{R})} \leq C(T). \tag{67}$$

Therefore, by (66) and (67),

$$\frac{d}{dt} \|\omega(t, \cdot)\|_{L^2(\mathbb{R})}^2 \leq C(T) \|\omega(t, \cdot)\|_{L^2(\mathbb{R})}^2.$$

The Gronwall Lemma and (62) give (8).

## 4. Conclusion

By proving several a priori estimates and using the Aubin-Lions Lemma we show the convergence of the solution of (2) converges to the unique one of (1) as the coefficient  $\beta$  goes to 0.

## References

---

- [1] A. Biswas, H. Triki, Manel Labidi. Bright and dark solitons of the Rosenau-Kawahara equation with power law nonlinearity. *Physics of Wave Phenomena*. 2011; 19(1): 24-29.
- [2] Qodrat Ebadi, Aida Mojaver, Houria Triki, Ahmet Yildirim. Topological solitons and other solutions of the Rosenau-KdV equation with power law nonlinearity. *Romanian Journal of Physics*. 2013; 58(1-2): 3-14.
- [3] Jinsong Hu, Youcai Xu, Bing Hu. Conservative linear difference scheme for Rosenau-Kdv equation. *Adv. Math. Phys.* 2013.
- [4] Takuji Kawahara. Oscillatory solitary waves in dispersive media. *Journal of the Physical Society of Japan*. 1972; 33: 260-264.
- [5] M. Labidi, A. Biswas. Application of He's principles to Rosenau-Kawahara equation. *Math. Eng. Sci. Aerospace*. 2011; 2(2): 183-187.
- [6] P. Razborova, B. Ahmed, A. Biswas. Solitons, shock waves and conservation laws of Rosenau-KdV-RLW equation with power law nonlinearity. *Appl. Math. Inf. Sci.* 2014; 8(2): 485-491.
- [7] D. J. Korteweg, G. de Vries. On the change of form of long waves advancing in a rectangular canal, and on a new type of long stationary waves. *Philos. Mag.* 2009; 39(240):422-443.
- [8] Yassine Benia, Andrea Scapellato. Existence of solution to Korteweg-de Vries equation in a non-parabolic domain. *Nonlinear Anal.* 2020; 195: 111758.
- [9] Giuseppe Maria Coclite, Lorenzo di Ruvo. On the solutions for an Ostrovsky type equation. *Nonlinear Anal. Real World Appl.* 2020; 55: 103141.
- [10] C. E. Kenig, G. Ponce, L. Vega. Well-posedness and scattering results for the generalized korteweg-de vries equation via the contraction principle. *Commun. Pure Appl. Math.* 1993; 46: 527-620.
- [11] A. J. Mendez. On the propagation of regularity for solutions of the fractional Korteweg-de Vries equation. *J. Differ. Equ.* 2020; 269(11): 9051-9089.
- [12] N. A. Kudryashov. On new travelling wave solutions of the KdV and the KdV-Burgers equations. *Commun. Nonlinear Sci. Numer. Simul.* 2009; 14: 1891-1900.
- [13] Giuseppe Maria Coclite, Lorenzo di Ruvo. A singular limit problem for conservation laws related to the Rosenau equation. *Juor. Abstr. differ. Equ. Appl.* 2017; 8(3): 24-47.
- [14] P. G. LeFloch, R. Natalini. Conservation laws with vanishing nonlinear diffusion and dispersion. *Nonlinear Anal. Ser. A: Theory Methods*. 1992; 36(2): 212-230.
- [15] M. E. Schonbek. Convergence of solutions to nonlinear dispersive equations. *Comm. Partial Differential Equations*. 1982; 7(8): 959-1000.
- [16] Amin Esfahani. Solitary wave solutions for generalized Rosenau-KdV equation. *Communications in Theoretical Physics*. 2011; 55(3): 396-398.
- [17] P. Rosenau. A quasi-continuous description of a nonlinear transmission line. *Phys. Scr.* 1986; 34: 827-829.
- [18] P. Rosenau. Dynamics of dense discrete systems. *Prog. Theor. Phys.* 1988; 79: 1028-1042.
- [19] Giuseppe Maria Coclite, Lorenzo di Ruvo. Well-posedness of the classical solutions for a Kawahara-Korteweg-de Vries-type equation. *Journal of Evolution Equations*. 2020. Available from: <https://doi.org/10.1007/s00028-020-00594-x>.
- [20] M. A. Park. On the Rosenau equation. *Mat. Appl. Comput.* 1990; 9: 145-152.
- [21] J. M. Zuo. Solitons and periodic solutions for the Rosenau-KdV and Rosenau-Kawahara equations. *Appl. Math. Comput.* 2009; 215(2): 835-840.
- [22] P. Razborova, H. Triki, A. Biswas. Perturbation of dispersive shallow water waves. *Ocean Eng.* 2013; 63: 1-7.
- [23] Maobo Zheng, Jun Zhou. An average linear difference scheme for the generalized Rosenau-KdV equation. *J. Appl. Math.* 2014. Available from: <http://dx.doi.org/10.1155/2014/202793>.
- [24] Giuseppe Maria Coclite, Lorenzo di Ruvo. A singular limit problem for conservation laws related to the Rosenau-Korteweg-de Vries equation. *Journal de Mathématiques Pures et Appliquées*. 2017; 107(9): 315-335.
- [25] J. Janwised, B. Wongsaijai, T. Mouktonglang, K. Poochinapan. A Modified three-Level average linear-implicit finite difference method for the Rosenau-Burgers equation. *Advances in Mathematical Physics*. Hindawi Publishing Corporation; 2014.
- [26] L. Liu, M. Mei, Y. S. Wong. Asymptotic behavior of solutions to the Rosenau-Burgers equation with a periodic initial

- boundary. *Nonlinear Analysis*. 2007; 67: 2527-2539.
- [27] Giuseppe Maria Coclite, Lorenzo di Ruvo. Well-posedness results for the continuum spectrum pulse equation. *Mathematics*. 2019; 7: 1006.
- [28] G. M. Coclite, M. Garavello. A Time Dependent Optimal Harvesting Problem with Measure Valued Solutions. *SIAM Journal on Control and Optimization*. 2017; 55(2): 913-935.
- [29] Giuseppe Maria Coclite, Mauro Garavello, Laura V. Spinolo. Optimal strategies for a time-dependent harvesting problem. *Discrete & Continuous Dynamical Systems-S*. 2016; 11(5): 865-900.
- [30] J. Simon. Compact sets in the space  $L^p(O, T; B)$ . *Annali di Matematica Pura ed Applicata*. 1986; 146: 65-94.
- [31] Giuseppe Maria Coclite, Lorenzo di Ruvo. Convergence of the ostrovsky equation to the ostrovsky-hunter one. *J. Differential Equations*. 2014; 256: 3245-3277.
- [32] Giuseppe Maria Coclite, Lorenzo di Ruvo. Wellposedness of the Ostrovsky-Hunter Equation under the combined effects of dissipation and short wave dispersion. *J. Evol. Equ.* 2016; 16: 365-389.