



Politecnico di Bari

Repository Istituzionale dei Prodotti della Ricerca del Politecnico di Bari

Assessment and source identification of pollution risk for touristic ports: Heavy metals and polycyclic aromatic hydrocarbons in sediments of 4 marinas of the Apulia region (Italy)

This is a pre-print of the following article

Original Citation:

Assessment and source identification of pollution risk for touristic ports: Heavy metals and polycyclic aromatic hydrocarbons in sediments of 4 marinas of the Apulia region (Italy) / Mali, Matilda; Dell'Anna, Maria Michela; Mastroilli, Pietro; Damiani, Leonardo; Piccinni, Alberto Ferruccio. - In: MARINE POLLUTION BULLETIN. - ISSN 0025-326X. - 114:2(2017), pp. 768-777. [10.1016/j.marpolbul.2016.10.063]

Availability:

This version is available at <http://hdl.handle.net/11589/99371> since: 2021-03-08

Published version

DOI:10.1016/j.marpolbul.2016.10.063

Terms of use:

(Article begins on next page)

Highlights

- Touristic ports are driving force for the local development of coastal regions
- Touristic port activity can affect the quality of port area and the nearby coast
- Cumulative indexes and PAH content ratio were used to assess port pollution sources
- The inner basin of the ports resulted unusually more affected than the outer one;
- Hydrodynamic or other factors need to be studied for a spot-on risk assessment

1
2
3 **Assessment and source identification of pollution risk for touristic ports:**

4 **Heavy metals and polycyclic aromatic hydrocarbons in sediments of 4**

5 **marinas of the Apulia region (Italy)**

6
7
8 *Matilda Mali^{a*}, Maria Michela Dell'Anna, Piero Mastroilli, Leonardo Damiani, Alberto*

9
10 *Ferruccio Piccinni*

11
12
13
14
15 ^aDICATECh, Politecnico di Bari, via Orabona, 4 I-70125 Bari, Italy

16
17
18 *Corresponding author e-mail address: matilda.mali@poliba.it (M. Mali); tel:+39 080 5963666;

19
20 fax: +39 080 5963414

21
22
23
24
25 **Abstract**

26
27 The Apulia region in Italy has the longest Adriatic coastline; thus, maritime tourism is the
28 driving force for its economic development. Pollution risk for four representative touristic ports
29 of the region was assessed by determining the concentrations of 10 metals, 16 polycyclic
30 aromatic hydrocarbons (PAHs) congeners, and the main nutrients. The cumulative mean Effects
31 Range–Median quotient (mERMq) was used to assess the hazard degree, while the distribution
32 patterns and content ratios of different PAH sediment concentrations were investigated to
33 identify the pollution sources. Principal component analyses indicated an anomalous pollution
34 trend for one of the small touristic ports assessed; this trend emerged from contamination by
35 heavy metals and PAHs to a larger extent than expected, considering the main activity in this
36 port, especially in its inner basin. The reason of this anomaly is thought to be the hydrodynamic
37 and/or other stress factors.
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56

57 **Keywords**

Hazard degree, port sediment, touristic port, heavy metals, PAHs, cumulative indices

Introduction

In a globalized economy, local development based on regional natural resources allows the regions to compete on a global scale. Maritime tourism is a driving force for the local development of regions with beaches, and touristic ports are important attraction points in such regions. Their economic benefits are based, on the one side, on environmental quality and scenic views of the nearby coastal areas and, on the other side, on the efficiency of the offered tourist services; they are therefore vulnerable to human activities that can adversely affect the ecosystem of the nearby areas. Thus, regions with beautiful sceneries, natural hydrologic structures, clean water, fresh air, and species diversity can suffer from pollution and lose their attractiveness (Mihalic̃, 2000; Ameer A. 2008; [Dias et al. 2014](#); [Anfuso et al. 2014](#); [Garcia et al. 2014](#); [Dalmora et al. 2016](#)).

The Apulia region is situated in southeast Italy and has the longest coastline. It has over 800 km of coastline, 6 certified blue-flag beaches, 30 important beach resorts, 3 regional areas of excellence, 3 marine reserves, and over 30 natural protected areas. Alongside the beach trail, many touristic ports play an important role for the local development of the region (MacCallum et al. 2011). In this context, the port expansion required to meet the increasing demands of residents and tourists would definitely facilitate touristic, commercial, and economic growth. Despite this benefit, the activities carried out in the touristic ports are also likely to cause deterioration of coastal landscape, thereby affecting the quality of marine water in the surrounding areas. (Petrosillo et al. 2010; Afuso et al. 2014; Pérez et al., 2014; Sanchís et al., 2015).

Trace metals and polycyclic aromatic hydrocarbons (PAHs) are the two major contaminant categories investigated in the marine environment, because of (1) their high toxicity and

1 environmental persistence and (2) their consequent inclusion as priority substances in the Water
2 Framework Directive 2000/60/EC (EC 2000). These contaminants can be introduced into the
3
4 port basins through various ways, for example, ship-borne oil discharges, dry-docking
5
6 operations, discharging of bilge oil, urban runoff, domestic wastewater discharges, and
7
8 atmospheric deposition across the air–sea interface. After being transported into the aquatic
9
10 ecosystems, these contaminants are absorbed by suspended solids and accumulate in port
11
12 sediments where they become even more resistant to degradation (Zakaria et al. 2002; De Luca
13
14 et al. 2005). When the concentration of these contaminants trapped in port sediments increases
15
16 significantly, they are leached to the surrounding water column, thereby leading to degradation
17
18 of the quality of the aquatic ecosystems (Lee et al. 2001; Tsapakis et al. 2006; Abdulla & Linden
19
20 2008; [Olieviro et al. 2012](#); [Ribeiro et al. 2013](#); [Civeira et al. 2016](#); [Mali et al. 2015](#); [Mali et al.](#)
21
22 2016).

23
24
25
26
27
28 To assess the effect of human activities on the environmental quality of marine areas in the
29
30 Apulia region, four ports were selected as they represent the numerous touristic ports operating
31
32 in the Apulian Adriatic coast (Figure 1a). The four selected ports – Torre a Mare (TM), Palese
33
34 (PL), San Giorgio (SG), and Monopoli (MP) – are characterized by different human activities
35
36 and different hydrodynamic morphologies. TM is a touristic port dealing with yachting and
37
38 landing of small-sized ships. It has a particular hydrodynamic structure that facilitates an
39
40 irreversible silting process of the port promoted by the materials that are periodically brought by
41
42 Levante wind toward the access channel and then transported into the inner basin (Figure S1).
43
44 MP is a commercial port dealing with yachting, fishery, and very important touristic/commercial
45
46 activities. The latter ones are concentrated mostly in the oldest piers. PL, similar to TM, is a
47
48 small-sized touristic port, currently designated for landing of small ships and yachts. SG port is a
49
50 natural bay and is not equipped with yachting facilities. Nevertheless, it is used for local fishery
51
52 activity or touristic needs. The distribution of 10 metals and metalloids (Al, As, Cd, Cr, Cu, Hg,
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Ni, Pb, Zn, and V) and 16 priority PAHs in sediments from all the four ports *was analyzed*. The
2 degree of potential health and environmental risk was assessed by calculating the cumulative
3
4 mean Effects Range–Median quotient (mERMq) value, according to Long et al. (2006); this
5
6 worldwide accepted method is an efficient tool that considers multiple factors affecting the
7
8 hazard of sediments. In the case of the PAH contamination degree assessment, several pollution
9
10 indices were used and the obtained results were compared to each other.
11
12

13 Finally, multivariate statistical analyses were carried out to identify specific port areas that are
14
15 strongly affected by heavy metal and PAH contamination and to establish their correlation with
16
17 pollution sources loading on port basins.
18
19
20
21
22
23

24 **1. Materials and methods**

25 *2.1 Geographical, morphological, and environmental settings*

26
27 The study area is a part of the Apulia region, a WNW–ESE elongated Italian region, flanked by
28
29 the Adriatic Sea, and characterized by three large Meso-Cenozoic limestone landmasses from the
30
31 north to south: the Gargano Promontory at the north, the Murge Plateau in the middle, and the
32
33 Salento Peninsula at the south. Several minor touristic ports punctuate the entire coastline, and
34
35 the four selected ports (MP, TM, SG, and PL) are located in the Murge Plateau landmass. With
36
37 regard to the geomorphological features, the coastal ecosystem is constituted by micritic and
38
39 calcarenitic limestones and sands, similar to most of the Adriatic coasts. Siliciclastic sediments
40
41 input from the exposed region to the shelf is low since few rivers flow in Apulia, thus supplying
42
43 small amounts of terrigenous material to their mouths. The only major watercourse of the area is
44
45 the Ofanto river, which is 170 km long and is the longest river flowing into the South Adriatic
46
47 Sea. Numerous ephemeral watercourses rhythmically affect the study area, but their contribution
48
49 to the coastal sediments is scarce because of their ephemeral and torrential regime. The land use
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 around the considered ports consists of residential neighborhoods. Numerous highly populated
2 cities (Barletta, Trani, Bisceglie, Bari, Polignano, and Monopoli) are located alongside the coast.
3
4 A prime cause of pollution is the lone incident of bombing during World War II that affected
5 only one of the selected ports (MP port). Moreover, there are currently no heavy industrial
6 activities in the area. The maritime transportation, fishery industry, and bathing and yachting
7 activities are the principal environmental stressors for the quality of the harbor basins. The effect
8 of such anthropogenic pressure can be observed often in the summer season with the increasing
9 fishery and yachting activities. The domestic sewage water from the tourism firms operating
10 alongside the coast is thought to affect the environment of the coastal area and related port
11 basins.
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26

27 **Figure 1 near here**

28 **Caption to Figure 1 near here**

29 *2.2 Sampling*

30
31
32
33
34
35
36 The sediment samples used in the present study were collected from May 2010 to September
37 2011. Sediments from the bottom of sea were sampled using a vibro-corer PF1, equipped with a
38 liner and a support vessel and provided with the differential GPS system for positioning of
39 sampling cores (Table S1). The sample cores were up to 1.50–2.00 m deep and could reach the
40 rocky hard bottom of the harbor sediment. Each core was divided into several 50-cm length
41 subcores. Aliquots of wet sediment samples were transferred from the liner into cleaned plastic
42 bags and stored at 4°C during transportation to the laboratory.
43
44
45
46
47
48
49
50
51
52
53
54
55

56 *2.3 Analytical methods*

1 Sediments stored at -20°C after sampling were dried at 40°C for 48 h. Each sediment was
2 classified according to Shepard (1954) into four classes: gravel: $> 2\text{ mm}$; sand: $2-0.063\text{ mm}$;
3 $0.063-0.002\text{ mm}$; and clay $< 0.002\text{ mm}$. A set of ASTM sieves was used for the granulometric
4 separation (Romano et al., 2001).
5
6

7
8
9 The trace metal concentrations were measured by inductively coupled plasma mass spectrometry
10 (ICP/MS X Series, Thermo Fisher Scientific) after sample mineralization by total acid digestion
11 (HCl, HNO_3 , and HF) (Pellegrini et al., 2001). The $<63\text{-}\mu\text{m}$ fraction dried at 105°C was used for
12 the determination of the metal and metalloids. The results are expressed on the dry weight basis.
13
14 The detection limits (LODs) were calculated from 3 replicates of procedural blanks. The LODs
15 estimated were expressed in mg/kg and were equal to 1 ppb for all metals and metalloids
16 analyzed. Marine Sediment Reference Material NIST 2702 (Inorganics in Marine Sediment) was
17 used to control the analysis quality: the agreement between the analytical results for certified and
18 measured values was satisfactory with recoveries between 80% and 100% for all metals and
19 metalloids analyzed.
20
21

22
23
24 The PAHs considered for this study were as follows: naphthalene (NA), acenaphthene (Ap),
25 acenaphthylene (Apl), phenanthrene (Phe), anthracene (An), fluoranthene (Flt), benz[a]
26 anthracene (BaA), chrysene (Chry), benz[b]fluoranthene (BbF), benz[k]fluoranthene (BkF),
27 benzo[a]pyrene (BaPy), benzo[e]pyrene (BePy), dibenz[a,h]anthracene (DBA), fluorene (fl),
28 indeno[1,2,3-cd] (Ind), and pyrene (PY). These PAHs were first extracted with
29 cyclohexane/methanol mixture and then determined by high-pressure liquid chromatography,
30 according to Ausili's (2001) method.
31
32

33
34
35 Total nitrogen (N_{tot}) was determined by the Kjeldahl method (Kjeldahl, J. 1983), while total
36 phosphorous (P_{tot}) by the elemental analysis procedure with a Perkin-Elmer 240B CHN
37 Elemental Analyzer (Aspilla et al., 1976).
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1 Total organic carbon (TOC) was determined by the elemental analysis procedure with a Perkin–
2 Elmer 240B CHN Elemental Analyzer, after removing carbonates by reaction with hydrochloric
3 acid (Nieuwenhuize et al., 1994).
4
5

6
7 All analyses were conducted in the ARPA (Apulian Regional Agency for Environmental
8 Protection) laboratories according to standardized protocols.
9
10

11 12 13 14 15 16 17 *2.4 Statistical analyses*

18
19 Unscrambler Program (v. 9.2 Camo, Oslo, Norway) and Statistica7Software were used to
20 perform statistical analyses. The PCA was carried out to assess geochemical associations and to
21 ascertain the relationship among contaminants in the port sediment and their possible common
22 sources (Lu et al., 2010; Varol et al. 2011; [Silva et al. 2009; 2012; Rodriguez-Iruretagiiena et al.,](#)
23 [2015](#)).
24
25
26
27
28
29
30
31
32
33

34 *2.5 Pollution risk degree for heavy metals*

35
36 The pollution risk associated with heavy metal contamination was determined by calculating
37 the mean ERM quotient (mERMq) (Long et al. 2006) on the basis of the relevant ERM
38 values (Long et al., 1995), according to the following equations:
39
40
41
42

$$43 \quad mERMq = \sum \left(\frac{ERMq}{n} \right) \quad \text{Eq (1a)}$$

$$44 \quad ERMq = \frac{Ci}{ERMi} \quad \text{Eq (1b)}$$

45
46
47
48
49
50
51
52 where Ci = the total content of the selected contaminant, $ERMi$ = the ERM value of the
53 selected contaminant, and n = the number of variables.
54
55
56

57 *2.6 Ecological risk degree for PAHs*

1 The diagnostic content ratio procedure (Soclo et al. 2000; Riccardi et al. 2013; Garcia et al.
2 2014) was used to identify the PAH origin. The diagnostic content ratios (ppb/ppb) used were
3
4 Phe/An, Flt/Py, BaA/(BaA+Chry), and low-molecular-weight PAHs/high-molecular-weight
5
6 PAHs (LMWPs/HMWPs).
7

8
9 The ecological risk associated with PAHs was calculated by the same cumulative index mERMq
10
11 method (Long et al., 2006) used for heavy metals and metalloids to perform a correct
12
13 comparative analysis.
14
15
16
17

18 **2. Results and Discussion**

19 *3.1 Sediment characteristics*

20
21
22 The sediments of the four considered ports were composed of a sandy fraction in the range of
23
24 21–97%_w, a coarse fraction in the range of 3–44%_w, and a fine grain-sized fraction (silt and
25
26 clay) in the range of 2–78%_w (Table S2). The sandy fraction represents the main component for
27
28 PL and SG ports, characterized by a scarce amount of fine grain-sized fraction (less than 4%_w).
29
30 On the contrary, the fine grain-sized fraction is present in much higher amount in TM and MP
31
32 samples, where differences were invariably observed between samples coming from the inner
33
34 and the outer (at the entrance of the port) zones, respectively, with the amount of fine fraction
35
36 higher (ranging from 17 to 78%_w) in the inner sites than in sediments collected at the entrance of
37
38 the port (fine fraction in the range of 4–10%_w).
39
40
41
42
43
44

45
46 Several factors influence the distribution of fine-grained sediments in the marine ecosystem, the
47
48 most important being sediment transportation and sedimentary processes (Tavakoly et al. 2011;
49
50 Tavakoly et al. 2014). SG is an almost natural bay; thus, it is expected to be sandy, because it is
51
52 well documented that natural bays have predominantly sandy sediments, with rare occurrences of
53
54 gravel (McManus et al. 1998).
55
56
57
58
59
60
61
62
63
64
65

PL is sandy similar to SG, although the former is equipped with artificial port facilities (piers, breakwaters, and jetties). It seems that these facilities have a negligible effect on the hydrodynamics of the port area and thus on the fine-sized distribution of the PL sediments. The TM sediments are mainly constituted by mud sediments, especially in the inner basins and the MP port. The accumulation of mud sediments in TM and MP ports is probably facilitated by the presence of artificial port facilities such as breakwaters and headlands, which commonly protect the port structure from waves and marine currents, thereby minimizing the hydrodynamic energy inside the port area (Hancock et al. 2001). Thus, it is conceivable that the particular hydrodynamic morphology of the port affects the sediment composition.

3.2 Contaminant patterns

3.2.1 HMe distribution

Table 1 presents the statistical concentration data (minimum and maximum concentrations and standard deviations) for metals in all the 42 samples from the four considered ports. The metal distribution pattern, considering the entire database of port samples, was in the following order of concentrations: Zn>Pb>Cu>V>Cr>Ni>As>Hg>Cd, which differs substantially from both the upper crust concentrations (Wedepohl and Turekian, 1961) and the crustal average shale abundances (Wedepohl, 1995) (Table S5).

Metals & Metalloids*	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn	V
Wedepohl & Turekian (1961) (Upper Crust)	2.0	0.1	35.0	14.3	0.06	18.6	17.0	53.0	52.0
Wedepohl (1995) (Average crust)	1.7	0.1	126.0	25.0	0.04	56.0	14.8	98.0	65.0

Min	3.8	0.05	0.1	2.7	0.0	3.9	5.0	11.0	10.6
Max	19.0	0.98	69.0	248.0	1.4	28.0	256.0	456.0	71.0
SDev	3.1	0.21	17.7	54.0	0.3	7.5	64.8	98.6	17.3

Table 1. Upper crust concentrations (Wedepohl and Turekian, 1961), crustal average shale abundances (Wedepohl, 1995), and statistical concentration data (minimum and maximum concentrations and standard deviations) for metals and metalloids in all the 42 samples from the four considered ports. Concentrations are expressed in mg/kg.

Analysis of the concentration of heavy metals in each port revealed that sediments collected in TM and MP are very different from those coming from PL and SG. Figure 2 shows the mean and standard deviation values for HMe concentration calculated for each port. Higher concentrations of Pb, V, Cd, Hg, and Cu with respect to the crustal shale averages (Wedepohl, 1995) were observed in MP and TM ports, while lower HMe concentrations were determined in SG and PL samples; this finding follows the crustal average order (Wedepohl, 1995). The HMe contamination observed in MP could be explained by its commercial and touristic activities; therefore, different pollution sources could be identified (vessel batteries, discharge of wastewater, fuel and oil spills, bilge water, runoff water, etc.). On the contrary, the unusually high HMe concentrations detected in TM could be related to the particular hydrodynamic morphology of the port structure that facilitates the transport to the inner basins of fine sediments and related contaminants coming from the external coastal area.

Figure 2 near here

Caption to Figure 2 near here

In addition, differences were noted between the concentrations of heavy metals determined within the sampling sites located in the inner basin with respect to those located at the entrance of the port (Figure 3). An exception was noted for Hg, as the concentration of Hg determined within the inner sampling sites was similar to that determined at the entrance basin sampling

1 sites. It is well known that fine particle sediments tend to trap and concentrate contaminants
2 (Salomons and Fôrstner, 1984; Horowitz, 1991); therefore, the transport and fate of heavy metals
3 are always associated with fine sediment dynamics. As discussed above, the morphology of the
4 port structure facilitates the accumulation of silty sediments in the inner basin of the commercial
5 ports, and consequently, a high level of heavy metal concentration was registered at the ports
6 (Palanques et al. 1995; Villaescusa-Celaya et al. 2000; Fdez-Ortiz de Vallejuelo S et al. 2010;
7 Oliveira et al. 2013).

8
9
10
11
12
13
14
15
16
17 **Figure 3 near here**

18
19 **Caption to Figure 3 near here**

20 21 22 23 24 *3.2.2 PAH distribution*

25
26 The total PAH concentration in the sediments from the four studied ports varied from 100 to
27 21500 ppb of dry matrix, ranging from 50 to 800 ppb for PL and SG samples and from 90 to
28 21500 ppb for TM and MP cores (Table S3). The lowest concentrations were observed in the
29 sediments collected from PL and SG because they are almost uncontaminated natural bays. TM
30 samples were moderately polluted by PAHs (90–2020 ppb) because of shipping activities, while
31 the highest concentrations of total PAHs (up to 21460 ppb) were found in MP samples from the
32 old basin, located near densely populated areas (Figure 4). In addition, the old basin morphology
33 of MP produces a hydrodynamic stagnation zone in the inner harbor parts (sites MP25-
34 MP26_MP27-MP28, Table S1), which causes the accumulation of contaminants (HMes and
35 PAHs) in the stagnation area due to local hydraulic head gradients.

36
37 The most abundant PAHs in the sediments were those formed by 4–5 and 6 aromatic rings
38 (mainly fluoranthrene (Fln), pyrene (Py), benzo [a] anthracene (BaA), benzo[b] pyrene (BbPy),
39 and indene [1,2,] pyrene). The presence of HMWPs is due to their high sorption capacity and
40 resistance to degradation (Wang et al., 2009) and has been frequently observed in marine
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

sediments (Readman et al. 1987; Riccardi et al. 2013; Garcia et al. 2014), while the predominance of phenanthrene, fluoranthene and pyrene is symptomatic of the diesel fuel spill caused by shipping activities.

Figure 4 near here

Caption to Figure 4 near here

3.3 Assessment of ecological risk degree

3.3.1. Risk degree for HMeS

The pollution risk associated with heavy metal contamination was determined by calculating the mean ERM quotient (mERMq) (Long et al. 2006) on the basis of the relevant ERM values (Long et al., 1995). The quotient provides a more realistic measurement of risk toxicity than the simple concentration data, as it weighs the magnitude by which individual metals exceed the respective ERM values, thus ranking the risk categories of the sites with multiple contaminants. The obtained results, reported in Figure 5, are presented according to the toxicity categories established by Long et al., 2006 (Table S4).

Figure 5 near here

Caption to Figure 5 near here

As expected, the calculated mERMq values showed a range from severe to moderate toxicity for the samples of MP and TM with the exception of two sampling points located at the entrance of TM port (TM_E and TM_F), while a slight toxicity was observed for the sediments from SG and PL. No sampling point with severe toxicity (mERMq value >1.5) was detected in any of the four ports.

3.2.2 Risk degree according to PAHs

PAHs can be divided into three main classes according to their features. The first one is composed of PAHs originating from incomplete combustion at high temperature of recent and fossil organic material (pyrolytic origin); the second one includes PAHs released by the slow maturation of organic matter under the geochemical gradient conditions (petrogenic origin); and the last class contains PAHs derived from short-term diagenetic degradation of biogenic precursors (diagenetic origin). However, anthropogenic activity is generally considered to be the major source of PAHs in the world (Zakaria et al. 2002). Each reported source generates characteristic PAH patterns, and it is therefore possible to identify the process that generated PAHs at a certain site.

In the present study, the diagnostic isomeric content ratios of specific PAHs were used to distinguish between the various origins and to determine pollution sources and estimate the environmental risks of the port areas. The simultaneous evaluation of four isomeric content ratios was used to distinguish pyrolytic from petrogenic processes.

Phenanthrene is the most thermodynamically stable isomer with three condensed aromatic rings, and the ratio between phenanthrene and anthracene amounts (Ph/An) is commonly used to distinguish petrogenic source from pyrolytic input (Budzinski et al. 1997). Because this ratio is temperature-dependent (Alberty and Reif, 1988), high-temperature processes such as combustion of organic matter and fossil fuels (coal and petroleum) generate PAHs characterized by a low Ph/An ratio (<10), whereas the slow maturation of organic matter during catagenesis leads to higher Ph/An ratio values (Baumard et al. 1998a; Baumard et al. 1998b; Stogiannidis and Lame, 2015). Petrogenic PAHs are introduced in the port area through accidental oil spills, discharge from routine tanker operations, municipal and urban runoff, and other sources. Thus, two different classes of sediments can be defined: the first class with a Ph/An content ratio > 10

1 indicating petrogenic inputs and the second one characterized by a Ph/An content ratio < 10
2 indicating the dominance of pyrolytic sources. Since the bordering values are not very precise
3 and the limits between the two processes are not clear-cut (Budzinski et al., 1997), a
4 simultaneous comparison with other isomeric content ratios is recommended to provide a good
5 estimation of the different PAH sources (Raoux, 1991). The isomeric content ratio between
6 fluoranthene and pyrene concentrations (Flt/Py) is also used for PAH source identification.
7 Values greater than 1 are characteristic of the pyrolytic origin, whereas values lower than 1 are
8 related to petroleum hydrocarbons (Perra et al. 2009) By plotting the two ratios (Ph/An against
9 Flt/Py), two zones with two different pollution sources, pyrolytic and petrogenic, were defined.
10 (Figure 6). Our results show that the samples of MP and TM sediments have PAHs derived from
11 mixed pyrolytic and petrogenic sources, while the samples collected in the two natural bays PL
12 and SG are more of a petrogenic origin (Ph/An > 10 and/or Flt/Py > 1).
13
14 To corroborate our source identification, cross-plot for other two content ratios was carried out
15 (Figure 7): that between chrysene and benzo[a]anthracene (Chry/BaA) and that of the
16 LMWPs/HMWPs (Yunker et al., 2002). Chrysene and benzo(a)anthracene derived from the
17 processes of organic matter combustion at high temperatures have a Chry/BaA content ratio of
18 less than 1.0 (Parlanti, 1990), which is characteristic of a pyrolytic origin. On the contrary, the
19 low maturation of organic matter during burial in the sedimentary matrix results in
20 Chry/BaA > 1.0. An LMWPs/HMWPs content ratio of less than 1.0 is characteristic of a pyrolytic
21 source (Soclo et al., 2000), while spillage of fuel oil or light-refined petroleum products yields
22 values of LMWPs/HMWPs > 1.0 (Liu et al. 2008).
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Figure 6 near here

Caption to Figure 6 near here

Figure 7 near here

Caption to Figure 7 near here

The ecological risk associated with PAHs (organic pollutants) was calculated by the mERMq method (Long et al., 2006). The obtained results are comparable with the mERMq values calculated for metals and metalloids. The mERMq values for organic pollutants fitted well with those determined for metals and metalloids, and the strong relationship ($R^2=0.89$) between them suggests a common contamination source for both these classes of contaminants (Figure 9). Nevertheless, the high PAH mERMq values found in MP and TM harbors indicate a severe PAH pollution risk for the two commercial ports.

Figure 8 near here

Caption to Figure 8 near here

Figure 9 near here

Caption to Figure 9 near here

3.3.3 Total organic carbon and total nitrogen.

The C/N ratio of organic matter in surficial marine sediments has been frequently used as an indicator of terrigenous addition to marine sediments (Trask, 1955; Pocklington, Leonard, 1979; Faganelli et al. 1988; Mayers, 1994; [Sánchez et al. 2013](#)). The C/N content ratio for each sample site was calculated as an organic carbon source indicator, and the obtained results are

1 plotted in Figure 10. The C/N ratios ranged from 14 to 49 and increased with the increasing fine
2 fraction within the sediment samples. These C/N ratios are much higher than the typical C/N
3 values reported for most coastal sediments (C/N=6–10) (Faganelli et al. 1988; Oliveira et al.
4 2011; Oliveira et al. 2012) indicating that the organic matter inputs to harbor sediments are
5 highly affected by terrestrial and anthropogenic sources, especially in the case of TM and MP.
6
7
8
9
10
11
12
13
14
15
16

17 **Figure 10 near here**

18 **Caption to Figure 10 near here**
19
20
21
22
23
24
25

26 *3.4 Principal Component Analyses*

27
28
29
30
31 To identify the common pollution sources and to relate them with the determined contamination
32 degree, a PCA was performed on the entire dataset, as a non-supervised technique for comparing
33 the compositional patterns between sediment samples and for identifying factors influencing
34 them. The analysis was carried out for each contaminant class. For this purpose, two datasets
35 were considered: the first one was constituted by a matrix with 10 variables in columns (Zn, Pb,
36 Cu, V, Cr, Ni, As, Hg, Cd, and Al) and the sediment samples in rows (42), and the second one
37 was a matrix with 16 PAHs in columns and the 42 samples in rows.
38
39
40
41
42
43
44
45
46
47

48 Before the analysis, the data were pretreated by scaling and mean-centering.

49
50 For metals and metalloids, a model with two PCs (76% and 9% of explained variance,
51 respectively) was selected, explaining 85% of the total variance, since the results did not
52 significantly change when the third principal component (PC3 = 5% of explained variance) was
53 considered using the Full Cross Validation method (Figure 11).
54
55
56
57
58
59
60
61
62
63
64
65

Figure 11 near here

Caption to Figure 11 near here

The score plot in the space formed by the two PCs (Figure 11(a)) reveals the existence of two different groups of sediments with two different pollution profiles. The first group was formed by sediment samples from MP and TM ports. The loading plots show the influence of almost all heavy metal concentrations on this group. The second group includes all sediments collected from the natural bays (SG and PL) and the sampling sites located at the entrance of the two ports of TM and MP (TM_E/TM_F and MP_23) indicated by the black arrows in Figure 11(a). This group was not affected by any metal concentrations. Two subclusters are noted in the loading plots: the samples of the inner basin of MP and TM ports are affected by As and Hg amounts [distinguished by strong positive loadings on PC2 (Hg: 0.46; As: 0.453)]; both of them are considered as metals with a high level of toxicity in environmental studies. The second subcluster in the loading plots is influenced by Cd, Cu, Pb, V, and Zn concentrations, characterized by lower positive loadings on PC2 and higher positive loadings on PC1. The PC space configuration suggests that these metals may have a common source, probably fossil fuel combustion, losses of fuel during maneuvering of vessel within the port (Cd and Pb), and use of metal antifungal products (Cu, V, and Zn).

A PCA of PAH congeners was also carried out to analyze their sources, and the obtained results confirmed the HMe trend. Further, in this case, a model with two PCs was considered since they explain 95% of the total variance (93% and 3% of the explained variance by each respective PC) (Figure 12 (a) and (b)). The score and loading space configurations of PC1 and PC2 show that the natural bays (PL and SG) results are not affected by any PAH congener, while the two ports (TM and MP) are influenced by the PAH content. In particular, 2- to 3-ring PAHs with low

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

molecular weight affected TM samples, while the MP port was influenced predominantly by 4- to 6-ring PAHs with high molecular weight.

Figure 12 near here

Caption to Figure 12 near here

3.5 Pollution source identification

By performing the ratio analyses, cumulative indices, and multivariate statistical analyses used to identify possible sources of heavy metals and PAHs, the following considerations can be proposed:

i) The strong pollution determined for Pb, Cd, Cu, Zn, and Hg within the ports (MP and TM) suggests that fuel additives and combustion and corrosion of alloys used in vessel components are the main pollution source for these metals. Cu and Cd can also be related to the use of antifungal organometallic compound and to the maritime batteries, often spilled on the bottom sea area.

ii) The C/N ratio was high for MP and TM and was moderately high for SG, with only PL showing a normal C/N ratio with respect to the unpolluted marine ecosystem (i.e., C/N in the range from 6 to 10). The high C/N ratio could be due to domestic and municipal wastewater discharge in the coastal area, especially for TM and MP ports.

iii) On the basis of PAH ratio analyses, the main sources of PAHs in the sediments of MP and TM ports are anthropogenic with a dominant effect of pyrolytic genesis. The PAH contamination trend is similar to that found for metal pollution. In addition, only for the TM port, besides fossil fuels burned by yachts and vessels, PAHs may have other petrogenic sources such as petrol, coal, grass, and wood.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

iv) The higher mERMq values for PAHs with respect to those determined for metals and metalloids suggest that the major risk for the four considered port basins is constituted by organic pollutants. Because of their high carcinogenic effect, the meticulous identification and prevention of fuel losses within the port area should be strictly and continuously monitored.

4. Conclusion

The different distribution patterns determined within the four studied tourist ports showed that multiple sources contributed to the increase in the content of pollutants (metals and PAHs), and human activity is one of the main sources of contamination of the coastal area. Generally, low hazard degrees were determined for the natural bay (SG) and one of the touristic ports (PL) that are not subjected to heavy anthropogenic activities. With regard to TM and MP, heavy metal pollution was moderate, but large risk indices were estimated for PAHs, which should be considered of high concern. In fact, the mERMq values calculated for PAHs suggest a relatively elevated level of toxicity of TM and MP sediments. A special consideration should be made for the TM port. In fact, although TM is a small touristic port, its hazard degree resembles that of MP, which, on the contrary, is a bigger port characterized by a higher extent of commercial and touristic activities. The serious threats determined for TM sediments are probably related to the hydrodynamic port structure that facilitates the entrance of contaminated fine sediments from the outer area and hampers their exit from the inner basin. This anomalous behavior observed for TM suggests that other factors such as hydrodynamic parameters may play a role in determining pollution risk. Further investigations are warranted to verify this hypothesis.

Acknowledgments.

The authors thank ARPA Puglia for all the chemical analyses and the Municipality of Bari for encouraging this study.

1
2 **References**
3

4 Albery, R.A. & Reif, A.K., 1988. Standard chemical thermodynamic properties of polycyclic
5 aromatic hydrocarbons and their isomer groups. I. Benzene series. *J. Phys. Chem. Ref. Data* 17,
6
7 241-253.
8
9

10
11 Anfuso, G., Williams A.T., Cabrera Hernández J.A., Pranzini E., 2014. Coastal scenic
12 assessment and tourism management in western Cuba. *Tourism Management*; Vol. 42: 307–320;
13
14 <http://dx.doi.org/10.1016/j.tourman.2013.12.001>.
15
16

17
18 Ameer A., PhD, Olof Linden, PhD (editors). 2008. Maritime traffic effects on biodiversity in the
19 Mediterranean Sea: Review of impacts, priority areas and mitigation measures. Malaga, Spain:
20
21 IUCN Centre for Mediterranean Cooperation. 184.
22
23

24
25 Aspilla, K.I. Agemain, H and Chau, A. 1976. A semi-automated method for the determination of
26 inorganic, organic and total phosphorus in sediments. *Analyst*, 101 187-197.
27
28

29
30 Ausili, A., 2001. Analisi degli idrocarburi policiclici aromatici – Sedimenti. Scheda 9. In:
31
32 Cicero, A.M., Di Girolamo, I. (Eds.), *Metodologie Analitiche di Riferimento del Programma di*
33
34 *Monitoraggio per il controllo dell'ambiente marino-costiero (triennio 2001–2003)*. Ministero
35
36 dell'Ambiente e della Tutela del Territorio. ICRAM©, Roma.
37
38

39
40 Baumard P, Budzinski H, Garrigues P, Sorbe JC, Burgeot T, Bellocq J: Concentration of PAH in
41 various marine organisms in relation to those in sediments to throphic level. 1998a. *Mar Poll*
42
43 *Bull* 36:951–960.
44
45

46
47 Baumard P, Budzinski H, Michon Q, Garrigues P, Burgeot T, Bellocq J. 1998b. Origin and
48 bioavailability of PAHs in the Mediterranean Sea from mussel and sediment records. *Estuarine*
49
50 *Coastal Shelf Sci.* 47, 77–90
51
52
53
54
55
56
57
58
59

1 Budzinski H, Jones I, Bellocq J, Pie´rard C, Garrigues P. 1997. Evaluation of sediment
2 contamination by polycyclic aromatic hydrocarbons in the Gironde Estuary. *Mar. Chem.* 58:85–
3
4 97.
5

6
7 Civeira M.S., Ramos C.G., Oliveira, M. L.S. Kautzmann, R.M., Taffarel S.R., Teixeira E.C.,
8
9 Silva, Luis F.O., 2016. Nano-mineralogy of suspended sediment during the beginning of coal
10 rejects spill. *Chemosphere (Oxford)*, v. 145, p. 142-147.
11

12
13 Dalmora A.C., Ramos C.G., Querol X, Kautzmann R.M., Olivira M.L.S. Taffarel S.R., Moreno
14 T., Silva L.F.O. 2016. Nanoparticulate mineral matter from basalt dust wastes. *Chemosphere*
15
16
17
18
19
20
21 (Oxford), v. 144, p. 2013-2017, 2016.

22
23 Dias C.L., Olivira M.L.S. Hower J.C., Taffarel S.R., Kautzmann R.M., Silva L.F.O.
24 Nanominerals and ultrafine particles from coal fires from Santa Catarina, South Brazil.
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000

1 Garcia, K.O.; Teixeira, E.C.; Agudelo-Castaneda, D.M.; Braga, M.; Alabarse, P.G.; Wiegand, F.;

2 Kautzmann, R.M.; Silva, L.F.O. Assessment of nitro-polycyclic aromatic hydrocarbons in PM1

3

4 near an area of heavy-duty traffic. *Science of the Total Environment*, v. 479-480, p. 57-65, 2014.

5

6 Hancock, G.J., Olley, J.M. and Wallbrink, P.J. 2001. Sediment transport and accumulation in

7

8 Westernport, CSIRO Land and Water Technical Report, 47/01.

9

10

11 Horowitz, A.J., 1991. *A Primer on Sediments-trace Element Chemistry*, 2nd Edition. Lewis

12

13 Publishers, Chelsea, MI

14

15 Kjeldahl, J. (1883) "Neue Methode zur Bestimmung des Stickstoffs in organischen Körpern"

16

17 (New method for the determination of nitrogen in organic substances), *Zeitschrift für analytische*

18

19 *Chemie*, 22 (1): 366-383

20

21

22

23

24 Lee, K.T., Tanabe, S., Koh, C.H., 2001. Distribution of organochlorine pesticides in sediments

25

26 from Kyeonggi Bay and nearby areas, Korea. *Environ. Pollut.* 114. 207–213.

27

28 Liu Y., Chen L., Jianfu Z., Qinghui H., Zhiliang Z. Hongwen G. 2008. Distribution and sources

29

30 of polycyclic aromatic hydrocarbons in surface sediments of rivers and an estuary in Shanghai,

31

32 China. *Environmental Pollution*, 154, 2, 298–305.

33

34

35

36 Long, E.R., D.D. Macdonald, S.L. Smith and F.D Calder, 1995. Incidence of adverse biological

37

38 effects with ranges of chemical concentrations in marine and estuarine sediments. *Environmental*

39

40 *Management* 19: 81-97.

41

42

43

44 Long, E.R., Ingersoll, C.G. and MacDonald, D.D. 2006. Calculation and uses of mean sediment

45

46 quality guideline quotients: A critical review. *Environmental Science & Technology* 40, 1726-

47

48 1736.

49

50

51 Lu, X. Wang, L. Li, L.Y. Huang, L. Kang D. 2010. Multivariate statistical analysis of heavy

52

53 metals in street dusts of Baoji, NW China; *J. Hazard. Mater.*, 173, 744–749

54

55

56 MacCallum, N., Baert T., Rosato P., Barbieri S. 2011. *Sustainable Tourism and Local*

57

58 *Development in Apulia Region*". *OECD Local Economic and Employment Development*

59

60

61

62

63

64

65

(LEED) Working Papers, No. 2011/02, OECD Publishing, Paris. DOI:

<http://dx.doi.org/10.1787/5kghrjkgphd5-en>

Mali M., Dell'Anna MM., Mastrorilli P., Damiani L., Ungaro N., Gredilla A., Fdez-Ortiz de Vallejuelo S. 2016. Identification of hot spots within harbour sediments through a new cumulative hazard index. Case study: Port of Bari, Italy. *Ecol. Indicators* 60, 548–556.

Mali M., Dell'Anna MM, Mastrorilli P., Damiani L., Ungaro N., Belviso C., Fiore S., 2015. Are conventional statistical techniques exhaustive for defining metal background concentrations in harbour sediments? A case study: The Coastal Area of Bari (Southeast Italy). *Chemosphere* 138: 708–717.

McManus, J., Komar, D.P., Bostrom, G., Cobert, G. Marra, J.J. 1998. Sediment sources and the history of accumulation in Tillamook bay, Oregon. Final Report. Tillamook Bay National Estuary Project. Online access on July 5th, 2016.

https://nrimp.dfw.state.or.us/web%20stores/data%20libraries/files/Watershed%20Councils/Watershed%20Councils_469_DOC_TBaySedimentStudy.pdf

Mihalic, T., 2000. Environmental management of a tourist destination: a factor of tourism competitiveness. *Tourism Manage.* 21, 65–78.

Nieuwenhuize, J., Maas, Y.E.M., Middleburg, J.J., 1994. Rapid analysis of organic carbon and nitrogen in particulate materials. *Mar. Chem.* 45, 217–224

Oliveira CR, Dos Santos Madureira LA. 2011. Assessment and sources of nonaromatic hydrocarbons in surface sediments of three harbors in Santa Catarina State, Brazil. *Environ Monit Assess.* 173(1-4): 91-105. doi: 10.1007/s10661-010-1373-z.

Oliveira, M. L.S. ; Ward, C.R.; French, D.; Hower, J.C.; Querol, X.; Silva, L.F.O., 2012. Mineralogy and leaching characteristics of beneficiated coal products from Santa Catarina, Brazil. *International Journal of Coal Geology*, v. 94: 314-325;

1 Oliveira, M.L.S.; Ward, C.R.; Sampaio, C.H.; Querol, X.; Cutruneo, C.M.N.L.; Taffarel, Silvio
2 R.; Silva, L.F.O. 2013. Partitioning of Mineralogical and Inorganic Geochemical Components of
3
4 Coals from Santa Catarina, Brazil, by Industrial Beneficiation Processes. *International Journal of*
5
6 *Coal Geology*, v. 116: 75-92.

7
8
9 Palanques A., Diaz J.I. & Farran M., 1995. Contamination of heavy metals Heavy metals in the
10 suspended and surface sediment of the Gulf of Cadiz (Spain): the role of sources, currents,
11
12 pathways and sinks. *Oceanologica Acta*, 18(4), 469-477.

13
14
15 Parlanti, E., 1990. Utilisation des hydrocarbures comme traceurs d'origine de la matiere
16 organique sedimentaire en milieu marin. Etude du Golfe du Lyon et du Golfe de Gascogne
17
18 (Programme Ecomarge) PhD thesis Nr 495, University Bordeaux I, Bordeaux, France; 289 pp.

19
20
21 Perra, G., Renzi M., Guerranti C., Focardi S.E., 2009. Perra Polycyclic aromatic hydrocarbons
22 pollution in sediments: distribution and sources in a lagoon system (Orbetello, Central Italy);
23
24 TWB, *Transit. Waters Bull.* 3(1), 45-58.

25
26
27 Petrosillo I. Vassallo P., Valente D., Mensa J.A., Mauro F., Zurlini G., 2010. Mapping the
28 environmental risk of a tourist harbor in order to foster environmental security: Objective vs.
29
30 subjective assessments. *Mar. Poll. Bull.* 60: 1051–1058.

31
32
33 Raoux C. 1991. Modelisation du mecanisme de contamination par les hydrocarbures aromatiques
34 polycycliques (HAP) des sediments costiers de Mediterranee: Consequences sur la
35
36 biodisponibilite´ des HAP dans les milieu marins. PhD thesis. University Bordeaux I, Bordeaux,
37
38 France.

39
40
41 Readman, J.W., R.F.C. Mantoura and M.M. Rhead, 1987. A record of polycyclic aromatic
42
43 hydrocarbon PAH pollution obtained from accreting sediments of the Tamar Estuary, UK:
44
45 evidence for not equilibrium behaviour of PAH. *Sci. Total Environ*, 66: 73-94.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Ribeiro, J.; Taffarel, S. R.; Sampaio, C. H.; Flores, D.; Silva, L.F.O. Mineral speciation and fate of some hazardous contaminants in coal waste pile from anthracite mining in Portugal. *International Journal of Coal Geology*, v. 109-110, p. 15-23, 2013.

Riccardi C, Di Filippo P, Pomata D, Di Basilio M, Spicaglia S., 2013. Identification of hydrocarbon sources in contaminated soils of three industrial areas. *Sci. Total. Environ.* 450, 13–22.

Rodriguez-Iruretagiiena, A.; Fdez-Ortiz de Vallejuelo, S.; Gredilla, A.; Ramos, C.G.; Oliveira, M.L.S.; Arana, G.; De Diego, A.; Madariaga, J.M.; Silva, L.F.O. Fate of hazardous elements in agricultural soils surrounding a coal power plant complex from Santa Catarina (Brazil). *Science of the Total Environment*, v. 508, p. 374-382, 2015.

Romano, E., Gabellini, M., 2001. Analisi delle caratteristiche granulometriche – Sedimenti, scheda 3. In: Cicero, A.M., Di Girolamo, I. (Eds.), *Metodologie Analitiche di Riferimento del Programma di Monitoraggio per il controllo dell'ambiente marino costiero (triennio 2001–2003)*. Ministero dell'Ambiente e della Tutela del Territorio. ICRAM©, Rome.

Salomons, W., Forstner, U., 1984. *Metals in the Hydrocycle*. Springer-Verlag, Berlin, Heidelberg/New York/Tokyo, ISBN 3540127550.

Sánchez A, López-Ortiz B.E., Aguiñiga-García S. Balart E. 2013. Distribution and composition of organic matter in sediments of the oxygen minimum zone of the Northeastern Mexican Pacific: paleoceanographic implications. *Journal of Iberian Geology* 39 (1) 2013: 111-120: http://dx.doi.org/10.5209/rev_JIGE.2013.v39.n1.41753.

Shepard F.P., 1954. Nomenclature based on sand–silt–clay ratios. *J. Sed. Petr.*, 24, 151-158.

Silva, L.F.O. ; Oliveira M.L.S., da Boit, K.M.; Finkelman, R.B., 2009. Characterization of Santa Catarina (Brazil) coal with respect to human health and environmental concerns. *Environmental Geochemistry and Health*, v. 31, p. 475-485, 2009.

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Silva, L.F.O.; Sampaio, C.H.; Guedes, A.; Fdez-Ortiz de Vallejuelo, S.; Madariaga, J.M. Multianalytical approaches to the characterisation of minerals associated with coals and the diagnosis of their potential risk by using combined instrumental microspectroscopic techniques and thermodynamic speciation. *Fuel*, v. 94, p. 52-63, 2012.

Soclo, H.H. Garrigues, P. Ewald. M. 2000. Origin of polycyclic aromatic hydrocarbons (PAHs) in coastal marine sediments: case studies in Cotonou (Benin) and Aquitaine (France) areas. *Marine Pollution Bulletin*, 40, 387–396.

Tavakoly Sanny S.B., Sulaiman A.B., Monazami G., Salleh A., 2011. Assessment of Sediment Quality According To Heavy Metal Status in the West Port of Malaysia, *World Academy of Sciences, Engineering and Technology*, 2011, 633;

Tavakoly Sany SB, Hashim R, Rezayi M, Salleh A, Safari O., 2014. A review of strategies to monitor water and sediment quality for a sustainability assessment of marine environment *Environ Sci. Pollut. Res.* 21, 813–833.

Tsapakis, M., Apostolaki, M., Eisenreich, S., and Stephanou, E. G., 2006. Atmospheric deposition and marine sedimentation fluxes of polycyclic aromatic hydrocarbons in the Eastern Mediterranean Basin, *Environ. Sci. Technol.*, 40, 4922–4927

Turekian, K.K., Wedepohl, K.H., 1961. Distribution of the elements in some major units of the earth's crust. *Bull. Geol. Soc. Amer.* 72, 175–192.

Varol, M., 2011. Assessment of heavy metal contamination in sediments of the Tigris River (Turkey) using pollution indices and multivariate statistical techniques. *J. Hazard. Mater.* 195, 355– 364.

Villaescusa-Celaya, J.A., Gutiérrez-Galindo, E.A. Flores-Muñoz G. 2000. Heavy metals in the fine fraction of coastal sediments from Baja California (Mexico) and California (USA), *Environ. Poll.* 108 (3), 453–462

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65

Wang D, Tian F, Yang M, Liu C, Li Y. 2009. Application of positive matrix factorization to identify potential sources of PAHs in soil of Dalian, China. *Environ Pollut.*, 157, 1559–1564.

Wedehpol, K.H., 1995. The composition of the continental crust. *Geochim. Cosmochim. Ac* 59, 1217-1232.

Yunker MB, Macdonald RW, Vingarzan R, Mitchell RH, Goyette D., 2002. PAHs in the Fraser River basin: a critical appraisal of PAH ratios as indicators of PAH source and composition. *Organic Geochem* 33, 489–515.

Zakaria MP, Takada H, Tsutsumi S, Ohno K, Yamada J, Kound E, Kumata H., 2002. Distribution of polycyclic aromatic hydrocarbons (PAHs) in rivers and estuaries in Malaysia: a widespread input of petrogenic PAHs. *Environ. Sci. Technol.* 36:1907–1918.

Metals & Metalloids*	As	Cd	Cr	Cu	Hg	Ni	Pb	Zn	V
Wedepohl & Turekian (1961) (Upper Crust)	2.0	0.1	35.0	14.3	0.06	18.6	17.0	53.0	52.0
Wedepohl (1995) (Average crust)	1.7	0.1	126.0	25.0	0.04	56.0	14.8	98.0	65.0
Min	3.8	0.05	0.1	2.7	0.0	3.9	5.0	11.0	10.6
Max	19.0	0.98	69.0	248.0	1.4	28.0	256.0	456.0	71.0
SDev	3.1	0.21	17.7	54.0	0.3	7.5	64.8	98.6	17.3

Table 1. Upper crust concentrations (Wedepohl and Turekian, 1961), crustal average shale abundances (Wedepohl, 1995) and statistical concentration data (minimum and maximum concentrations and standard deviations) for metals and metalloids in all of the 42 samples of the four considered port. Concentrations are expressed in mg/kg.

Figure(s)




Figure 1




Figure 2




Figure 3




Figure 4




Figure 9




Figure 10




Figure 11




Figure 12

Figure's captions

Figure 1: Map of study area and sampling sites within port; a) Apulian Coast; b) sampling sites of PL port, c) sampling sites of SG bay, d) sampling sites of TM port, e) sampling sites of MP port.

Figure 2. Means and Standard Deviations of concentration of heavy metals calculated from data obtained in all sampling stations for each port ($\text{mg}\cdot\text{kg}^{-1}$). Color code: blue = MP; red = PL; green = SG; light blue = TM; brown = Wedepohl values).

Figure 3. Means and Standard Deviations (SDev) of concentration of heavy metals metals calculated from data obtained, the red ones from sampling sites located in the inner basins of the four ports, and the blue ones from sites located in the entrance areas (all the concentration are expressed in $\text{mg}\cdot\text{kg}^{-1}$).

Figure 4. Means and Standard Deviations (SDev) of concentration of PAHs calculated from data obtained in all sampling stations for each port ($\text{mg}\cdot\text{kg}^{-1}$). Color code: blue = MP; red = PL; green = SG; light blue = TM).

Figure 5. Distribution of mERMq for HMEs the four considered ports.

Figure 6. Cross plot of Ph/An content ratio values against Flt/Py content ratio values for sediments from different ports (MP, TM, SG and PL)

Figure 7. Cross plot of LMWPs/HMWPs content ratio against Chry/BaA content ratio or sediments from different ports (MP, TM, SG and PL).

Figure 8. The distribution of mERMq values obtained for PAHs within sediments samples

Figure 9. Cross plot of mERMq values for PAHs and Metals.

Figure 10. C/N content ratio values for each sampling site of the four ports.

Figure 11. Score (a and c) and Loading (b) plot of PC1/PC2 projection. Explained variance (d) of selected PCs. In figure (a) red, green, blue and light blue points indicate samples coming from PL, SG, MP and TM bays, respectively. In figure (c) points of TM and MP located at the port entrance and all sample points coming from SG and PL are represented in blue, while inner points of MP and TM are represented in red.

Figure 12. Score plot and Loading plot of the projection of PC1/PC2 for PAHs. Red and green points indicate samples coming from PL and SB bay respectively, while blue and brown indicates samples coming from MP and TM respectively. See section 2.3 for the meaning of the PAH abbreviations in the loadings plot. ^(c) (d)