



Politecnico di Bari

Repository Istituzionale dei Prodotti della Ricerca del Politecnico di Bari

Nonlinear Schrödinger equation in the Bopp-Podolsky electrodynamics: Solutions in the electrostatic case

This is a pre-print of the following article

Original Citation:

Nonlinear Schrödinger equation in the Bopp-Podolsky electrodynamics: Solutions in the electrostatic case / D'Avenia, Pietro; Siciliano, Gaetano. - In: JOURNAL OF DIFFERENTIAL EQUATIONS. - ISSN 0022-0396. - STAMPA. - 267:2(2019), pp. 1025-1065. [10.1016/j.jde.2019.02.001]

Availability:

This version is available at <http://hdl.handle.net/11589/163475> since: 2021-03-11

Published version

DOI:10.1016/j.jde.2019.02.001

Publisher:

Terms of use:

(Article begins on next page)

NONLINEAR SCHRÖDINGER EQUATION IN THE BOPP-PODOLSKY ELECTRODYNAMICS: SOLUTIONS IN THE ELECTROSTATIC CASE

PIETRO D'AVENIA AND GAETANO SICILIANO

ABSTRACT. We study the following nonlinear Schrödinger-Bopp-Podolsky system

$$\begin{cases} -\Delta u + \omega u + q^2 \phi u = |u|^{p-2} u & \text{in } \mathbb{R}^3 \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2 \end{cases}$$

with $a, \omega > 0$. We prove existence and nonexistence results depending on the parameters q, p . Moreover we also show that, in the radial case, the solutions we find tend to solutions of the classical Schrödinger-Poisson system as $a \rightarrow 0$.

CONTENTS

1. Introduction	2
Notations	4
2. Deduction of the Schrödinger-Bopp-Podolsky system	5
3. Preliminaries	6
3.1. The operator $-\Delta + a^2 \Delta^2$	7
3.2. The functional setting	11
3.3. The Mountain Pass Geometry	12
4. Existence for small q 's in the case $p \in (2, 6)$	13
5. Existence for all q 's in the case $p \in (3, 6)$	16
5.1. The case $p \in [4, 6)$.	16
5.2. The case $p \in (3, 4)$	17
6. The behaviour as $a \rightarrow 0$ in the radial case	19
Appendix A. Properties of solutions and nonexistence	21
A.1. Regularity of the solutions.	21
A.2. A useful identity	22
A.3. The Pohožaev identity	22
A.4. A nonexistence result	24
Appendix B. Proof of Lemma 4.5	24
References	29

2010 *Mathematics Subject Classification.* 35J48, 35J50, 35Q60.

Key words and phrases. Elliptic systems, Schrödinger-Bopp-Podolsky equations, Variational Methods, Standing waves solutions.

P. d'Avenia was partially supported by a grant of the group GNAMPA of INdAM and by FRA2016 of Politecnico di Bari. G. Siciliano was partially supported by Fapesp, CNPq and Capes, Brazil. This work was partially carried out during a stay of P. d'Avenia at the Universidade de São Paulo. He would like to express his deep gratitude to the Instituto de Matemática e Estatística, Departamento de Matemática, for the warm hospitality.

1. INTRODUCTION

In this paper we consider the system

$$(1.1) \quad \begin{cases} -\Delta u + \omega u + q^2 \phi u = |u|^{p-2}u \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi u^2 \end{cases} \quad \text{in } \mathbb{R}^3$$

where $u, \phi : \mathbb{R}^3 \rightarrow \mathbb{R}$, $\omega, a > 0$, $q \neq 0$.

Such a system appears when we couple a Schrödinger field $\psi = \psi(t, x)$ with its electromagnetic field in the Bopp-Podolsky electromagnetic theory, and, in particular, in the electrostatic case for standing waves $\psi(t, x) = e^{i\omega t}u(x)$, see Section 2 for more details.

The Bopp-Podolsky theory, developed by Bopp [11], and independently by Podolsky [34], is a second order gauge theory for the electromagnetic field. As the Mie theory [33] and its generalizations given by Born and Infeld [12–15], it was introduced to solve the so called *infinity problem* that appears in the classical Maxwell theory. In fact, by the well known Gauss law (or Poisson equation), the electrostatic potential ϕ for a given charge distribution whose density is ρ satisfies the equation

$$(1.2) \quad -\Delta \phi = \rho \quad \text{in } \mathbb{R}^3.$$

If $\rho = 4\pi\delta_{x_0}$, with $x_0 \in \mathbb{R}^3$, the fundamental solution of (1.2) is $\mathcal{G}(x - x_0)$, where

$$\mathcal{G}(x) = \frac{1}{|x|},$$

and the electrostatic energy is

$$\mathcal{E}_M(\mathcal{G}) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla \mathcal{G}|^2 = +\infty.$$

Thus, equation (1.2) is *replaced* by

$$-\operatorname{div} \left(\frac{\nabla \phi}{\sqrt{1 - |\nabla \phi|^2}} \right) = \rho \quad \text{in } \mathbb{R}^3$$

in the Born-Infeld theory and by

$$-\Delta \phi + a^2 \Delta^2 \phi = \rho \quad \text{in } \mathbb{R}^3$$

in the Bopp-Podolsky one. In both cases, if $\rho = 4\pi\delta_{x_0}$, we are able to write explicitly the solutions of the respective equations and to see that their energy is finite. In particular, when we consider the operator $-\Delta + a^2 \Delta^2$, we have that $\mathcal{K}(x - x_0)$, with

$$\mathcal{K}(x) := \frac{1 - e^{-|x|/a}}{|x|},$$

is the fundamental solution of the equation

$$-\Delta \phi + a^2 \Delta^2 \phi = 4\pi\delta_{x_0},$$

it has no singularity in x_0 since it satisfies

$$\lim_{x \rightarrow x_0} \mathcal{K}(x - x_0) = \frac{1}{a},$$

and its energy is

$$\mathcal{E}_{BP}(\mathcal{K}) = \frac{1}{2} \int_{\mathbb{R}^3} |\nabla \mathcal{K}|^2 + \frac{a^2}{2} \int_{\mathbb{R}^3} |\Delta \mathcal{K}|^2 < +\infty$$

(see Section 3.1 for more details).

Moreover the Bopp-Podolsky theory may be interpreted as an effective theory for short distances (see [27]) and for large distances it is experimentally indistinguishable from the Maxwell one. Thus

the Bopp-Podolsky parameter $a > 0$, which has dimension of the inverse of mass, can be interpreted as a cut-off distance or can be linked to an effective radius for the electron. For more physical details we refer the reader to the recent papers [7, 10, 17, 18, 20, 21] and to references therein.

Finally we point out that the operator $-\Delta + \Delta^2$ appears also in other different interesting mathematical and physical situations (see [8, 25] and their references).

Before stating our results, few preliminaries are in order. We introduce here the space \mathcal{D} as the completion of $C_c^\infty(\mathbb{R}^3)$ with respect to the norm $\sqrt{\|\nabla\phi\|_2^2 + a^2\|\Delta\phi\|_2^2}$. We refer to Section 3 for more properties on this space.

Then, fixed $a > 0$ and $q \neq 0$, we say that a pair $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ is a solution of (1.1) if

$$\begin{aligned} \int_{\mathbb{R}^3} \nabla u \nabla v + \omega \int_{\mathbb{R}^3} uv + q^2 \int_{\mathbb{R}^3} \phi uv &= \int_{\mathbb{R}^3} |u|^{p-2} uv \quad \text{for all } v \in H^1(\mathbb{R}^3), \\ \int_{\mathbb{R}^3} \nabla \phi \nabla \xi + a^2 \int_{\mathbb{R}^3} \Delta \phi \Delta \xi &= 4\pi \int_{\mathbb{R}^3} \phi u^2 \quad \text{for all } \xi \in \mathcal{D}. \end{aligned}$$

By standard arguments the solutions we find are easily seen to be positive. Moreover we say that a solution (u, ϕ) is nontrivial whenever $u \not\equiv 0$.

Then our results can be stated as follows.

Theorem 1.1 (Existence for small charges). *If $p \in (2, 6)$ then there exists $q_* > 0$ such that, for all $q \in (-q_*, q_*) \setminus \{0\}$, problem (1.1) admits a nontrivial solution.*

The above existence result can be extended to any value of q provided that a restriction on p is made.

Theorem 1.2 (Existence for arbitrary charges). *If $p \in (3, 6)$ then, for all $q \neq 0$, problem (1.1) admits a nontrivial solution.*

To prove our existence results we use Variational Methods. Indeed the solutions can be found as critical points of a smooth functional \mathcal{J}_q defined on $H^1(\mathbb{R}^3)$. However we need to face with the following difficulties. The geometry of the functional strongly depends on the values of the parameters p and q , that may allow or prevents the existence of critical points. Moreover, a fundamental tool in Critical Point Theory is the following compactness condition: we say that the functional \mathcal{J}_q satisfies the *Palais-Smale* ((PS) for short) condition if any (PS) sequence $\{u_n\} \subset H^1(\mathbb{R}^3)$, that is a sequence satisfying

$$\{\mathcal{J}_q(u_n)\} \text{ bounded and } \mathcal{J}_q'(u_n) \rightarrow 0 \text{ in } H^{-1}(\mathbb{R}^3) \text{ as } n \rightarrow +\infty,$$

admits a convergent subsequence.

In particular, to find a bounded (PS) sequence in the case $p \in (2, 3]$, a suitable truncation, introduced in [30] and already used successfully in recent papers as [3, 26, 31], is performed. Once we have a bounded (PS) sequence, the invariance by translations of the problem clearly leads to a second difficulty: the lack of compactness. To overcome this problem a useful Splitting Lemma (see Lemma 4.5) is given.

Another difficulty which appears is due to the fact that the kernel \mathcal{K} is not homogeneous. This makes difficult the use of rescaling of type $t \mapsto u(t^\alpha \cdot)$ and hence arguments as in [35] cannot be used. However we can take some advantage from the fact that $\mathcal{K} \leq \mathcal{G}$.

Let us observe that, due to the invariance of \mathcal{J}_q under the group induced by the action of rotations on $H^1(\mathbb{R}^3)$, we can restrict ourselves to $H_r^1(\mathbb{R}^3)$, the subspace of radial functions, which is a natural constraint: if $u \in H_r^1(\mathbb{R}^3)$ is a critical point of $\mathcal{J}_q|_{H_r^1(\mathbb{R}^3)}$, then it is a critical point for the functional on the entire $H^1(\mathbb{R}^3)$. Then the same results as in Theorem 1.1 and Theorem 1.2 hold in the radial setting (with even a simpler argument in order to recover compactness). Actually

in this case we can say even more: the solutions found tend to solutions of the Schrödinger-Poisson system

$$(1.3) \quad \begin{cases} -\Delta u + \omega u + q^2 \phi u = |u|^{p-2}u & \text{in } \mathbb{R}^3, \\ -\Delta \phi = 4\pi u^2 \end{cases}$$

obtained formally by (1.1) by setting $a = 0$. Indeed we have the following

Theorem 1.3. *For $q \neq 0$ fixed according to the restriction in the Theorems 1.1 and 1.2, let $(u^a, \phi^a) \in H_r^1(\mathbb{R}^3) \times \mathcal{D}_r$ be solutions of (1.1). Then*

$$u^a \rightarrow u^0 \text{ in } H_r^1(\mathbb{R}^3) \quad \text{and} \quad \phi^a \rightarrow \phi^0 \text{ in } D_r^{1,2}(\mathbb{R}^3) \quad \text{as } a \rightarrow 0,$$

where $(u^0, \phi^0) \in H_r^1(\mathbb{R}^3) \times D_r^{1,2}(\mathbb{R}^3)$ is a solution of (1.3).

Let us recall that $D^{1,2}(\mathbb{R}^3)$ is the usual Sobolev space defined as the completion of $C_c^\infty(\mathbb{R}^3)$ with respect to the norm $\|\nabla \phi\|_2$ and $D_r^{1,2}(\mathbb{R}^3), \mathcal{D}_r$ are the respective subspaces of radial functions.

We point out that there is a wide literature on the coupling of matter with its own electromagnetic field for many different situations. See e.g. [3, 24] for the Maxwell theory, [4, 9, 23, 26] for the Born-Infeld one, [19] for the Chern-Simon one, and references therein. However, to the best of our knowledge, this is the first time that a system like (1.1), which involves the Bopp-Podolsky electromagnetic theory, appears in the mathematical literature.

The paper is organised as follows.

In Section 2 the Schrödinger-Bopp-Podolsky system is obtained in the framework of the Abelian Gauge Theories. Then the ansatz of stationary solutions in the electrostatic case is done and (1.1) is deduced.

In Section 3 we give general preliminaries in order to attack our problem. In particular we first define the right spaces in which find the solutions. Then we show rigorously that \mathcal{K} is the fundamental solution of the operator $-\Delta + a^2 \Delta^2$. Probably this result is known, but we were not able to find it in the literature. In particular Lemma 3.3 is interesting of its right. Moreover the smooth energy functional \mathcal{J}_q is defined in such a way that its critical points are exactly solutions of (1.1) and its geometric properties are proved.

In Section 4 we prove Theorem 1.1. Indeed the hypothesis of small charges allows to have the Mountain Pass Geometry for any value of $p \in (2, 6)$. Here the boundedness of the (PS) sequences is obtained by means of the key Proposition 4.3. On the other hand the compactness is recovered by means of the Splitting Lemma 4.5.

Section 5 is devoted to the proof of Theorem 1.2. In this case, that is for $p \in (3, 6)$, the Mountain Pass Geometry holds for every q . However the boundedness of the (PS) sequences is obtained in two different way: it is *standard* if $p \in [4, 6)$ while it is obtained by means of the monotonicity trick (see [29, 37]) if $p \in (3, 4)$. Even now the compactness (and then the existence of a solution) can be recovered by using the Splitting Lemma; nevertheless, just to use a different (and simpler) argument, we prefer to give the proof in the radial setting.

In Section 6 we study the behaviour of the radial solutions with respect to a . After proving Lemma 6.1 which may be of some interest in other contexts, we prove Theorem 1.3.

In Appendix A we collect few facts concerning the regularity of our solutions, we prove the Pohožaev identity and give the proof of some nonexistence results in the cases $p \geq 6$ and $p \leq 2$. In particular, to achieve our goal, in the case $p \leq 2$ we use an interesting identity, obtained in Section A.2, which could be useful for further developments.

Finally in Appendix B we give the proof of the Splitting Lemma 4.5.

Notations. As a matter of notations through the paper

- we denote with $\|\cdot\|_p$ the usual $L^p(\mathbb{R}^3)$ norm;

- $\langle \cdot, \cdot \rangle$ is the scalar product in $H^1(\mathbb{R}^3)$ which gives rise to the norm $\|\cdot\| = \sqrt{\|\nabla \cdot\|_2^2 + \omega \|\cdot\|_2^2}$;
- $p' = p/(p-1)$ is the conjugate exponent of p ;
- we use the symbol $o_n(1)$ for a vanishing sequence in the specified space;
- we use C, C_1, C_2, \dots to denote suitable positive constants whose value may also change from line to line;
- if not specified, the domain of the integrals is \mathbb{R}^3 .

Other notations will be introduced whenever we need. Moreover, for simplicity, from now on we will consider positive q 's.

2. DEDUCTION OF THE SCHRÖDINGER-BOPP-PODOLSKY SYSTEM

Let us consider the nonlinear Schrödinger Lagrangian density

$$\mathcal{L}_{\text{Sc}} = i\hbar\bar{\psi}\partial_t\psi - \frac{\hbar^2}{2m}|\nabla\psi|^2 + \frac{2}{p}|\psi|^p,$$

where $\psi : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{C}$, $\hbar, m, p > 0$, and let (ϕ, \mathbf{A}) be the gauge potential of the electromagnetic field (\mathbf{E}, \mathbf{H}) , namely $\phi : \mathbb{R}^3 \rightarrow \mathbb{R}$ and $\mathbf{A} : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ satisfy

$$\mathbf{E} = -\nabla\phi - \frac{1}{c}\partial_t\mathbf{A}, \quad \mathbf{H} = \nabla \times \mathbf{A}.$$

The coupling of the field ψ with the electromagnetic field (\mathbf{E}, \mathbf{H}) through the minimal coupling rule, namely the study of the interaction between ψ and its own electromagnetic field, can be obtained replacing in \mathcal{L}_{Sc} the derivatives ∂_t and ∇ respectively with the covariant ones

$$D_t = \partial_t + \frac{iq}{\hbar}\phi, \quad \mathbf{D} = \nabla - \frac{iq}{\hbar c}\mathbf{A},$$

q being a *coupling* constant. This leads to consider

$$\begin{aligned} \mathcal{L}_{\text{CSc}} &= i\hbar\bar{\psi}D_t\psi - \frac{\hbar^2}{2m}|\mathbf{D}\psi|^2 + \frac{2}{p}|\psi|^p \\ &= i\hbar\bar{\psi}\left(\partial_t + \frac{iq}{\hbar}\phi\right)\psi - \frac{\hbar^2}{2m}\left|\left(\nabla - \frac{iq}{\hbar c}\mathbf{A}\right)\psi\right|^2 + \frac{2}{p}|\psi|^p. \end{aligned}$$

Now, to get the total Lagrangian density, we have to add to \mathcal{L}_{CSc} the Lagrangian density of the electromagnetic field.

The Bopp-Podolsky Lagrangian density (see [34, Formula (3.9)]) is

$$\begin{aligned} \mathcal{L}_{\text{BP}} &= \frac{1}{8\pi} \left\{ |\mathbf{E}|^2 - |\mathbf{H}|^2 + a^2 \left[(\text{div } \mathbf{E})^2 - \left| \nabla \times \mathbf{H} - \frac{1}{c}\partial_t\mathbf{E} \right|^2 \right] \right\} \\ &= \frac{1}{8\pi} \left\{ \left| \nabla\phi + \frac{1}{c}\partial_t\mathbf{A} \right|^2 - |\nabla \times \mathbf{A}|^2 \right. \\ &\quad \left. + a^2 \left[\left(\Delta\phi + \frac{1}{c}\text{div } \partial_t\mathbf{A} \right)^2 - \left| \nabla \times \nabla \times \mathbf{A} + \frac{1}{c}\partial_t(\nabla\phi + \frac{1}{c}\partial_t\mathbf{A}) \right|^2 \right] \right\}. \end{aligned}$$

Thus the total action is

$$\mathcal{S}(\psi, \phi, \mathbf{A}) = \int \mathcal{L} dx dt$$

where $\mathcal{L} := \mathcal{L}_{\text{CSc}} + \mathcal{L}_{\text{BP}}$ is the total Lagrangian density.

The Euler-Lagrange equations of \mathcal{S} are given by

$$\left\{ \begin{aligned} & i\hbar \left(\partial_t + \frac{iq}{\hbar} \phi \right) \psi + \frac{\hbar^2}{2m} \left(\nabla - \frac{iq}{\hbar c} \mathbf{A} \right)^2 \psi + |\psi|^{p-2} \psi = 0 \\ & -\operatorname{div} \left(\nabla \phi + \frac{1}{c} \partial_t \mathbf{A} \right) + a^2 \left[\Delta \left(\Delta \phi + \frac{1}{c} \operatorname{div} \partial_t \mathbf{A} \right) - \frac{1}{c} \partial_t \operatorname{div} \left(\nabla \times \nabla \times \mathbf{A} + \frac{1}{c} \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) \right) \right] \\ & \quad = 4\pi q |\psi|^2 \\ & -\frac{\hbar q}{mc} \Im \left[\left(\nabla \bar{\psi} + \frac{iq}{\hbar c} \mathbf{A} \bar{\psi} \right) \psi \right] - \frac{1}{4\pi} \left\{ \frac{1}{c} \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) + \nabla \times \nabla \times \mathbf{A} \right\} \\ & \quad + \frac{a^2}{4\pi} \left[\frac{1}{c} \nabla \partial_t \left(\Delta \phi + \frac{1}{c} \operatorname{div} \partial_t \mathbf{A} \right) - \nabla \times \nabla \times \nabla \times \nabla \times \mathbf{A} - \frac{1}{c^2} \partial_{tt} \nabla \times \nabla \times \mathbf{A} \right. \\ & \quad \left. - \frac{1}{c} \nabla \times \nabla \times \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) - \frac{1}{c^3} \partial_{ttt} (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) \right] = 0. \end{aligned} \right.$$

If we consider $\psi(t, x) = e^{iS(t, x)} u(t, x)$ with $S, u : \mathbb{R} \times \mathbb{R}^3 \rightarrow \mathbb{R}$ the Euler-Lagrange equations are

$$\left\{ \begin{aligned} & -\frac{\hbar^2}{2m} \Delta u + \left[\frac{\hbar^2}{2m} \left| \nabla S - \frac{q}{\hbar c} \mathbf{A} \right|^2 + \hbar \partial_t S + q\phi \right] u = |u|^{p-2} u \\ & \partial_t u^2 + \frac{\hbar}{m} \operatorname{div} \left[\left(\nabla S - \frac{q}{\hbar c} \mathbf{A} \right) u^2 \right] = 0 \\ & -\operatorname{div} \left(\nabla \phi + \frac{1}{c} \partial_t \mathbf{A} \right) + a^2 \left[\Delta \left(\Delta \phi + \frac{1}{c} \operatorname{div} \partial_t \mathbf{A} \right) - \frac{1}{c} \partial_t \operatorname{div} \left(\nabla \times \nabla \times \mathbf{A} + \frac{1}{c} \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) \right) \right] \\ & \quad = 4\pi q |u|^2 \\ & \frac{\hbar q}{mc} \left(\nabla S - \frac{q}{\hbar c} \mathbf{A} \right) u^2 - \frac{1}{4\pi} \left\{ \frac{1}{c} \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) + \nabla \times \nabla \times \mathbf{A} \right\} \\ & \quad + \frac{a^2}{4\pi} \left[\frac{1}{c} \nabla \partial_t \left(\Delta \phi + \frac{1}{c} \operatorname{div} \partial_t \mathbf{A} \right) - \nabla \times \nabla \times \nabla \times \nabla \times \mathbf{A} - \frac{1}{c^2} \partial_{tt} \nabla \times \nabla \times \mathbf{A} \right. \\ & \quad \left. - \frac{1}{c} \nabla \times \nabla \times \partial_t (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) - \frac{1}{c^3} \partial_{ttt} (\nabla \phi + \frac{1}{c} \partial_t \mathbf{A}) \right] = 0. \end{aligned} \right.$$

Finally, if we consider standing waves $\psi(t, x) = e^{i\omega t/\hbar} u(x)$ in the purely electrostatic case ($\phi = \phi(x)$ and $\mathbf{A} = \mathbf{0}$), the second and fourth equation are satisfied and we get

$$(SBP) \quad \begin{cases} -\frac{\hbar^2}{2m} \Delta u + \omega u + q\phi u = |u|^{p-2} u \\ -\Delta \phi + a^2 \Delta^2 \phi = 4\pi q u^2. \end{cases}$$

Normalising the constants \hbar and m and renaming the unknown ϕ it is easy to see that solutions of (1.1) give rise to solutions of (SBP). Hence from now on we will refer to system (1.1).

3. PRELIMINARIES

In this section we give some preliminary results that will be useful for our arguments. In particular we give some fundamental properties on the operator $-\Delta + a^2 \Delta^2$. Then we introduce the functional whose critical points are weak solutions of our problem and we conclude the section showing that, at least for small q 's, such a functional satisfies the geometrical assumptions of the Mountain Pass Theorem.

3.1. The operator $-\Delta + a^2\Delta^2$. Let \mathcal{D} be the completion of $C_c^\infty(\mathbb{R}^3)$ with respect to the norm $\|\cdot\|_{\mathcal{D}}$ induced by the scalar product

$$\langle \varphi, \psi \rangle_{\mathcal{D}} := \int \nabla \varphi \nabla \psi + a^2 \int \Delta \varphi \Delta \psi.$$

Then \mathcal{D} is an Hilbert space continuously embedded into $D^{1,2}(\mathbb{R}^3)$ and consequently in $L^6(\mathbb{R}^3)$. It is interesting to note also the following result.

Lemma 3.1. *The space \mathcal{D} is continuously embedded in $L^\infty(\mathbb{R}^3)$.*

Proof. Let $\varphi \in C_c^\infty(\mathbb{R}^3)$, $x \in \mathbb{R}^3$, and Q be a unitary cube containing x . Arguing as in Brezis [16, Proof of Theorem 9.12], using the Sobolev inequality applied to φ and to $\nabla \varphi$, and since

$$\sum_{i,j} \int \partial_{ij} \varphi \partial_{ij} \varphi = \int \Delta \varphi \Delta \varphi,$$

we have

$$|\varphi(x)| \leq |\bar{\varphi}| + C \|\nabla \varphi\|_{L^6(Q)} \leq C \|\varphi\|_{W^{1,6}(Q)} \leq C \|\varphi\|_{W^{1,6}(\mathbb{R}^3)} \leq C (\|\nabla \varphi\|_2 + \|\Delta \varphi\|_2) \leq C \|\varphi\|_{\mathcal{D}}.$$

Here $\bar{\varphi}$ is the mean of φ on Q and C 's do not depend on Q and φ . Therefore, standard density arguments allow to conclude. \square

The next Lemma gives a useful characterization of the space \mathcal{D} .

Lemma 3.2. *The space $C_c^\infty(\mathbb{R}^3)$ is dense in*

$$\mathcal{A} := \{\phi \in D^{1,2}(\mathbb{R}^3) : \Delta \phi \in L^2(\mathbb{R}^3)\}$$

normed by $\sqrt{\langle \phi, \phi \rangle_{\mathcal{D}}}$ and, therefore, $\mathcal{D} = \mathcal{A}$.

Proof. Let $\phi \in \mathcal{A}$, $\rho \in C_c^\infty(\mathbb{R}^3; \mathbb{R}_+)$, $\|\rho\|_1 = 1$, and $\{\rho_n\} \subset C_c^\infty(\mathbb{R}^3)$ the sequence of mollifiers given by $\rho_n(x) = n^3 \rho(nx)$. Define $\phi_n := \rho_n * \phi \in C^\infty(\mathbb{R}^3)$. Since, recalling the well known properties of the mollifiers,

$$\partial_i \phi_n = \rho_n * \partial_i \phi \in L^2(\mathbb{R}^3), \quad i = 1, 2, 3, \quad \Delta \phi_n = \rho_n * \Delta \phi \in L^2(\mathbb{R}^3),$$

and

$$\|\nabla \phi_n - \nabla \phi\|_2 \rightarrow 0, \quad \|\Delta \phi_n - \Delta \phi\|_2 \rightarrow 0,$$

we have

$$(3.1) \quad \phi_n \in C^\infty(\mathbb{R}^3) \cap \mathcal{A} \quad \text{and} \quad \|\phi_n - \phi\|_{\mathcal{D}} \rightarrow 0.$$

Let now $\xi \in C^\infty(\mathbb{R}^3) \cap \mathcal{A}$, $\zeta \in C_c^\infty(\mathbb{R}^3; [0, 1])$ with $\zeta(x) = 1$ in $B(0, 1)$, $\text{supp}(\zeta) \subset B(0, 2)$ and define

$$\xi_n := \zeta(\cdot/n) \xi \in C_c^\infty(\mathbb{R}^3).$$

We have

$$\begin{aligned} \nabla \xi_n &= \zeta(\cdot/n) \nabla \xi + \frac{1}{n} \xi \nabla \zeta(\cdot/n), \\ \Delta \xi_n &= \zeta(\cdot/n) \Delta \xi + \frac{2}{n} \nabla \xi \nabla \zeta(\cdot/n) + \frac{1}{n^2} \xi \Delta \zeta(\cdot/n). \end{aligned}$$

Noticing that

$$\frac{1}{n^2} \int \xi^2(x) \left| \nabla \zeta\left(\frac{x}{n}\right) \right|^2 \leq \frac{1}{n^2} \left(\int_{|x| \geq n} \xi^6 \right)^{1/3} \left(\int \left| \nabla \zeta\left(\frac{x}{n}\right) \right|^3 \right)^{2/3} = C \left(\int_{|x| \geq n} \xi^6 \right)^{1/3} \rightarrow 0$$

and, analogously,

$$\frac{2}{n} \nabla \xi \nabla \zeta(\cdot/n), \quad \frac{1}{n^2} \xi \Delta \zeta(\cdot/n) \rightarrow 0 \quad \text{in } L^2(\mathbb{R}^3),$$

as $n \rightarrow +\infty$, we infer

$$\begin{aligned}\|\nabla \xi - \nabla \xi_n\|_2^2 &\leq 2\|(1 - \zeta(\cdot/n))\partial_i \xi\|_2^2 + o_n(1) = o_n(1) \\ \|\Delta \xi - \Delta \xi_n\|_2^2 &\leq 2\|(1 - \zeta(\cdot/n))\Delta \xi\|_2^2 + o_n(1) = o_n(1)\end{aligned}$$

showing that $\|\xi_n - \xi\|_{\mathcal{D}} \rightarrow 0$. This joint with (3.1) concludes the proof. \square

For every fixed $u \in H^1(\mathbb{R}^3)$, the Riesz Theorem implies that there exists a unique solution $\phi_u \in \mathcal{D}$ of the second equation in (1.1). To write *explicitly* such a solution (see also [34, Formula (2.6)]), we consider

$$\mathcal{K}(x) = \frac{1 - e^{-|x|/a}}{|x|}.$$

We have the following fundamental properties.

Lemma 3.3. *For all $y \in \mathbb{R}^3$, $\mathcal{K}(\cdot - y)$ solves in the sense of distributions*

$$-\Delta \phi + a^2 \Delta^2 \phi = 4\pi \delta_y.$$

Moreover

- (i) if $f \in L_{\text{loc}}^1(\mathbb{R}^3)$ and, for a.e. $x \in \mathbb{R}^3$, the map $y \in \mathbb{R}^3 \mapsto f(y)/|x - y|$ is summable, then $\mathcal{K} * f \in L_{\text{loc}}^1(\mathbb{R}^3)$;
- (ii) if $f \in L^p(\mathbb{R}^3)$ with $1 \leq p < 3/2$, then $\mathcal{K} * f \in L^q(\mathbb{R}^3)$ for $q \in (3p/(3 - 2p), +\infty]$.

In both cases $\mathcal{K} * f$ solves

$$(3.2) \quad -\Delta \phi + a^2 \Delta^2 \phi = 4\pi f$$

in the sense of distributions and we have the following distributional derivatives

$$\nabla(\mathcal{K} * f) = (\nabla \mathcal{K}) * f \quad \text{and} \quad \Delta(\mathcal{K} * f) = (\Delta \mathcal{K}) * f \quad \text{a.e. in } \mathbb{R}^3.$$

Proof. Let us consider for simplicity $y = 0$ and prove that for every $\varphi \in C_c^\infty(\mathbb{R}^3)$

$$-\int \mathcal{K} \Delta \varphi + a^2 \int \mathcal{K} \Delta^2 \varphi = 4\pi \varphi(0).$$

Of course it is enough to show that

$$(3.3) \quad \lim_{r \rightarrow 0^+} I(r) = 4\pi \varphi(0)$$

where

$$I(r) := -\int_{|x|>r} \mathcal{K} \Delta \varphi + a^2 \int_{|x|>r} \mathcal{K} \Delta^2 \varphi.$$

Since φ has compact support, we consider the annulus $A := \{x \in \mathbb{R}^3 : r < |x| < R\}$ for R large enough and a standard integration by parts shows that

$$\begin{aligned}I(r) &= -\int_A \mathcal{K} \Delta \varphi + a^2 \int_A \mathcal{K} \Delta^2 \varphi \\ &= \int_A \varphi(-\Delta \mathcal{K} + a^2 \Delta^2 \mathcal{K}) + \int_{|x|=r} \varphi(\nabla \mathcal{K} - a^2 \nabla \Delta \mathcal{K}) \cdot \nu + \int_{|x|=r} \mathcal{K}(a^2 \nabla(\Delta \varphi) - \nabla \varphi) \cdot \nu \\ &\quad + a^2 \int_{|x|=r} \Delta \mathcal{K} \nabla \varphi \cdot \nu - a^2 \int_{|x|=r} \Delta \varphi \nabla \mathcal{K} \cdot \nu\end{aligned}$$

where ν is the unit outward normal to A . Since φ is continuous, $(a^2\nabla(\Delta\varphi) - \nabla\varphi) \cdot \nu$, $\nabla\varphi \cdot \nu$, and $\Delta\varphi$ are bounded, \mathcal{K} can be extended continuously in 0 by setting $\mathcal{K}(0) = 1/a$, and, for $x \neq 0$,

$$(3.4) \quad \nabla\mathcal{K} = -\frac{x}{|x|^3} + \frac{x}{|x|^3} \left(\frac{|x|}{a} + 1 \right) e^{-|x|/a},$$

$$(3.5) \quad \begin{aligned} \Delta\mathcal{K} &= -\frac{e^{-|x|/a}}{a^2|x|}, \\ \nabla\Delta\mathcal{K} &= \frac{x}{a^2|x|^3} \left(\frac{|x|}{a} + 1 \right) e^{-|x|/a}, \\ \Delta^2\mathcal{K} &= \frac{1}{a^2} \left(\Delta\mathcal{K} + \operatorname{div} \frac{x}{|x|^3} \right), \end{aligned}$$

we have

$$\begin{aligned} -\Delta\mathcal{K} + a^2\Delta^2\mathcal{K} &= 0 \text{ in } A \implies \int_A \varphi(-\Delta\mathcal{K} + a^2\Delta^2\mathcal{K}) = 0, \\ (\nabla\mathcal{K} - a^2\nabla\Delta\mathcal{K}) \cdot \nu &= \frac{1}{r^2} \text{ on } |x| = r \implies \int_{|x|=r} \varphi(\nabla\mathcal{K} - a^2\nabla\Delta\mathcal{K}) \cdot \nu \\ &= \int_S \varphi(r\sigma) d\sigma \rightarrow 4\pi\varphi(0) \text{ as } r \rightarrow 0^+, \\ \mathcal{K} \leq 1/a &\implies \int_{|x|=r} \mathcal{K}(a^2\nabla(\Delta\varphi) - \nabla\varphi) \cdot \nu \rightarrow 0 \text{ as } r \rightarrow 0^+, \\ |\Delta\mathcal{K}| \leq 1/(a^2|x|) &\implies \int_{|x|=r} \Delta\mathcal{K} \nabla\varphi \cdot \nu \rightarrow 0 \text{ as } r \rightarrow 0^+, \\ \nabla\mathcal{K} \cdot \nu &= \frac{1}{r^2} \left[1 - \left(\frac{r}{a} + 1 \right) e^{-\frac{r}{a}} \right] \text{ on } |x| = r \implies \int_{|x|=r} \Delta\varphi \nabla\mathcal{K} \cdot \nu \rightarrow 0 \text{ as } r \rightarrow 0^+, \end{aligned}$$

where S is the unit sphere in \mathbb{R}^3 . Thus (3.3) is proved.

To get (i), we observe that, by Fubini Theorem, for all balls $B \subset \mathbb{R}^3$

$$\int_B |\mathcal{K} * f| \leq \int_{\mathbb{R}^3} \left(\int_B \mathcal{K}(x-y) dx \right) |f(y)| dy$$

and, since $\mathcal{K} \leq |\cdot|^{-1}$, we can conclude arguing as in [32, Proof of Theorem 6.21].

Since $\mathcal{K} \in L^\tau(\mathbb{R}^3)$ for $\tau \in (3, +\infty]$, the Young inequality (see e.g. [32, Inequality (4), p 99]) allows to get (ii), since

$$\frac{1}{q} = \frac{1}{p} + \frac{1}{\tau} - 1 < \frac{1}{p} + \frac{1}{3} - 1 = \frac{3-2p}{3p}.$$

Moreover the fact that $\mathcal{K} * f$ solves (3.2) in the sense of distributions, namely, for all $\varphi \in C_c^\infty(\mathbb{R}^3)$

$$-\int (\mathcal{K} * f) \Delta\varphi + a^2 \int (\mathcal{K} * f) \Delta^2\varphi = 4\pi \int f\varphi,$$

is a consequence of the Fubini Theorem and of the first part of this Lemma.

To conclude, let us consider, for instance, the assumptions in (i). The proof of the remaining case is similar.

We claim that the functions $(\nabla\mathcal{K}) * f$ and $(\Delta\mathcal{K}) * f$ are well defined a.e. in \mathbb{R}^3 . In fact, by (3.4), we have that, for every $i = 1, 2, 3$,

$$(3.6) \quad |\partial_i \mathcal{K}(x)| \leq \frac{1}{|x|^2} \left(1 - e^{-|x|/a} - \frac{|x|}{a} e^{-|x|/a} \right) \leq C \quad \text{for } |x| \text{ small}$$

and, since

$$(3.7) \quad |\partial_i \mathcal{K}(x)| \leq \frac{1}{|x|^2} + \frac{1}{|x|^2} \left(\frac{|x|}{a} + 1 \right) e^{-|x|/a},$$

then, in particular,

$$|\partial_i \mathcal{K}(x)| \leq \frac{2}{|x|^2} + \frac{1}{a|x|} \leq \frac{C}{|x|} \quad \text{for } |x| \text{ large.}$$

Thus, if $r > 0$ is sufficiently large, for a.e. x in \mathbb{R}^3 , using the summability of the map $y \in \mathbb{R}^3 \mapsto f(y)/|x-y|$, we deduce

$$\begin{aligned} |[\partial_i \mathcal{K} * f](x)| &\leq \int_{B_r(x)} |\partial_i \mathcal{K}(x-y)| |f(y)| dy + \int_{B_r^c(x)} |\partial_i \mathcal{K}(x-y)| |f(y)| dy \\ &\leq C \left[\int_{B_r(x)} |f(y)| dy + \int_{B_r^c(x)} \frac{|f(y)|}{|x-y|} dy \right] < +\infty. \end{aligned}$$

Moreover, by (3.5), $|\Delta \mathcal{K}| \leq 1/(a^2 |\cdot|)$ and so, arguing again as in [32, Proof of Theorem 6.21] we get the claim.

Then, since

$$|\mathcal{K}(x)|, |\partial_i \mathcal{K}(x)|, |\Delta \mathcal{K}(x)| \leq \frac{C}{|x|},$$

and the map $y \in \mathbb{R}^3 \mapsto f(y)/|x-y|$ is summable, [32, Theorem 6.21] implies that

$$\mathcal{K} * f, \partial_i \mathcal{K} * f, \Delta \mathcal{K} * f \in L_{\text{loc}}^1(\mathbb{R}^3)$$

and so, for all $\varphi \in C_c^\infty(\mathbb{R}^3)$,

$$\begin{aligned} (x, y) &\mapsto \partial_i \varphi(x) \mathcal{K}(x-y) f(y) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3), \\ (x, y) &\mapsto \Delta \varphi(x) \mathcal{K}(x-y) f(y) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3), \\ (x, y) &\mapsto \varphi(x) \partial_i \mathcal{K}(x-y) f(y) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3), \\ (x, y) &\mapsto \varphi(x) \Delta \mathcal{K}(x-y) f(y) \in L^1(\mathbb{R}^3 \times \mathbb{R}^3). \end{aligned}$$

Hence, by Fubini's Theorem and using a limit argument as in the first part of this proof we have that for all $\varphi \in C_c^\infty(\mathbb{R}^3)$

$$\int (\mathcal{K} * f) \partial_i \varphi = - \int \varphi (\partial_i \mathcal{K}) * f \quad i = 1, 2, 3$$

and

$$\int (\mathcal{K} * f) \Delta \varphi = \int \varphi (\Delta \mathcal{K}) * f.$$

The proof is thereby completed. □

Then, if we fix $u \in H^1(\mathbb{R}^3)$, the unique solution in \mathcal{D} of the second equation in (1.1) is

$$(3.8) \quad \phi_u := \mathcal{K} * u^2.$$

Actually the following useful properties hold.

Lemma 3.4. *For every $u \in H^1(\mathbb{R}^3)$ we have:*

- (i) for every $y \in \mathbb{R}^3$, $\phi_{u(\cdot+y)} = \phi_u(\cdot+y)$;
- (ii) $\phi_u \geq 0$;
- (iii) for every $s \in (3, +\infty]$, $\phi_u \in L^s(\mathbb{R}^3) \cap C_0(\mathbb{R}^3)$;
- (iv) for every $s \in (3/2, +\infty]$, $\nabla \phi_u = \nabla \mathcal{K} * u^2 \in L^s(\mathbb{R}^3) \cap C_0(\mathbb{R}^3)$;
- (v) $\phi_u \in \mathcal{D}$;
- (vi) $\|\phi_u\|_6 \leq C \|u\|^2$;

(vii) ϕ_u is the unique minimizer of the functional

$$E(\phi) = \frac{1}{2} \|\nabla \phi\|_2^2 + \frac{a^2}{2} \|\Delta \phi\|_2^2 - \int \phi u^2, \quad \phi \in \mathcal{D}.$$

Moreover

(viii) if $v_n \rightharpoonup v$ in $H^1(\mathbb{R}^3)$, then $\phi_{v_n} \rightharpoonup \phi_v$ in \mathcal{D} .

Proof. Let us fix $u \in H^1(\mathbb{R}^3)$. Items (i) and (ii) are obvious.

Being $\mathcal{K} \in L^\tau(\mathbb{R}^3)$ for $\tau \in (3, +\infty]$, by well known properties of the convolution product, we have (iii).

Moreover, since $\nabla \mathcal{K}$ is bounded near 0 (see (3.6)) and, by (3.7), decays as $|\cdot|^{-2}$ at infinity, then $\nabla \mathcal{K} \in L^\tau(\mathbb{R}^3)$ for $\tau \in (3/2, +\infty]$ and so we get (iv).

Property (v) holds since $\nabla \phi_u \in L^2(\mathbb{R}^3)$ and $\Delta \phi_u = \Delta \mathcal{K} * u^2 \leq 1/(a^2 |\cdot|) * u^2 \in L^2(\mathbb{R}^3)$.

Multiplying the second equation in (1.1) by the solution ϕ_u , integrating and using Lemma 3.1, we find

$$\|\phi_u\|_{\mathcal{D}}^2 \leq C \|u^2\|_1 \|\phi_u\|_\infty \leq C \|u\|^2 \|\phi_u\|_{\mathcal{D}}$$

and then

$$\|\phi_u\|_6 \leq C \|\phi_u\|_{\mathcal{D}} \leq C \|u\|^2.$$

obtaining (vi). Property (vii) is also trivial.

We conclude observing that for every $\varphi \in C_c^\infty(\mathbb{R}^3)$ we have

$$\langle \phi_{v_n}, \varphi \rangle_{\mathcal{D}} = 4\pi \int v_n^2 \varphi \rightarrow 4\pi \int v^2 \varphi = \langle \phi_v, \varphi \rangle_{\mathcal{D}}$$

and we obtain (viii) by density. \square

3.2. The functional setting. It is easy to see that the critical points of the C^1 functional

$$F_q(u, \phi) = \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 + \frac{q^2}{2} \int \phi u^2 - \frac{q^2}{16\pi} \|\nabla \phi\|_2^2 - \frac{a^2 q^2}{16\pi} \|\Delta \phi\|_2^2 - \frac{1}{p} \|u\|_p^p$$

on $H^1(\mathbb{R}^3) \times \mathcal{D}$ are weak solutions of (1.1). Indeed if $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ is a critical point of F_q then

$$0 = \partial_u F_q(u, \phi)[v] = \int \nabla u \nabla v + \omega \int uv + q^2 \int \phi uv - \int |u|^{p-2} uv, \quad \text{for all } v \in H^1(\mathbb{R}^3),$$

and

$$0 = \partial_\phi F_q(u, \phi)[\xi] = \frac{q^2}{2} \int u^2 \xi - \frac{q^2}{8\pi} \int \nabla \phi \nabla \xi - \frac{a^2 q^2}{8\pi} \int \Delta \phi \Delta \xi, \quad \text{for all } \xi \in \mathcal{D}.$$

However the functional F_q is strongly unbounded from below and above and hence the usual techniques of the Critical Point Theory cannot be used immediately. Hence we adopt a reduction procedure which is successfully used also with other system of equations involving the coupling between matter and electromagnetic field. Here we just revise the main argument. We refer the reader to [5, 6] for the details. Then, first of all one observes that actually $\partial_\phi F_q$ is a C^1 function. Thus, if G_Φ is the graph of the map $\Phi : u \in H^1(\mathbb{R}^3) \mapsto \phi_u \in \mathcal{D}$, an application of the Implicit Function Theorem gives

$$G_\Phi = \{(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D} : \partial_\phi F_q(u, \phi) = 0\} \quad \text{and} \quad \Phi \in C^1(H^1(\mathbb{R}^3); \mathcal{D}).$$

Then we define the *reduced* functional

$$\mathcal{J}_q(u) := F_q(u, \Phi(u)) = \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 + \frac{q^2}{4} \int \phi_u u^2 - \frac{1}{p} \|u\|_p^p,$$

which is of class C^1 on $H^1(\mathbb{R}^3)$ and, for all $u, v \in H^1(\mathbb{R}^3)$

$$\begin{aligned}\mathcal{J}'_q(u)[v] &= \partial_u F_q(u, \Phi(u))[v] + \partial_\phi F_q(u, \Phi(u)) \circ \Phi'(u)[v] \\ &= \partial_u F_q(u, \Phi(u))[v] \\ &= \int \nabla u \nabla v + \omega \int uv + q^2 \int \phi_u uv - \int |u|^{p-2} uv.\end{aligned}$$

Rigorously the functional \mathcal{J}_q should depend also on $a > 0$. However, for the sake of simplicity, we do not write explicitly this dependence, which is deserved in Section 6 where the limit as $a \rightarrow 0$ is considered.

We have that the following statements are equivalent:

- (i) the pair $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ is a critical point of F_q , i.e. (u, ϕ) is a solution of (1.1);
- (ii) u is a critical point of \mathcal{J}_q and $\phi = \phi_u$.

In virtue of this, to solve problem (1.1) is equivalent to find the critical points of \mathcal{J}_q , namely to solve

$$-\Delta u + \omega u + q^2 \phi_u u = |u|^{p-2} u \quad \text{in } \mathbb{R}^3.$$

3.3. The Mountain Pass Geometry. We conclude this section showing that the functional \mathcal{J}_q satisfies the geometrical assumptions of the Mountain Pass Theorem [2].

Lemma 3.5. *The functional \mathcal{J}_q satisfies:*

- (i) $\mathcal{J}_q(0) = 0$;
- (ii) there exist $\delta, \rho > 0$ such that $\mathcal{J}_q(u) > \delta$ for all $u \in H^1(\mathbb{R}^3)$ with $\|u\| = \rho$;
- (iii) there exists $w \in H^1(\mathbb{R}^3)$ with $\|w\| > \rho$ such that $\mathcal{J}_q(w) < 0$ for every $q \neq 0$ if $p \in (3, 6)$, and for q small enough if $p \in (2, 3]$.

Proof. Property (i) is trivial.

Moreover, since $\phi_u = \mathcal{K} * u^2 \geq 0$, by Sobolev inequality we have

$$\mathcal{J}(u) \geq C_1 \|u\|^2 - C_2 \|u\|^p$$

and so, if we take $\rho > 0$ small enough, we get (ii).

To prove (iii), we fix $u \neq 0$ in $H^1(\mathbb{R}^3)$ and distinguish two cases:

Case 1: $p \in (3, 6)$.

Let $u_\tau = \tau^2 u(\tau \cdot)$. We have

$$\begin{aligned}\mathcal{J}_q(u_\tau) &= \frac{\tau^3}{2} \|\nabla u\|_2^2 + \frac{\tau}{2} \omega \|u\|_2^2 + q^2 \frac{\tau^3}{4} \iint \frac{1 - e^{-\frac{|x-y|}{\tau a}}}{|x-y|} u^2(x) u^2(y) - \frac{\tau^{2p-3}}{p} \|u\|_p^p \\ &\leq \frac{\tau^3}{2} \|\nabla u\|_2^2 + \frac{\tau}{2} \omega \|u\|_2^2 + q^2 \frac{\tau^3}{4} \iint \frac{u^2(x) u^2(y)}{|x-y|} - \frac{\tau^{2p-3}}{p} \|u\|_p^p\end{aligned}$$

and the conclusion easily follows considering $\tau \rightarrow +\infty$. Actually, if $p \in (4, 6)$, the simpler curve $u_\tau = \tau u$ also works.

Case 2: $p \in (2, 3]$.

Let $u_\tau = \tau^{\frac{p}{p-2}} u(\tau \cdot)$. We have

$$(3.9) \quad \mathcal{J}_q(u_\tau) \leq \frac{\tau^{\frac{p+2}{p-2}}}{2} \|\nabla u\|_2^2 + \frac{\tau^{\frac{6-p}{p-2}}}{2} \omega \|u\|_2^2 + q^2 \frac{\tau^{\frac{10-p}{p-2}}}{4} \iint \frac{u^2(x) u^2(y)}{|x-y|} - \frac{\tau^{\frac{p^2-3p+6}{p-2}}}{p} \|u\|_p^p$$

and since

$$\frac{6-p}{p-2} < \frac{p+2}{p-2} < \frac{p^2-3p+6}{p-2} < \frac{10-p}{p-2},$$

arguing as in [35, Corollary 4.4] we have that, if $q = 0$, the right hand side of 3.9 is unbounded from below (considering $\tau \rightarrow +\infty$) and thus, for q small enough, its infimum is strictly negative and we conclude. \square

Remark 3.6. We observe explicitly that δ, ρ and w do not depend on q , neither on a ; indeed the term involving these two parameters has been successfully thrown away.

4. EXISTENCE FOR SMALL q 'S IN THE CASE $p \in (2, 6)$

In this section we prove an existence result for small q and $p \in (2, 6)$. Actually, as we will see in the next section, such a result will be improved in the case $p \in (3, 6)$: we will be able to find solutions of (1.1) for all $q \neq 0$.

Let us consider, for every $T > 0$, the truncated functional

$$\mathcal{J}_{q,T}(u) := \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 + \frac{q^2}{4} K_T(u) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u^2(x) u^2(y) - \frac{1}{p} \|u\|_p^p$$

where

$$K_T(u) := \chi \left(\frac{\|u\|^2}{T^2} \right)$$

and $\chi \in C_0^\infty(\mathbb{R}, \mathbb{R})$ satisfies

$$\chi(s) := \begin{cases} \chi(s) = 1 & \text{for } s \in [0, 1], \\ 0 \leq \chi \leq 1 & \text{for } s \in [1, 2], \\ \chi(s) = 0 & \text{for } s \in [2, +\infty[, \\ \|\chi'\|_\infty \leq 2. \end{cases}$$

Observe that

$$\begin{aligned} \mathcal{J}'_{q,T}(u)[u] &= \|\nabla u\|_2^2 + \omega \|u\|_2^2 + q^2 K_T(u) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u^2(x) u^2(y) \\ &\quad + \frac{q^2}{2T^2} \chi' \left(\frac{\|u\|^2}{T^2} \right) \|u\|^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u^2(x) u^2(y) - \|u\|_p^p \end{aligned}$$

Arguing as in Lemma 3.5 we have

Lemma 4.1. *The functional $\mathcal{J}_{q,T}$ satisfies the geometric assumption of the Mountain Pass Theorem, namely:*

- (i) $\mathcal{J}_{q,T}(0) = 0$;
- (ii) there exist $\delta, \rho > 0$ such that for all $u \in H^1(\mathbb{R}^3)$ with $\|u\| = \rho$, $\mathcal{J}_{q,T}(u) \geq \delta$;
- (iii) there exists $w \in H^1(\mathbb{R}^3)$ with $\|w\| > \rho$ such that $\mathcal{J}_{q,T}(w) < 0$.

Proof. Property (i) is trivial.

By Sobolev inequality,

$$\mathcal{J}_{q,T}(u) \geq \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 - \frac{1}{p} \|u\|_p^p \geq \frac{1}{2} \|u\|^2 - C \|u\|^p$$

an so, taking ρ small enough, we get (ii).

Finally, let us fix $\psi \in C_0^\infty(\mathbb{R}, \mathbb{R})$ with $\|\psi\| = 1$ and consider $\psi_t := t\psi$ for $t > 0$. If t is sufficiently large, then

$$\mathcal{J}_{q,T}(\psi_t) = \frac{t^2}{2} - \frac{t^p}{p} \|\psi\|_p^p < 0$$

and we get (iii). \square

Remark 4.2. As in Remark 3.6 we have that δ, ρ and w do not depend on q, a, T .

In virtue of the above Lemma we can define the Mountain Pass level for $\mathcal{J}_{q,T}$, namely

$$c_{q,T} := \inf_{\gamma \in \Gamma_{q,T}} \max_{t \in [0,1]} \mathcal{J}_{q,T}(\gamma(t)) > 0,$$

where $\Gamma_{q,T} := \{\gamma \in C([0,1], H^1(\mathbb{R}^3)) : \gamma(0) = 0, \mathcal{J}_{q,T}(\gamma(1)) < 0\}$. By the Ekeland Variational Principle (see also [38]) there exists a (PS) sequence $\{u_n\} \subset H^1(\mathbb{R}^3)$ for $\mathcal{J}_{q,T}$ at level $c_{q,T}$. We can define also

$$c_q := \inf_{\gamma \in \Gamma_q} \max_{t \in [0,1]} \mathcal{J}_q(\gamma(t)) > 0,$$

$\Gamma_q := \{\gamma \in C([0,1], H^1(\mathbb{R}^3)) : \gamma(0) = 0, \mathcal{J}_q(\gamma(1)) < 0\}$, the Mountain Pass level associated to \mathcal{J}_q . Since $\mathcal{J}_{q,T} \leq \mathcal{J}_q$, we have that $c_{q,T} \leq c_q$.

We show now that, for a suitable $\bar{T} > 0$, we have that $\|u_n\| \leq \bar{T}$ and then, being $K_{\bar{T}}(u_n) = 1$, $\{u_n\}$ is also a (PS) sequence for the untruncated functional \mathcal{J}_q , at least for small values of q .

Lemma 4.3. *There exists $\bar{T} > 0$ independent on q and $q_* := q(\bar{T}) > 0$ such that if $q < q_*$, then*

$$\limsup_n \|u_n\| \leq \bar{T}.$$

Proof. Assume by contradiction that

$$(4.1) \quad \limsup_n \|u_n\| > T.$$

Since

$$\begin{aligned} p\mathcal{J}_{q,T}(u_n) - \mathcal{J}'_{q,T}(u_n)[u_n] &= \left(\frac{p}{2} - 1\right) \|u_n\|^2 + \left(\frac{p}{4} - 1\right) q^2 K_T(u_n) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \\ &\quad - \frac{q^2}{2T^2} \chi' \left(\frac{\|u_n\|^2}{T^2} \right) \|u_n\|^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \end{aligned}$$

and so

$$\begin{aligned} (4.2) \quad &\left(\frac{p}{2} - 1\right) \|u_n\|^2 - \|\mathcal{J}'_{q,T}(u_n)\| \|u_n\| \leq \left(\frac{p}{2} - 1\right) \|u_n\|^2 + \mathcal{J}'_{q,T}(u_n)[u_n] \\ &= p\mathcal{J}_{q,T}(u_n) - \left(\frac{p}{4} - 1\right) q^2 K_T(u_n) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \\ &\quad + \frac{q^2}{2T^2} \chi' \left(\frac{\|u_n\|^2}{T^2} \right) \|u_n\|^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \\ &\leq p\mathcal{J}_{q,T}(u_n) + \left| \frac{p}{4} - 1 \right| q^2 K_T(u_n) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \\ &\quad + \frac{q^2}{2T^2} \left| \chi' \left(\frac{\|u_n\|^2}{T^2} \right) \right| \|u_n\|^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y). \end{aligned}$$

Let $w \in H^1(\mathbb{R}^3)$ be as in (iii) of Lemma 4.1. Since $\mathcal{J}_{q,T}(u_n) \rightarrow c_{q,T}$ as $n \rightarrow +\infty$, there exists $\nu \in \mathbb{N}$ such that for all $n \geq \nu$

$$\begin{aligned} \mathcal{J}_{q,T}(u_n) &\leq 2c_{q,T} \leq 2 \max_{t \in [0,1]} \mathcal{J}_{q,T}(tw) \\ &\leq 2 \max_{t \in [0,1]} \left[\frac{t^2}{2} \|w\|^2 - \frac{t^p}{p} \|w\|_p^p \right] + \frac{q^2}{2} \max_{t \in [0,1]} \left[t^4 K_T(tw) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} w^2(x) w^2(y) \right] \\ &:= 2I_1 + \frac{q^2}{2} I_2. \end{aligned}$$

Observe that $I_1 > 0$.

If $t^2 \|w\|^2 > 2T^2$ then $I_2 = 0$ and if $t^2 \|w\|^2 \leq 2T^2$, then

$$I_2 \leq \frac{4T^4}{\|w\|^4} \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} w^2(x) w^2(y).$$

Thus

$$(4.3) \quad \mathcal{J}_{q,T}(u_n) \leq C_1 + q^2 C_2 T^4.$$

Analogously, we get that

$$(4.4) \quad K_T(u_n) \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \leq C_3 \|u_n\|^4 \leq 4C_3 T^4$$

and

$$(4.5) \quad \left| \chi' \left(\frac{\|u_n\|^2}{T^2} \right) \right| \|u_n\|^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u_n^2(x) u_n^2(y) \leq C_4 T^6.$$

Putting (4.3)–(4.5) in (4.2) we have

$$\left(\frac{p}{2} - 1 \right) \|u_n\|^2 - \|\mathcal{J}'_{q,T}(u_n)\| \|u_n\| \leq C_5 + q^2 C_6 T^4.$$

On the other hand, since $\|\mathcal{J}'_{q,T}(u_n)\| \rightarrow 0$ as $n \rightarrow +\infty$ and by (4.1), we have

$$\left(\frac{p}{2} - 1 \right) \|u_n\|^2 - \|\mathcal{J}'_{q,T}(u_n)\| \|u_n\| \geq C_7 T^2 - T$$

and so

$$C_7 T^2 - T \leq C_5 + q^2 C_6 T^4$$

which gives a contradiction if $q = q(T)$ is sufficiently small and for large T . \square

Remark 4.4. Observe that the above proof shows that \bar{T} and $q_* = q(\bar{T})$ do not depend on $a > 0$.

Hence, for every $q \in (0, q_*)$, we have a bounded (PS) sequence $\{u_n\}$, which actually depends on q and a , for the functional $\mathcal{J}_{q,\bar{T}}$ at the level $c_{q,\bar{T}}$.

However since the bound is exactly \bar{T} , which gives $\mathcal{J}_{q,\bar{T}}(u_n) = \mathcal{J}_q(u_n)$ and $c_{q,\bar{T}} = c_q$, we have

$$(4.6) \quad \mathcal{J}_q(u_n) \rightarrow c_q > 0, \quad \mathcal{J}'_q(u_n) \rightarrow 0, \quad \text{as } n \rightarrow +\infty.$$

Moreover we can assume that $u_n \rightharpoonup u_0$ in $H^1(\mathbb{R}^3)$.

The next result helps us to recover the compactness of the bounded (PS) sequence $\{u_n\}$ we have found. For the reader convenience, we give its proof in Appendix B.

Lemma 4.5 (Splitting). *Let $\{u_n\}$ be a bounded (PS) sequence for \mathcal{J}_q at level $d > 0$ and assume that $u_n \rightharpoonup u_0$ in $H^1(\mathbb{R}^3)$. Then, up to subsequences, either u_n strongly converges to u_0 , or there exist $\ell \in \mathbb{N}$, $\{z_n^{(k)}\} \subset \mathbb{R}^3$ for $1 \leq k \leq \ell$, $w_1, \dots, w_\ell \in H^1(\mathbb{R}^3)$ such that*

- (i) $|z_n^{(k)}| \rightarrow +\infty$ for all $1 \leq k \leq \ell$ and $|z_n^{(k)} - z_n^{(h)}| \rightarrow +\infty$ for all $1 \leq k \neq h \leq \ell$, as $n \rightarrow +\infty$;
- (ii) $w_k \neq 0$ and $\mathcal{J}'_q(w_k) = 0$ for all $1 \leq k \leq \ell$;
- (iii) $\left\| u_n - u_0 - \sum_{k=1}^{\ell} w_k(\cdot + z_n^{(k)}) \right\| = o_n(1)$;
- (iv) $d = \mathcal{J}_q(u_0) + \sum_{k=1}^{\ell} \mathcal{J}_q(w_k)$;
- (v) $\mathcal{J}_q(u_n) = \mathcal{J}_q(u_0) + \sum_{k=1}^{\ell} \mathcal{J}_q(w_k) + o_n(1)$.

Then we can easily conclude the proof of Theorem 1.1.

Indeed let $\{u_n\}$ the bounded (PS) sequence for \mathcal{J}_q at level $c_q > 0$ obtained in (4.6). By Lemma 4.5 we have the following possibilities:

- if $u_n \rightarrow u_0$ we have finished being u_0 a solution;
- if there exists $w \in \{u_0, w_1, \dots, w_\ell\}$ such that $\mathcal{J}_q(w) < 0$ we have finished, being w a nontrivial solution;
- if $\mathcal{J}_q(u_0), \mathcal{J}_q(w_1), \dots, \mathcal{J}_q(w_\ell) \geq 0$, by (iv) in Lemma 4.5 we have that

$$c_q = \mathcal{J}_q(u_0) + \sum_{k=1}^{\ell} \mathcal{J}_q(w_k) > 0$$

and we conclude.

From now on we will denote a generic solution by u .

5. EXISTENCE FOR ALL q 'S IN THE CASE $p \in (3, 6)$

In this section we prove the existence of solutions of (1.1) for every $q \neq 0$, but only for p large, and, as we said in the Introduction, for radial symmetric functions. Observe that if u is radial, also ϕ_u is.

First we give some convergence properties in the radial setting recalling that $H_r^1(\mathbb{R}^3)$ is compactly embedded in $L^s(\mathbb{R}^3)$ for $s \in (2, 6)$ by the celebrated Strauss Lemma [36].

Lemma 5.1. *If $u_n \rightharpoonup u$ in $H_r^1(\mathbb{R}^3)$, then*

- (i) $\phi_{u_n} \rightarrow \phi_u$ in \mathcal{D} ;
- (ii) $\int \phi_{u_n} u_n^2 \rightarrow \int \phi_u u^2$.

Proof. To prove (i) we define the linear and continuous operators on \mathcal{D}

$$T_n(\varphi) = \int \varphi u_n^2 \quad \text{and} \quad T(\varphi) = \int \varphi u^2$$

represented, by the Riesz Theorem, respectively by ϕ_{u_n} and ϕ_u . Then, by the Hölder inequality,

$$\|\phi_{u_n} - \phi_u\|_{\mathcal{D}} = \|T_n - T\|_{\mathcal{D}'} \leq C \|u_n^2 - u^2\|_{6/5} \rightarrow 0.$$

Moreover from (i) and the Hölder inequality we easily get (ii). \square

5.1. The case $p \in [4, 6)$. In this case any (PS) sequence for \mathcal{J}_q is bounded. In fact, if $\{u_n\} \subset H^1(\mathbb{R}^3)$ is a (PS) sequence, that is

$$|\mathcal{J}_q(u_n)| \leq M, \quad \mathcal{J}'_q(u_n) = o_n(1),$$

then,

$$pM + c\|u_n\| \geq p\mathcal{J}_q(u_n) - \mathcal{J}'_q(u_n)[u_n] = \frac{p-2}{2}\|u\|^2 + q^2 \frac{p-4}{4} \int \phi_u u^2 \geq C\|u_n\|^2,$$

from which the boundedness of $\{u_n\}$ follows.

The next Lemma is standard, since we have compactness.

Lemma 5.2. *Let $p \in (2, 6)$. Any bounded sequence $\{u_n\} \subset H_r^1(\mathbb{R}^3)$ such that $\mathcal{J}'_q(u_n) \rightarrow 0$ has a convergent subsequence.*

Proof. We can assume, up to subsequence, that $u_n \rightharpoonup u$ in $H_r^1(\mathbb{R}^3)$ and $u_n \rightarrow u$ in $L^p(\mathbb{R}^3)$ for $p \in (2, 6)$. By Lemma 5.1 we have also that $\phi_{u_n} \rightarrow \phi_u$ in \mathcal{D} . By defining the Riesz isomorphism $R = -\Delta + \omega I : H_r^1(\mathbb{R}^3) \rightarrow H_r^{-1}(\mathbb{R}^3)$, by

$$R(u_n) + q^2 \phi_{u_n} u_n = |u_n|^{p-2} u_n + o_n(1)$$

we have

$$u_n = -q^2 R^{-1}(\phi_{u_n} u_n) + R^{-1}(|u_n|^{p-2} u_n) + o_n(1).$$

It is standard to see that each term in the right hand side is convergent in $H_r^1(\mathbb{R}^3)$, however, for the reader's convenience, we give a short proof. Observe that

$$\|\phi_{u_n} u_n\|_{3/2} \leq \|u_n\|_2 \|\phi_{u_n}\|_6 \leq \|u_n\| \|\phi_{u_n}\|_{\mathcal{D}} \leq C$$

and, since by duality, $L_r^{3/2}(\mathbb{R}^3) \hookrightarrow H_r^{-1}(\mathbb{R}^3)$ we deduce $\{\phi_{u_n} u_n\}$ is convergent in $H_r^{-1}(\mathbb{R}^3)$, and consequently $R^{-1}(\phi_{u_n} u_n)$ also is. Analogously,

$$\||u_n|^{p-2} u_n\|_{p'} = \|u_n\|_p^{p-1} \leq C \|u_n\|^{p-1} \leq C$$

and again the compact embedding of $L_r^{p'}(\mathbb{R}^3)$ into $H_r^{-1}(\mathbb{R}^3)$ guarantees that $\{|u_n|^{p-2} u_n\}$ is convergent into $H_r^{-1}(\mathbb{R}^3)$ and so we conclude. \square

Putting together Lemma 3.5, Lemma 5.2 and the boundedness of the (PS) sequences, the Mountain Pass Theorem allows to get a solution of (1.1).

5.2. The case $p \in (3, 4)$. To study this case, we apply the following result

Theorem 5.3 ([29, Theorem 1.1]). *Let X be a Banach space with a norm $\|\cdot\|$ and let $L \subset \mathbb{R}^+$ be an interval. We consider a family $\{I_\lambda\}$ of C^1 functionals on X of the form*

$$I_\lambda(u) = A(u) - \lambda B(u), \quad \forall \lambda \in L,$$

where $B(u) \geq 0$ for all $u \in X$ and either $A(u) \rightarrow +\infty$ or $B(u) \rightarrow +\infty$ as $\|u\| \rightarrow +\infty$. We assume that there exist two functions $v_1, v_2 \in X$ such that

$$c_\lambda = \inf_{\gamma \in \Gamma} \max_{t \in [0,1]} I_\lambda(\gamma(t)) > \max\{I_\lambda(v_1), I_\lambda(v_2)\}, \quad \forall \lambda \in L,$$

$\Gamma = \{\gamma \in C([0,1], X) : \gamma(0) = v_1, \gamma(1) = v_2\}$. Then for almost all $\lambda \in L$, there exists a bounded (PS) sequence $\{u_n(\lambda)\} \subset X$ of I_λ at level c_λ .

Let then, for $\lambda \in [1/2, 1]$,

$$\mathcal{J}_{q,\lambda}(u) := \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 + \frac{q^2}{4} \int \phi_u u^2 - \frac{\lambda}{p} \|u\|_p^p.$$

The Mountain Pass Geometry for $\mathcal{J}_{q,\lambda}$, which can be proved arguing as in Lemma 3.5, ensures that

$$c_\lambda := \inf_{\gamma \in \Gamma_{q,\lambda}} \max_{t \in [0,1]} \mathcal{J}_{q,\lambda}(\gamma(t)) > 0,$$

where $\Gamma_{q,\lambda} = \{\gamma \in C([0,1]; H_r^1(\mathbb{R}^3)) : \gamma(0) = 0, \gamma(1) = w\}$ and $w \in H_r^1(\mathbb{R}^3)$ is such that $\mathcal{J}_{q,\lambda}(w) < 0$. Then Theorem 5.3 gives, for a sequence $\{\lambda_j\} \subset [1/2, 1]$ such that $\lim_j \lambda_j = 1$, a bounded (PS) sequence $\{u_{n,\lambda_j}\}$ at level c_{λ_j} for the functional $\mathcal{J}_{q,\lambda_j}$.

Observe that, for all $j \in \mathbb{N}$, $c_{\lambda_j} \in [c_1, c_{1/2}]$.

In view of Lemma 5.2 we can assume that, up to subsequence, for every $j \in \mathbb{N}$, $\{u_{n,\lambda_j}\}$ strongly converges to some $u_{\lambda_j} \in H_r^1(\mathbb{R}^3)$ satisfying

$$\mathcal{J}_{q,\lambda_j}(u_{\lambda_j}) = c_{\lambda_j}, \quad \mathcal{J}'_{q,\lambda_j}(u_{\lambda_j}) = 0.$$

In particular, being such u_{λ_j} a solution of the equation

$$-\Delta u + \omega u + q^2 \phi_u u = \lambda_j |u|^{p-2} u \quad \text{in } \mathbb{R}^3,$$

and, arguing as in Appendix A.3, it satisfies the Pohozaev identity

$$\frac{1}{2} \|\nabla u_{\lambda_j}\|_2^2 + \frac{3}{2} \omega \|u_{\lambda_j}\|_2^2 - \frac{q^2}{16\pi} \|\nabla \phi_j\|_2^2 + \frac{q^2 a^2}{16\pi} \|\Delta \phi_j\|_2^2 + \frac{3}{2} q^2 \int \phi_j u_{\lambda_j}^2 - \frac{3\lambda_j}{p} \|u_{\lambda_j}\|_p^p = 0$$

with $\phi_j := \phi_{u_{\lambda_j}}$, which can be written also as

$$(5.1) \quad -\|\nabla u_{\lambda_j}\|_2^2 - \frac{q^2}{16\pi} \|\nabla \phi_j\|_2^2 + \frac{q^2 a^2}{16\pi} \|\Delta \phi_j\|_2^2 + \frac{3}{4} q^2 \int \phi_j u_{\lambda_j}^2 + 3c_{\lambda_j} = 0.$$

Moreover

$$(5.2) \quad \begin{aligned} pc_{\lambda_j} &= p\mathcal{J}_{q,\lambda_j}(u_{\lambda_j}) - \mathcal{J}'_{q,\lambda_j}(u_{\lambda_j})[u_{\lambda_j}] \\ &= \left(\frac{p}{2} - 1\right) \|\nabla u_{\lambda_j}\|_2^2 + \left(\frac{p}{2} - 1\right) \omega \|u_{\lambda_j}\|_2^2 + q^2 \left(\frac{p}{4} - 1\right) \int \phi_j u_{\lambda_j}^2. \end{aligned}$$

Thus, using (5.1) and (5.2), we get

$$(5.3) \quad \begin{aligned} (p-3) \|\nabla u_{\lambda_j}\|_2^2 + \frac{p-2}{2} \omega \|u_{\lambda_j}\|_2^2 &= \left(\frac{p}{2} - 2\right) \|\nabla u_{\lambda_j}\|_2^2 + \left(\frac{p}{2} - 1\right) \|\nabla u_{\lambda_j}\|_2^2 + \left(\frac{p}{2} - 1\right) \omega \|u_{\lambda_j}\|_2^2 \\ &= \left(\frac{5}{2}p - 6\right) c_{\lambda_j} + \frac{q^2}{8\pi} \left(\frac{p}{4} - 1\right) \left(-\|\nabla \phi_j\|_2^2 + a^2 \|\Delta \phi_j\|_2^2 + 4\pi \int \phi_j u_{\lambda_j}^2\right) \\ &= \left(\frac{5}{2}p - 6\right) c_{\lambda_j} + \frac{q^2 a^2}{4\pi} \left(\frac{p}{4} - 1\right) \|\Delta \phi_j\|_2^2 \\ &\leq \frac{5p-12}{2} c_{1/2} \end{aligned}$$

which gives directly the boundedness of $\{u_{\lambda_j}\}$.

We show now that $\{u_{\lambda_j}\}$ is indeed a (PS) sequence for the unperturbed functional \mathcal{J}_q . Indeed due to the boundedness of $\{u_{\lambda_j}\}$ in $H_r^1(\mathbb{R}^3)$:

$$\mathcal{J}_q(u_{\lambda_j}) = \mathcal{J}_{q,\lambda_j}(u_{\lambda_j}) - (1 - \lambda_j) \|u_{\lambda_j}\|_p^p = c_{\lambda_j} + o_j(1)$$

implying the boundedness of $\{\mathcal{J}_q(u_{\lambda_j})\}$. Moreover

$$\begin{aligned} \sup_{\|v\| \leq 1} |\mathcal{J}'_q(u_{\lambda_j})[v]| &= \sup_{\|v\| \leq 1} \left| \mathcal{J}'_{q,\lambda_j}(u_{\lambda_j})[v] - (1 - \lambda_j)p \int |u_{\lambda_j}|^{p-2} u_{\lambda_j} v \right| \\ &\leq \|\mathcal{J}'_{q,\lambda_j}(u_{\lambda_j})\| + (1 - \lambda_j)pC \|u_{\lambda_j}\|_p^{p-1} \\ &= o_j(1). \end{aligned}$$

But then, in view of Lemma 5.2, up to subsequence $u_{\lambda_j} \rightarrow \mathbf{u}$ and so $\mathcal{J}'_q(\mathbf{u}) = 0$, meaning that \mathbf{u} is a solution of (1.1) we were looking for.

We conclude the section with the following remark that will be useful in the next section.

Remark 5.4. *In the radial setting we can repeat the arguments in Section 4 replacing the Splitting Lemma 4.5 by standard arguments using the compact embedding of $H_r^1(\mathbb{R}^3)$ into $L^p(\mathbb{R}^3)$, $p \in (2, 6)$, Lemma B.1, and Lemma 5.1. In such a way, using the notations introduced in Lemma 4.3, for every $q < q_*$ we get a solution \mathbf{u} such that $\|\mathbf{u}\| \leq \bar{T}$.*

6. THE BEHAVIOUR AS $a \rightarrow 0$ IN THE RADIAL CASE

This section is devoted to the proof of Theorem 1.3. We begin by showing the following preliminary result.

Lemma 6.1. *Consider $f^0 \in L^{6/5}(\mathbb{R}^3)$, $\{f_a\}_{a \in (0,1)} \subset L^{6/5}(\mathbb{R}^3)$ and let*

$$\phi^0 \in D^{1,2}(\mathbb{R}^3) \text{ be the unique solution of } -\Delta\phi = f^0 \text{ in } \mathbb{R}^3$$

and

$$\phi^a \in \mathcal{D} \text{ be the unique solution of } -\Delta\phi + a^2\Delta^2\phi = f^a \text{ in } \mathbb{R}^3.$$

As $a \rightarrow 0$ we have:

- (i) if $f^a \rightharpoonup f^0$ in $L^{6/5}(\mathbb{R}^3)$, then $\phi^a \rightharpoonup \phi^0$ in $D^{1,2}(\mathbb{R}^3)$;
- (ii) if $f^a \rightarrow f^0$ in $L^{6/5}(\mathbb{R}^3)$, then $\phi^a \rightarrow \phi^0$ in $D^{1,2}(\mathbb{R}^3)$ and $a\Delta\phi^a \rightarrow 0$ in $L^2(\mathbb{R}^3)$.

Proof. By

$$\|\nabla\phi^a\|_2^2 + a^2\|\Delta\phi^a\|_2^2 = \int f^a\phi^a \leq C\|f^a\|_{6/5}\|\nabla\phi^a\|_2$$

we deduce

$$\|\nabla\phi^a\|_2 \leq C\|f^a\|_{6/5}, \quad \|a\Delta\phi^a\|_2 \leq C\|f^a\|_{6/5}.$$

Then there exists $\phi_* \in D^{1,2}(\mathbb{R}^3)$ such that $\phi^a \rightharpoonup \phi_*$ in $D^{1,2}(\mathbb{R}^3)$. Passing to the limit as $a \rightarrow 0$ in the identity

$$\int \nabla\phi^a \nabla\varphi + a^2 \int \Delta\phi^a \Delta\varphi = \int f^a\varphi, \quad \varphi \in C_c^\infty(\mathbb{R}^3),$$

and using that

$$\left| a^2 \int \Delta\phi^a \Delta\varphi \right| \leq a\|a\Delta\phi^a\|_2\|\Delta\varphi\|_2 \leq aC \rightarrow 0,$$

we get

$$\int \nabla\phi_* \nabla\varphi = \int f^0\varphi.$$

Then, by the uniqueness, $\phi_* = \phi^0$ proving (i).

Assume now that $f^a \rightarrow f^0$ in $L^{6/5}(\mathbb{R}^3)$. Of course we have

$$(6.1) \quad \|\nabla\phi^0\|_2^2 \leq \liminf_{a \rightarrow 0} \|\nabla\phi^a\|_2^2.$$

Let $\{\varphi_n\} \subset C_c^\infty(\mathbb{R}^3)$ such that $\varphi_n \rightarrow \phi^0$ in $D^{1,2}(\mathbb{R}^3)$ as $n \rightarrow +\infty$. Since ϕ^a minimizes the functional

$$E_a(\phi) = \frac{1}{2}\|\nabla\phi\|_2^2 + \frac{a^2}{2}\|\Delta\phi\|_2^2 - \int f^a\phi, \quad \phi \in \mathcal{D},$$

we have

$$\begin{aligned} \frac{1}{2}\|\nabla\phi^a\|_2^2 &= E_a(\phi^a) - \frac{a^2}{2}\|\Delta\phi^a\|_2^2 + \int f^a\phi^a \\ &\leq E_a(\varphi_n) + \int f^a\phi^a \\ &= \frac{1}{2}\|\nabla\varphi_n\|_2^2 + \frac{a^2}{2}\|\Delta\varphi_n\|_2^2 - \int f^a\varphi_n + \int f^a\phi^a \end{aligned}$$

and then

$$(6.2) \quad \limsup_{a \rightarrow 0} \frac{1}{2}\|\nabla\phi^a\|_2^2 \leq \frac{1}{2}\|\nabla\varphi_n\|_2^2 - \int f^0\varphi_n + \int f^0\phi^0.$$

Passing to the limit in n in (6.2) we get

$$(6.3) \quad \limsup_{a \rightarrow 0} \|\nabla \phi^a\|_2^2 \leq \|\nabla \phi^0\|_2^2.$$

By (6.1), (6.3) and the convergence $\phi^a \rightharpoonup \phi^0$ in $D^{1,2}(\mathbb{R}^3)$ we infer $\phi^a \rightarrow \phi^0$ in $D^{1,2}(\mathbb{R}^3)$. Finally we see that, for $a \rightarrow 0$,

$$\|a\Delta \phi^a\|_2^2 = \int f^a \phi^a - \|\nabla \phi^a\|_2^2 \rightarrow \int f^0 \phi^0 - \|\nabla \phi^0\|_2^2 = 0$$

and the proof is complete. \square

Now we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. From now on we fix q according to the restriction of Theorem 1.1, and let $\{\mathbf{u}^a, \phi^a\} \subset H_r^1(\mathbb{R}^3) \times \mathcal{D}_r$ be the family of the solutions of (1.1) for this fixed value q . We are using the notation $\phi^a := \phi_{\mathbf{u}^a}^a$. In contrast to the previous sections we use the explicit dependence on a also in the functional. Then, the functions $\{\mathbf{u}^a\}$ solve

$$-\Delta u + \omega u + q^2 \phi^a u = |u|^{p-2} u \text{ in } \mathbb{R}^3$$

and are critical point of \mathcal{J}_q^a at the Mountain Pass value $c_q^a > 0$.

Our first aim is to show the boundedness of $\{\mathbf{u}^a\}_{a \in (0,1)}$ in $H_r^1(\mathbb{R}^3)$.

Let

$$c_q^0 := \inf_{\gamma \in \Gamma^0} \max_{t \in [0,1]} \mathcal{J}^0(\gamma(t)) \geq c_q^a > 0,$$

where $\Gamma^0 := \{\gamma \in C([0,1], H_r^1(\mathbb{R}^3)) : \gamma(0) = 0, \mathcal{J}^0(\gamma(1)) < 0\}$,

$$\mathcal{J}^0(u) := \frac{1}{2} \|\nabla u\|_2^2 + \frac{\omega}{2} \|u\|_2^2 + \frac{q^2}{4} \iint \frac{u^2(x)u^2(y)}{|x-y|} - \frac{1}{p} \|u\|_p^p$$

is the functional related to (1.3).

We distinguish three cases.

Case A: $p \in [4, 6)$.

We have

$$c_q^0 \geq c_q^a = \mathcal{J}_q^a(\mathbf{u}^a) - \frac{1}{p} (\mathcal{J}_q^a)'(\mathbf{u}^a)[\mathbf{u}^a] = \frac{p-2}{2p} \|\mathbf{u}^a\|^2 + \frac{p-4}{4p} \int \phi^a (\mathbf{u}^a)^2 \geq \frac{p-2}{2p} \|\mathbf{u}^a\|^2.$$

Case B: $p \in (3, 4)$.

Arguing as in (5.3), since \mathbf{u}^a is a solution at the Mountain Pass level c_q^a , we infer

$$(p-3) \|\nabla \mathbf{u}^a\|_2^2 + \frac{p-2}{2} \omega \|\mathbf{u}^a\|_2^2 \leq \frac{5p-12}{2} c_q^a \leq \frac{5p-12}{2} c_q^0.$$

Case C: $p \in (2, 3]$.

By Remark 5.4 we already know that $\|\mathbf{u}^a\| \leq \bar{T}$, with \bar{T} that does not depend on a (see Remark 4.4).

Hence in any case $\{\mathbf{u}^a\}$ is bounded in $H_r^1(\mathbb{R}^3)$ and there exists $\mathbf{u}^0 \in H_r^1(\mathbb{R}^3)$ such that, up to subsequences, $\mathbf{u}^a \rightharpoonup \mathbf{u}^0$ in $H_r^1(\mathbb{R}^3)$ as $a \rightarrow 0$. In particular $(\mathbf{u}^a)^2 \rightarrow (\mathbf{u}^0)^2$ in $L^{6/5}(\mathbb{R}^3)$ and by (ii) of Lemma 6.1 we infer that $\phi^a \rightarrow \phi^0$, where $\phi^0 \in D_r^{1,2}(\mathbb{R}^3)$ is the unique solution of $-\Delta \phi = 4\pi \mathbf{u}_0^2$ in \mathbb{R}^3 . The fact that $\mathbf{u}^a \rightarrow \mathbf{u}^0$ in $H_r^1(\mathbb{R}^3)$ is done as in Lemma 5.2 since the proof can be merely repeated using Lemma 6.1.

Let now $\varphi \in C_c^\infty(\mathbb{R}^3)$ with $\text{supp}(\varphi) = \Omega$. We know that

$$\langle \mathbf{u}^a, \varphi \rangle + q^2 \int_\Omega \phi^a \mathbf{u}^a \varphi = \int_\Omega |\mathbf{u}^a|^{p-2} \mathbf{u}^a \varphi.$$

We want to pass to the limit as $a \rightarrow 0$ in each term. Of course

$$(6.4) \quad \langle \mathbf{u}^a, \varphi \rangle \rightarrow \langle \mathbf{u}^0, \varphi \rangle,$$

and, as follows by standard arguments,

$$\int_{\Omega} |\mathbf{u}^a|^{p-2} \mathbf{u}^a \varphi \rightarrow \int_{\Omega} |\mathbf{u}^0|^{p-2} \mathbf{u}^0 \varphi.$$

Moreover, since $\phi^a \rightarrow \phi^0$ in $L^6(\mathbb{R}^3)$, $\mathbf{u}^a \rightarrow \mathbf{u}^0$ in $L^{12/5}(\Omega)$ and $\varphi \in L^{12/5}(\Omega)$, by the Hölder inequality we easily get

$$(6.5) \quad \int_{\Omega} \phi^a \mathbf{u}^a \varphi \rightarrow \int_{\Omega} \phi^0 \mathbf{u}^0 \varphi.$$

Then by (6.4)-(6.5) we arrive at

$$\langle \mathbf{u}^0, \varphi \rangle + q^2 \int_{\Omega} \phi^0 \mathbf{u}^0 \varphi = \int_{\Omega} |\mathbf{u}^0|^{p-2} \mathbf{u}^0 \varphi$$

which shows that (\mathbf{u}^0, ϕ^0) solves (1.3). \square

APPENDIX A. PROPERTIES OF SOLUTIONS AND NONEXISTENCE

In this appendix we show that our solutions are indeed classical. Moreover we prove, by means of Nehari and Pohožaev type identities, some nonexistence results.

A.1. Regularity of the solutions. We remark here that the weak solutions we find are indeed classical solutions. This is based on standard bootstrap arguments that we briefly recall here.

Let us first observe that if $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ is a weak solution of (1.1) then $\psi := -a^2 \Delta \phi + \phi$ solves weakly, in any bounded domain Ω , the equation

$$-\Delta \psi = 4\pi u^2 \quad \text{in } \Omega.$$

Now, being $u^2 \in L^3(\mathbb{R}^3)$ it holds (see e.g. [28, Theorem 9.9])

$$(A.1) \quad -a^2 \Delta \phi + \phi = \psi \in W_{\text{loc}}^{2,3}(\mathbb{R}^3).$$

Since $\phi \in H_{\text{loc}}^1(\mathbb{R}^3)$ is a weak solution of (A.1) with $\psi \in W_{\text{loc}}^{2,2}(\mathbb{R}^3)$, by *higher interior regularity* (see e.g. [28, Theorem 8.10]), we deduce $\phi \in W_{\text{loc}}^{4,2}(\mathbb{R}^3)$ and by the Sobolev embedding (see e.g. [1, Theorem 5.4]) we deduce that $\phi \in C_{\text{loc}}^{2,\lambda}(\mathbb{R}^3)$, $\lambda \in (0, 1/2]$.

Then considering the equation

$$-\Delta u + \omega u + q^2 \phi u = |u|^{p-2} u$$

we deduce by bootstrap arguments that $u \in C_{\text{loc}}^{2,\lambda}(\mathbb{R}^3)$. But then, being

$$-\Delta \psi = 4\pi u^2 \in H_{\text{loc}}^2(\mathbb{R}^3),$$

it holds again by [28, Theorem 8.10] that

$$-a^2 \Delta \phi + \phi = \psi \in H_{\text{loc}}^4(\mathbb{R}^3)$$

and then, by higher interior regularity and Sobolev embedding, $\phi \in H_{\text{loc}}^6(\mathbb{R}^3) \hookrightarrow C_{\text{loc}}^{4,\lambda}(\mathbb{R}^3)$, $\lambda \in (0, 1/2]$.

A.2. A useful identity. We define the Fourier transform of a function $f \in L^1(\mathbb{R}^3)$ by the formula

$$\mathcal{F}(f)(x) := \frac{1}{(2\pi)^{3/2}} \int e^{-ixy} f(y) dy.$$

If $f \in L^2(\mathbb{R}^3)$ its Fourier transform is defined by the usual approximation procedure. With this definition we have, for $f, g \in L^2(\mathbb{R}^3)$,

$$\mathcal{F}(f * g) = (2\pi)^{3/2} \mathcal{F}(f) \mathcal{F}(g) \quad \text{and} \quad \int \mathcal{F}(f) \mathcal{F}(g) = \int fg.$$

Then, by classical results in Fourier analysis (observe that the functions involved are e.g. in $L^1(\mathbb{R}^3) \cap L^2(\mathbb{R}^3)$), we get for any $a > 0$,

$$\mathcal{F}(e^{-|\cdot|/a})(x) = \sqrt{\frac{2}{\pi}} \frac{2a^3}{(1 + a^2|x|^2)^2} \quad \text{and} \quad \mathcal{F}\left(\frac{e^{-|\cdot|/a}}{|\cdot|}\right)(x) = \sqrt{\frac{2}{\pi}} \frac{a^2}{1 + a^2|x|^2}.$$

Then by recalling (3.5) we have

$$(A.2) \quad \mathcal{F}(\Delta\mathcal{K} * \Delta\mathcal{K}) = (2\pi)^{3/2} \mathcal{F}(\Delta\mathcal{K}) \mathcal{F}(\Delta\mathcal{K}) = \frac{4\sqrt{2\pi}}{(1 + a^2|\cdot|^2)^2}$$

from which we deduce

$$\Delta\mathcal{K} * \Delta\mathcal{K} = \frac{2\pi}{a^3} e^{-|\cdot|/a}.$$

Since for $u \in H^1(\mathbb{R}^3)$ it is $\Delta\phi_u = \Delta\mathcal{K} * u^2$ (see Lemma 3.3 and item (v) in Lemma 3.4), in virtue of (A.2), we get

$$\begin{aligned} \|\Delta\phi_u\|_2^2 &= \int |\mathcal{F}(\Delta\phi_u)|^2 \\ &= (2\pi)^{3/2} \int \mathcal{F}(u^2) \mathcal{F}(\Delta\mathcal{K}) \mathcal{F}(\Delta\phi_u) \\ (A.3) \quad &= \int \mathcal{F}(u^2) \mathcal{F}(\Delta\mathcal{K} * \Delta\phi_u) \\ &= \int u^2 \Delta\mathcal{K} * \Delta\mathcal{K} * u^2 \\ &= \frac{2\pi}{a^3} \iint e^{-|x-y|/a} u^2(x) u^2(y) dx dy. \end{aligned}$$

In another words we have the identity

$$\int \left(\int \frac{e^{-|x-y|/a}}{|x-y|} u^2(y) dy \right)^2 dx = 2\pi a \iint e^{-|x-y|/a} u^2(x) u^2(y) dx dy$$

true for any $u \in H^1(\mathbb{R}^3)$.

A.3. The Pohožaev identity. Let $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ be a nontrivial solution of (1.1). Recall that $\phi = \phi_u$. We have

$$(A.4) \quad \|\nabla u\|_2^2 + \omega \|u\|_2^2 + q^2 \int \phi u^2 - \|u\|_p^p = 0$$

and

$$(A.5) \quad \|\nabla \phi\|_2^2 + a^2 \|\Delta\phi\|_2^2 = 4\pi \int \phi u^2,$$

that are usually called *Nehari* identities.

Moreover (u, ϕ) satisfies also the following *Pohožaev* identity

$$(A.6) \quad -\frac{1}{2}\|\nabla u\|_2^2 - \frac{3}{2}\omega\|u\|_2^2 + \frac{q^2}{16\pi}\|\nabla\phi\|_2^2 - \frac{q^2a^2}{16\pi}\|\Delta\phi\|_2^2 - \frac{3}{2}q^2\int\phi u^2 + \frac{3}{p}\|u\|_p^p = 0.$$

In fact, if (u, ϕ) solve (1.1), recalling the regularity proved in Section A.1, for every $R > 0$, we have

$$(A.7) \quad \int_{B_R} -\Delta u(x \cdot \nabla u) = -\frac{1}{2}\int_{B_R} |\nabla u|^2 - \frac{1}{R}\int_{\partial B_R} |x \cdot \nabla u|^2 + \frac{R}{2}\int_{\partial B_R} |\nabla u|^2,$$

$$(A.8) \quad \int_{B_R} \phi u(x \cdot \nabla u) = -\frac{1}{2}\int_{B_R} u^2(x \cdot \nabla \phi) - \frac{3}{2}\int_{B_R} \phi u^2 + \frac{R}{2}\int_{\partial B_R} \phi u^2,$$

$$(A.9) \quad \int_{B_R} u(x \cdot \nabla u) = -\frac{3}{2}\int_{B_R} u^2 + \frac{R}{2}\int_{\partial B_R} u^2,$$

$$(A.10) \quad \int_{B_R} |u|^{p-2}u(x \cdot \nabla u) = -\frac{3}{p}\int_{B_R} |u|^p + \frac{R}{p}\int_{\partial B_R} |u|^p,$$

where B_R is the ball of \mathbb{R}^3 centered in the origin and with radius R (see also [22]), and, since

$$\Delta^2\phi(x \cdot \nabla\phi) = \operatorname{div}\left(\nabla\Delta\phi(x \cdot \nabla\phi) - \Delta\phi\nabla\phi - \mathbb{F} + x\frac{(\Delta\phi)^2}{2}\right) + \frac{(\Delta\phi)^2}{2},$$

where $\mathbb{F}_i = \Delta\phi(x \cdot \nabla(\partial_i\phi))$, $i = 1, 2, 3$, then

$$(A.11) \quad \int_{B_R} \Delta^2\phi(x \cdot \nabla\phi) = \frac{1}{2}\int_{B_R} (\Delta\phi)^2 + \int_{\partial B_R} \left(\nabla\Delta\phi(x \cdot \nabla\phi) - \Delta\phi\nabla\phi - \mathbb{F} + x\frac{(\Delta\phi)^2}{2}\right) \cdot \nu.$$

Multiplying the first equation of (1.1) by $x \cdot \nabla u$ and the second equation by $x \cdot \nabla\phi$ and integrating on B_R , by (A.7), (A.8), (A.9), (A.10), and (A.11) we get

$$(A.12) \quad \begin{aligned} & -\frac{1}{2}\int_{B_R} |\nabla u|^2 - \frac{1}{R}\int_{\partial B_R} |x \cdot \nabla u|^2 + \frac{R}{2}\int_{\partial B_R} |\nabla u|^2 - \frac{3}{2}\omega\int_{B_R} u^2 + \frac{R}{2}\omega\int_{\partial B_R} u^2 \\ & - \frac{q^2}{2}\int_{B_R} u^2(x \cdot \nabla\phi) - \frac{3}{2}q^2\int_{B_R} \phi u^2 + q^2\frac{R}{2}\int_{\partial B_R} \phi u^2 = -\frac{3}{p}\int_{B_R} |u|^p + \frac{R}{p}\int_{\partial B_R} |u|^p \end{aligned}$$

and

$$(A.13) \quad \begin{aligned} 4\pi\int_{B_R} u^2(x \cdot \nabla\phi) &= -\frac{1}{2}\int_{B_R} |\nabla\phi|^2 - \frac{1}{R}\int_{\partial B_R} |x \cdot \nabla\phi|^2 + \frac{R}{2}\int_{\partial B_R} |\nabla\phi|^2 + \frac{a^2}{2}\int_{B_R} (\Delta\phi)^2 \\ &+ a^2\int_{\partial B_R} \left(\nabla\Delta\phi(x \cdot \nabla\phi) - \Delta\phi\nabla\phi - \mathbb{F} + x\frac{(\Delta\phi)^2}{2}\right) \cdot \nu. \end{aligned}$$

Substituting (A.13) into (A.12) we obtain

$$\begin{aligned} & -\frac{1}{2}\int_{B_R} |\nabla u|^2 - \frac{3}{2}\omega\int_{B_R} u^2 + \frac{q^2}{16\pi}\int_{B_R} |\nabla\phi|^2 - \frac{q^2a^2}{16\pi}\int_{B_R} (\Delta\phi)^2 - \frac{3}{2}q^2\int_{B_R} \phi u^2 + \frac{3}{p}\int_{B_R} |u|^p \\ &= \frac{1}{R}\int_{\partial B_R} |x \cdot \nabla u|^2 - \frac{R}{2}\int_{\partial B_R} |\nabla u|^2 - \frac{R}{2}\omega\int_{\partial B_R} u^2 - q^2\frac{R}{2}\int_{\partial B_R} \phi u^2 + \frac{R}{p}\int_{\partial B_R} |u|^p \\ & - \frac{q^2}{8\pi R}\int_{\partial B_R} |x \cdot \nabla\phi|^2 + \frac{q^2R}{16\pi}\int_{\partial B_R} |\nabla\phi|^2 \\ & + \frac{q^2a^2}{8\pi}\int_{\partial B_R} \left(\nabla\Delta\phi(x \cdot \nabla\phi) - \Delta\phi\nabla\phi - \mathbb{F} + x\frac{(\Delta\phi)^2}{2}\right) \cdot \nu. \end{aligned}$$

Using the same arguments as in [22, Proof of Theorem 1.1] we have that the right hand side tends to zero as $R \rightarrow +\infty$, since

$$\begin{aligned} \int_{\partial B_R} \nabla \Delta \phi (x \cdot \nabla \phi) \cdot \nu &= R \int_{\partial B_R} \frac{\partial \Delta \phi}{\partial \nu} \frac{\partial \phi}{\partial \nu} \rightarrow 0, \\ \int_{\partial B_R} \Delta \phi \nabla \phi \cdot \nu &= \int_{\partial B_R} \Delta \phi \frac{\partial \phi}{\partial \nu} \rightarrow 0, \\ \int_{\partial B_R} \mathbb{F} \cdot \nu &= R \int_{\partial B_R} \frac{\partial^2 \phi}{\partial \nu^2} \rightarrow 0, \\ \frac{1}{2} \int_{\partial B_R} (\Delta \phi)^2 x \cdot \nu &= \frac{R}{2} \int_{\partial B_R} (\Delta \phi)^2 \rightarrow 0, \end{aligned}$$

and so we get (A.6).

Finally we observe that, using (A.3), the Pohožaev identity (A.6) can be written also as

$$(A.14) \quad -\frac{1}{2} \|\nabla u\|_2^2 - \frac{3}{2} \omega \|u\|_2^2 - \frac{q^2}{4a} \iint \left[5 \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|/a} + e^{-\frac{|x-y|}{a}} \right] u^2(x) u^2(y) dx dy + \frac{3}{p} \|u\|_p^p = 0.$$

A.4. A nonexistence result. Using the identities recalled before, we are able to show nonexistence results for $p \leq 2$ and for $p \geq 6$.

In fact, if $(u, \phi) \in H^1(\mathbb{R}^3) \times \mathcal{D}$ is a nontrivial solution of (1.1) and $p \geq 6$, replacing (A.4) and (A.5) into (A.6) we get

$$\begin{aligned} 0 &= -\frac{1}{2} \|\nabla u\|_2^2 - \frac{3}{2} \omega \|u\|_2^2 + \frac{q^2}{16\pi} \|\nabla \phi\|_2^2 - \frac{q^2 a^2}{16\pi} \|\Delta \phi\|_2^2 - \frac{3}{2} q^2 \int \phi u^2 + \frac{3}{p} \|u\|_p^p \\ &= \left(\frac{3}{p} - \frac{1}{2} \right) \|\nabla u\|_2^2 + \left(\frac{3}{p} - \frac{3}{2} \right) \omega \|u\|_2^2 - \frac{q^2 a^2}{8\pi} \|\Delta \phi\|_2^2 + \left(\frac{3}{p} - \frac{5}{4} \right) q^2 \int \phi u^2 \\ &\leq -\omega \|u\|_2^2 < 0. \end{aligned}$$

Moreover, if $p \leq 2$, replacing (A.14) into (A.5) and using (3.8), we have

$$\begin{aligned} 0 &= \|\nabla u\|_2^2 + \omega \|u\|_2^2 + q^2 \int \phi u^2 - \|u\|_p^p \\ &= \left(1 - \frac{p}{6} \right) \|\nabla u\|_2^2 + \left(1 - \frac{p}{2} \right) \omega \|u\|_2^2 + \left(1 - \frac{5}{12p} \right) q^2 \iint \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|} u^2(x) u^2(y) dx dy \\ &\quad - q^2 \frac{p}{12a} \iint e^{-\frac{|x-y|}{a}} u^2(x) u^2(y) dx dy \\ &= \left(1 - \frac{p}{6} \right) \|\nabla u\|_2^2 + \left(1 - \frac{p}{2} \right) \omega \|u\|_2^2 \\ &\quad + \frac{q^2}{a} \iint \left[\left(1 - \frac{5}{12p} \right) \frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|/a} - \frac{p}{12} e^{-\frac{|x-y|}{a}} \right] u^2(x) u^2(y) dx dy \\ &\geq \frac{2}{3} \|\nabla u\|_2^2 + \frac{q^2}{6a} \iint \left[\frac{1 - e^{-\frac{|x-y|}{a}}}{|x-y|/a} - e^{-\frac{|x-y|}{a}} \right] u^2(x) u^2(y) dx dy > 0 \end{aligned}$$

since the function in the parenthesis is positive.

APPENDIX B. PROOF OF LEMMA 4.5

This appendix is devoted to the proof of the Splitting Lemma. To do this, we need some preliminary results.

Lemma B.1. *The weak limit of a (PS) sequence for \mathcal{J}_q in $H^1(\mathbb{R}^3)$ is a critical point of \mathcal{J}_q .*

Proof. Let $\{v_n\} \subset H^1(\mathbb{R}^3)$ be a (PS) sequence for \mathcal{J}_q and v its weak limit. Then, for all $\varphi \in C_0^\infty(\mathbb{R}^3)$ we have that

$$\mathcal{J}'_q(v_n)[\varphi] = \int \nabla v_n \nabla \varphi + \omega \int v_n \varphi + q^2 \int \phi_{v_n} v_n \varphi - \int |v_n|^{p-2} v_n \varphi \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Due to the strong convergence of v_n to v in $L_{\text{loc}}^q(\mathbb{R}^3)$ for $1 \leq q < 6$, to conclude it is enough to prove that

$$\int \phi_{v_n} v_n \varphi \rightarrow \int \phi_v v \varphi.$$

Observe that

$$\left| \int \phi_{v_n} v_n \varphi - \int \phi_v v \varphi \right| \leq \underbrace{\int \phi_{v_n} |v_n - v| |\varphi|}_{I_1} + \underbrace{\int |\phi_{v_n} - \phi_v| |v \varphi|}_{I_2}.$$

By the Hölder inequality, the boundedness of $\{\phi_{v_n}\}$ in $L^6(\mathbb{R}^3)$, see (vi) in Lemma 3.4, and the strong convergence of v_n to v in $L_{\text{loc}}^3(\mathbb{R}^3)$ we get

$$I_1 \leq \|\phi_{v_n}\|_6 \|v_n - v\|_{L^2(\text{supp } \varphi)} \|\varphi\|_{L^3(\text{supp } \varphi)} \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

On the other hand, using Lemma 3.4, item (viii) we infer

$$I_2 \leq \|\phi_{v_n} - \phi_v\|_{L^2(\text{supp } \varphi)} \|v \varphi\|_{L^2(\text{supp } \varphi)} \rightarrow 0 \quad \text{as } n \rightarrow +\infty,$$

completing the proof by density. \square

Lemma B.2. *For every $v \in H^1(\mathbb{R}^3)$ and $v_n \rightharpoonup 0$ in $H^1(\mathbb{R}^3)$, we have*

$$\mathcal{J}_q(v_n + v) - \mathcal{J}_q(v_n) - \mathcal{J}_q(v) \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

Proof. By $\|v_n + v\|^2 = \|v_n\|^2 + \|v\|^2 + o_n(1)$ and the Brezis-Lieb Lemma it is enough to show that

$$\int \phi_{v_n+v} (v_n + v)^2 - \int \phi_{v_n} v_n^2 - \int \phi_v v^2 \rightarrow 0 \quad \text{as } n \rightarrow +\infty.$$

But

$$\int \phi_{v_n+v} (v_n + v)^2 - \int \phi_{v_n} v_n^2 - \int \phi_v v^2 = 4 \int \phi_{v_n} v_n v + 2 \int \phi_{v_n} v^2 + 4 \int (\mathcal{K} * v_n v) v_n v + 4 \int \phi_v v_n v,$$

and each term in the right hand side above converges to zero. Let us see the proof of the second one, being the proof of the other terms completely analogous.

For a subset $A \subset \mathbb{R}^3$ let us denote with $\mathbf{1}_A$ its characteristic function. Let B_1 and B_2 be two spheres centered in 0 with radius R_1 and R_2 . We first write

$$\int \phi_{v_n} v^2 = \int (\mathcal{K} * \mathbf{1}_{B_1} v_n^2) v^2 + \int (\mathcal{K} * \mathbf{1}_{B_1^c} v_n^2) \mathbf{1}_{B_2} v^2 + \int (\mathcal{K} * \mathbf{1}_{B_1^c} v_n^2) \mathbf{1}_{B_2^c} v^2.$$

Then, since $\mathcal{K} \leq 1/a$, we easily get

$$\begin{aligned} \int (\mathcal{K} * \mathbf{1}_{B_1} v_n^2) v^2 &\leq \frac{1}{a} \|v_n\|_{L^2(B_1)}^2 \|v\|_2^2 \rightarrow 0, \\ \int (\mathcal{K} * \mathbf{1}_{B_1^c} v_n^2) \mathbf{1}_{B_2^c} v^2 &\leq \frac{1}{a} \|v_n\|_2^2 \|v\|_{L^2(B_2^c)}^2 < \frac{1}{n}, \end{aligned}$$

if $R_2 = R_2(n)$ is taken sufficiently large, and, using that $\mathcal{K} \leq |x|^{-1}$,

$$\int (\mathcal{K} * \mathbf{1}_{B_1^c} v_n^2) \mathbf{1}_{B_2} v^2 \leq \iint_{B_2 \times B_1^c} \frac{v_n^2(y) v^2(x)}{|x - y|} dx dy \leq \frac{\|v\|_2^2 \|v_n\|_2^2}{|R_1 - R_2|} < \frac{1}{n}$$

taking $R_1 = R_1(n)$ sufficiently large. \square

Let us recall the Lions Lemma

Lemma B.3. *Let $2 < r < 6$. There exists a constant $C > 0$ such that*

$$\forall u \in H^1(\mathbb{R}^3) : \|u\|_r \leq \left(\sup_{z \in \mathbb{Z}^3} \|u\|_{L^2(z+Q)} \right)^{(r-2)/r} \|u\|^{2/r}$$

where $Q = [0, 1]^3$.

As a consequence of this lemma we infer

Lemma B.4. *Let $\{v_n\} \subset H^1(\mathbb{R}^3)$ be a sequence such that $v_n \rightharpoonup 0$ in $H^1(\mathbb{R}^3)$. Then $\mathcal{J}'_q(v_n) \rightarrow 0$. If, in addition, $v_n \not\rightarrow 0$ in $H^1(\mathbb{R}^3)$, then, up to subsequences,*

$$\exists \{z_n\} \subset \mathbb{Z}^3 \text{ with } |z_n| \rightarrow +\infty \text{ such that } \lim_n \|v_n\|_{L^p(z_n+Q)} > 0.$$

Proof. Let $\varphi \in C_c^\infty(\mathbb{R}^3)$ and $\Omega := \text{supp } \varphi$. We have

$$\mathcal{J}'_q(v_n)[\varphi] = \langle v_n, \varphi \rangle + \int_{\Omega} \phi_{v_n} v_n \varphi - \int_{\Omega} |v_n|^{p-2} v_n \varphi.$$

Then the first part follows by observing that

$$\begin{aligned} \left| \int_{\Omega} \phi_{v_n} v_n \varphi \right| &\leq \frac{1}{a} \|v_n\|_2^2 \|v_n\|_{L^2(\Omega)} \|\varphi\|_{L^2(\Omega)} = o_n(1), \\ \left| \int_{\Omega} |v_n|^{p-2} v_n \varphi \right| &\leq \|v_n\|_{L^{p'}(\Omega)}^{p-1} \|\varphi\|_{L^p(\Omega)} = o_n(1), \end{aligned}$$

uniformly in φ . We conclude by density.

Assume now that $v_n \not\rightarrow 0$. Then there exists a subsequence, that we rename again v_n , such that $\|v_n\| \rightarrow \alpha > 0$. If $\liminf_n \sup_{z \in \mathbb{Z}^3} \|v_n\|_{L^p(z+Q)} = 0$, the Lions Lemma B.3 gives $\liminf_n \|v_n\|_p = 0$ and then we have

$$0 < \alpha = \liminf_n \|v_n\|^2 \leq \lim_n \mathcal{J}'_q(v_n)[v_n] + \liminf_n \|v_n\|_p^p = 0,$$

reaching a contradiction. Hence $\liminf_n \sup_{z \in \mathbb{Z}^3} \|v_n\|_{L^p(z+Q)} > 0$ and then there exists a sequence $\{z_n\} \subset \mathbb{Z}^3$ such that

$$\lim_n \|v_n\|_{L^p(z_n+Q)} > 0.$$

The sequence $\{z_n\}$ has to be unbounded. Otherwise, if for some $R > 0$ it is $z_n + Q \subset B_R$ for all $n \in \mathbb{N}$, we have the contradiction

$$0 < \lim_n \|v_n\|_{L^p(z_n+Q)} \leq \lim_n \|v_n\|_{L^p(B_R)} = 0,$$

concluding the proof. \square

Finally we recall two basic facts.

Lemma B.5. *Let $\{y_n\} \subset \mathbb{R}^3$, $v \in H^1(\mathbb{R}^3)$, $\{v_n\} \subset H^1(\mathbb{R}^3)$ be bounded.*

- (i) *If $|y_n| \rightarrow +\infty$, then $v(\cdot + y_n) \rightharpoonup 0$ in $H^1(\mathbb{R}^3)$.*
- (ii) *If $\{y_n\}$ is bounded, then, up to a subsequence,*

$$v_n \not\rightarrow 0 \text{ in } H^1(\mathbb{R}^3) \implies v_n(\cdot + y_n) \not\rightarrow 0 \text{ in } H^1(\mathbb{R}^3).$$

Proof. For the first part, if $w \in H^1(\mathbb{R}^3)$ and $\varepsilon > 0$, then there exists $\varphi_\varepsilon \in C_c^\infty(\mathbb{R}^3)$ such that $\|w - \varphi_\varepsilon\| \leq \varepsilon$. Consequently

$$|\langle v(\cdot + y_n), w \rangle| \leq |\langle v(\cdot + y_n), w - \varphi_\varepsilon \rangle| + |\langle v(\cdot + y_n), \varphi_\varepsilon \rangle| \leq \varepsilon \|v\| + o_n(1)$$

proving that $\limsup_n |\langle v(\cdot + y_n), w \rangle| \leq \varepsilon \|v\|$.

To show the second part, let $\varphi \in C_c^\infty(\mathbb{R}^3)$ and $y \in \mathbb{R}^3$ be such that $\langle v_n, \varphi \rangle \rightarrow \eta \neq 0$ and $y_n \rightarrow y$. We have

$$\langle v_n(\cdot + y_n), \varphi(\cdot + y) \rangle = \langle v_n, \varphi(\cdot + y - y_n) \rangle = \langle v_n, \varphi \rangle + \langle v_n, \varphi(\cdot + y - y_n) - \varphi \rangle.$$

Moreover, by the Lebesgue Theorem,

$$|\langle v_n, \varphi(\cdot + y - y_n) - \varphi \rangle| \leq C \|\varphi(\cdot - y_n) - \varphi(\cdot - y)\| = C \|\varphi(\cdot - y_n) - \varphi(\cdot - y)\|_{H^1(K)} = o_n(1)$$

where $K \subset \mathbb{R}^3$ is a suitable compact set, completing the proof. \square

Now we are able to give the proof of Lemma 4.5.

Proof of Lemma 4.5. By Lemma B.1 we know that $\mathcal{J}'_q(u_0) = 0$. Let us divide the proof in various steps.

STEP 1: We have two possibilities.

Case 1a: If $u_n \rightarrow u_0$ in $H^1(\mathbb{R}^3)$, then the first alternative in the Lemma follows and the proof is concluded.

Case 1b: If $u_n \not\rightarrow u_0$ in $H^1(\mathbb{R}^3)$, then we set $u_n^{(1)} := u_n - u_0$, which satisfies, in view of Lemma B.4 and Lemma B.2, we have:

- (1bi) $u_n^{(1)} \rightharpoonup 0$ in $H^1(\mathbb{R}^3)$,
- (1bii) $\mathcal{J}'_q(u_n^{(1)}) \rightarrow 0$,
- (1biii) $\mathcal{J}_q(u_n^{(1)}) \rightarrow d - \mathcal{J}_q(u_0)$.

Moreover, again by Lemma B.4, we have that

$$(B.1) \quad \exists \{z_n^{(1)}\} \subset \mathbb{Z}^3 \text{ with } |z_n^{(1)}| \rightarrow +\infty \text{ such that } \lim_n \|u_n^{(1)}\|_{L^p(z_n^{(1)}+Q)} > 0.$$

Setting $\tilde{u}_n^{(1)} := u_n^{(1)}(\cdot - z_n^{(1)})$, we easily get from (1bi)–(1biii) and (B.1) that

$$\{\tilde{u}_n^{(1)}\} \text{ is bounded in } H^1(\mathbb{R}^3), \quad \mathcal{J}'_q(\tilde{u}_n^{(1)}) \rightarrow 0, \quad \tilde{u}_n^{(1)} \not\rightarrow 0 \text{ in } H^1(\mathbb{R}^3).$$

Then

$$\tilde{u}_n^{(1)} \rightharpoonup w_1 \neq 0 \text{ in } H^1(\mathbb{R}^3)$$

and, by the invariance under translations of the functional and (1biii) we have

$$(B.2) \quad \mathcal{J}_q(\tilde{u}_n^{(1)}) = \mathcal{J}_q(u_n^{(1)}) \rightarrow d - \mathcal{J}_q(u_0),$$

so that $\{\tilde{u}_n^{(1)}\}$ is a bounded (PS) sequence for \mathcal{J}_q . By Lemma B.1,

$$\mathcal{J}'_q(w_1) = 0 \quad \text{with } w_1 \neq 0.$$

STEP 2: Now there are two possibilities.

Case 2a: If $\tilde{u}_n^{(1)} \rightarrow w_1$ in $H^1(\mathbb{R}^3)$, this means that

$$o_n(1) = \|u_n^{(1)} - w_1(\cdot + z_n^{(1)})\| = \|u_n - u_0 - w_1(\cdot + z_n^{(1)})\|$$

and then $\mathcal{J}_q(\tilde{u}_n^{(1)}) = \mathcal{J}_q(u_n^{(1)}) \rightarrow \mathcal{J}_q(w_1)$, which, taking into account (B.2) gives

$$d = \mathcal{J}_q(u_0) + \mathcal{J}_q(w_1)$$

and the Lemma is proved with $\ell = 1$.

Case 2b: If $\tilde{u}_n^{(1)} \not\rightarrow w_1$, then let $u_n^{(2)} := u_n^{(1)} - w_1(\cdot + z_n^{(1)}) \not\rightarrow 0$. The sequence $\{u_n^{(2)}\}$ satisfies:

- (2bi) $u_n^{(2)} \rightharpoonup 0$ in $H^1(\mathbb{R}^3)$,
- (2bii) $\mathcal{J}'_q(u_n^{(2)}) \rightarrow 0$,

(2biii) $\mathcal{J}_q(u_n^{(2)}) \rightarrow d - \mathcal{J}_q(u_0) - \mathcal{J}_q(w_1)$, since, by Lemma B.2,

$$\mathcal{J}_q(u_n^{(2)}) = \mathcal{J}_q(\tilde{u}_n^{(1)} - w_1) = \mathcal{J}_q(\tilde{u}_n^{(1)}) - \mathcal{J}_q(w_1) + o_n(1) = d - \mathcal{J}_q(u_0) - \mathcal{J}_q(w_1) + o_n(1).$$

Again we have also that

$$(B.3) \quad \exists \{z_n^{(2)}\} \subset \mathbb{Z}^3 \text{ with } |z_n^{(2)}| \rightarrow +\infty \text{ such that } \lim_n \|u_n^{(2)}\|_{L^p(z_n^{(2)}+Q)} > 0.$$

Setting $\tilde{u}_n^{(2)} := u_n^{(2)}(\cdot - z_n^{(2)})$, it holds as before that

$$\{\tilde{u}_n^{(2)}\} \text{ is bounded in } H^1(\mathbb{R}^3), \quad \mathcal{J}'_q(\tilde{u}_n^{(2)}) \rightarrow 0, \quad \tilde{u}_n^{(2)} \rightharpoonup w_2 \neq 0 \text{ in } H^1(\mathbb{R}^3).$$

Then $\{\tilde{u}_n^{(2)}\}$ is a bounded (PS) sequence for \mathcal{J}_q and by Lemma B.1,

$$(B.4) \quad \mathcal{J}'_q(w_2) = 0 \quad \text{with } w_2 \neq 0.$$

Moreover

$$(B.5) \quad |z_n^{(1)} - z_n^{(2)}| \rightarrow +\infty.$$

To see this, first observe that

$$\tilde{u}_n^{(1)} - w_1 = u_n^{(1)}(\cdot - z_n^{(1)}) - w_1 = u_n^{(2)}(\cdot - z_n^{(1)}) = \tilde{u}_n^{(2)}(\cdot + z_n^{(2)} - z_n^{(1)}).$$

Then if it were $|z_n^{(2)} - z_n^{(1)}| \leq R$, since $\tilde{u}_n^{(2)} \not\rightarrow 0$, by Lemma B.5 item (ii), we deduce

$$\tilde{u}_n^{(2)}(\cdot + z_n^{(2)} - z_n^{(1)}) \not\rightarrow 0,$$

which is a contradiction.

STEP3: Again we have two possibilities.

Case 3a: If $\tilde{u}_n^{(2)} \rightarrow w_2$ in $H^1(\mathbb{R}^3)$ this means that

$$(B.6) \quad \begin{aligned} o_n(1) &= \|\tilde{u}_n^{(2)} - w_2\| \\ &= \|u_n^{(2)} - w_2(\cdot + z_n^{(2)})\| \\ &= \|u_n^{(1)} - w_1(\cdot + z_n^{(1)}) - w_2(\cdot + z_n^{(2)})\| \\ &= \|u_n - u_0 - w_1(\cdot + z_n^{(1)}) - w_2(\cdot + z_n^{(2)})\| \end{aligned}$$

and then, being $\mathcal{J}_q(\tilde{u}_n^{(2)}) = \mathcal{J}_q(u_n^{(2)})$ the Lemma holds, in virtue of (B.3)–(B.6) with $\ell = 2$.

Case 3b: If $\tilde{u}_n^{(2)} \not\rightarrow w_2$, we argue as before repeating the procedure.

In this way we obtain at the generic

STEPm with the following alternatives:

Case ma: $\tilde{u}_n^{(m-1)} \rightarrow w_{m-1}$ in $H^1(\mathbb{R}^3)$ and the Lemma holds with $\ell = m - 1$.

Case mb: We have

- sequences of points $\{z_n^{(i)}\} \subset \mathbb{R}^3$ for $i = 1, \dots, m$ with $|z_n^{(i)}| \rightarrow +\infty$ for all $i = 1, \dots, m$ and $|z_n^{(i)} - z_n^{(j)}| \rightarrow +\infty$ for all $i, j = 1, \dots, m$ with $i \neq j$;
- functions $w_i \neq 0$ with $\mathcal{J}'_q(w_i) = 0$ for all $i = 1, \dots, m$;

and in this case the procedure continues.

However at some step $(\ell + 1)$ the first case has to occur stopping the process and proving the Lemma. That is there exists $\ell \in \mathbb{N}$ such that $\tilde{u}_n^{(\ell)} \rightarrow w_\ell$. To see this, we first observe that, for any $N \in \mathbb{N}$ we have that

$$(B.7) \quad \left\| u_n - u_0 - \sum_{i=1}^N w_i(\cdot + z_n^{(i)}) \right\|^2 = \|u_n\|^2 - \|u_0\|^2 - \sum_{i=1}^N \|w_i\|^2 + o_n(1)$$

Indeed expanding the left hand side above we have

$$(B.8) \quad \left\| u_n - u_0 - \sum_{i=1}^N w_i(\cdot + z_n^{(i)}) \right\|^2 = \|u_n\|^2 + \|u_0\|^2 + \left\| \sum_{i=1}^N w_i(\cdot + z_n^{(i)}) \right\|^2 - 2\langle u_n, u_0 \rangle \\ - 2 \sum_{i=1}^N \langle u_n, w_i(\cdot + z_n^{(i)}) \rangle - 2 \sum_{i=1}^N \langle u_0, w_i(\cdot + z_n^{(i)}) \rangle.$$

Now, since $|z_n^{(i)} - z_n^{(j)}| \rightarrow +\infty$ for $i \neq j$, by (i) in Lemma B.5,

$$(B.9) \quad \left\| \sum_{i=1}^N w_i(\cdot + z_n^{(i)}) \right\|^2 = \sum_{i=1}^N \left\| w_i(\cdot + z_n^{(i)}) \right\|^2 + 2 \sum_{i \neq j} \langle w_i(\cdot + z_n^{(i)}), w_j(\cdot + z_n^{(j)}) \rangle = \sum_{i=1}^N \|w_i\|^2 + o_n(1).$$

Analogously, being $|z_n^{(i)}| \rightarrow +\infty$,

$$(B.10) \quad \langle u_0, w_i(\cdot + z_n^{(i)}) \rangle = o_n(1).$$

Finally,

$$(B.11) \quad \langle u_n, w_i(\cdot + z_n^{(i)}) \rangle = \langle u_n - u_0 - \sum_{j=1}^{i-1} w_j(\cdot + z_n^{(j)}), w_i(\cdot + z_n^{(i)}) \rangle + \langle u_0 + \sum_{j=1}^{i-1} w_j(\cdot + z_n^{(j)}), w_i(\cdot + z_n^{(i)}) \rangle \\ = \langle u_n^{(i)}, w_i(\cdot + z_n^{(i)}) \rangle + \langle u_0, w_i(\cdot + z_n^{(i)}) \rangle + \sum_{j=1}^{i-1} \langle w_j, w_i(\cdot + z_n^{(i)} - z_n^{(j)}) \rangle \\ = \langle \tilde{u}_n^{(i)}, w_i \rangle + o_n(1) \\ = \|w_i\|^2 + o_n(1).$$

Then by plugging (B.9)-(B.11) into (B.8) we get (B.7).

In virtue of the fact that w_i are nontrivial critical points of \mathcal{J}_q , we have

$$\|w_i\|^2 \leq \|w_i\|^2 + \int \phi_{w_i} w_i^2 = \|w_i\|_p^p \leq C \|w_i\|^p$$

showing that $\{w_i\}$ are bounded away from zero in $H^1(\mathbb{R}^3)$.

Then, by (B.7) we deduce that the process has to stop, completing the proof of the Lemma. \square

REFERENCES

- [1] R.A. Adams, *Sobolev Spaces*, Academic Press, 1975. [21](#)
- [2] A. Ambrosetti, P.H. Rabinowitz, *Dual variational methods in critical point theory and applications*, J. Funct. Anal. **14** (1973), 349–381. [12](#)
- [3] A. Azzollini, P. d’Avenia, A. Pomponio, *On the Schrödinger-Maxwell equations under the effect of a general nonlinear term*, Ann. Inst. H. Poincaré Anal. Non Linéaire **27** (2010), no. 2, 779–791. [3](#), [4](#)
- [4] A. Azzollini, A. Pomponio, G. Siciliano, *On the Schrödinger-Born-Infeld system*, preprint, arXiv:[1710.03151](#). [4](#)
- [5] V. Benci and D. Fortunato, TMNA 1998. *An eigenvalue problem for the Schrödinger-Maxwell equations*, Topol. Methods Nonlinear Anal. **11** (1998), no. 2, 283–293. [11](#)
- [6] V. Benci and D. Fortunato, *Solitary waves of the nonlinear Klein-Gordon equation coupled with the Maxwell equations*, Rev. Math. Phys. **14** (2002), 409–420. [11](#)
- [7] M.C. Bertin, B.M. Pimentel, C.E. Valcárcel, G.E.R. Zambrano, *Hamilton-Jacobi formalism for Podolsky’s electromagnetic theory on the null-plane*, J. Math. Phys. **58** (2017), 082902. [3](#)
- [8] D. Bonheure, J.-B. Castéras, E. Moreira dos Santos, R. Nascimento, *Orbitally stable standing waves of a mixed dispersion nonlinear Schrödinger equation*, preprint, arXiv:[1710.09775](#). [3](#)
- [9] D. Bonheure, P. d’Avenia, A. Pomponio, *On the electrostatic Born-Infeld equation with extended charges*, Comm. Math. Phys. **346** (2016), no. 3, 877–906. [4](#)

- [10] C.A. Bonin, B.M. Pimentel, P.H. Ortega, *Multipole Expansion in Generalized Electrodynamics*, preprint, arXiv:1608.00902. [3](#)
- [11] F. Bopp, *Eine Lineare Theorie des Elektrons*, Ann. Phys. **430** (1940), 345–384. [2](#)
- [12] M. Born, *Modified field equations with a finite radius of the electron*, Nature **132** (1933), 282. [2](#)
- [13] M. Born, *On the quantum theory of the electromagnetic field*, Proc. Roy. Soc. London Ser. A **143** (1934), 410–437. [2](#)
- [14] M. Born, L. Infeld, *Foundations of the new field theory*, Nature **132** (1933), 1004. [2](#)
- [15] M. Born, L. Infeld, *Foundations of the new field theory*, Proc. Roy. Soc. London Ser. A **144** (1934), 425–451. [2](#)
- [16] H. Brezis, *Functional analysis, Sobolev spaces and partial differential equations*, Springer, 2011. [7](#)
- [17] R. Bufalo, B.M. Pimentel, D.E. Soto, *Causal approach for the electron-positron scattering in generalized quantum electrodynamics*, Phys. Rev. D **90** (2014), 085012. [3](#)
- [18] R. Bufalo, B.M. Pimentel, D.E. Soto, *Normalizability analysis of the generalized quantum electrodynamics from the causal point of view*, International Journal of Modern Physics A **32** (2017), 1750165. [3](#)
- [19] P.L. Cunha, P. d'Avenia, A. Pomponio, G. Siciliano, *A multiplicity result for Chern-Simons-Schrödinger equation with a general nonlinearity*, NoDEA Nonlinear Differential Equations Appl. **22** (2015), 1831–1850. [4](#)
- [20] R.R. Cuzinatto, C.A.M. de Melo, L.G. Medeiros, B.M. Pimentel, P.J. Pompeia, *Bopp-Podolsky black holes and the no-hair theorem*, Eur. Phys. J. C **78** (2018), 43. [3](#)
- [21] R.R. Cuzinatto, E.M. De Moraes, L.G. Medeiros, C. Naldoni De Souza, B.M. Pimentel, *De Broglie-Proca and Bopp-Podolsky massive photon gases in cosmology*, EPL **118** (2017), 19001. [3](#)
- [22] T. D'Aprile, D. Mugnai, *Non-Existence Results for the Coupled Klein-Gordon-Maxwell Equations*, Adv. Nonlinear Stud. **4** (2004), 307–322. [23](#), [24](#)
- [23] P. d'Avenia, L. Pisani, *Nonlinear Klein-Gordon equations coupled with Born-Infeld type equations*, Electron. J. Differential Equations 2002 (2002), 26. [4](#)
- [24] P. d'Avenia, L. Pisani, G. Siciliano, *Klein-Gordon-Maxwell systems in a bounded domain*, Discrete Contin. Dyn. Syst. **26** (2010), 135–149. [4](#)
- [25] G. Fibich, B. Ilan, G. Papanicolaou, *Self-Focusing with Fourth-Order Dispersion*, SIAM J. Appl. Math. **62** (2002), 1437–1462. [3](#)
- [26] G.M. Figueiredo, G. Siciliano, *Existence and asymptotic behaviour of solutions for a quasi-linear Schrödinger-Poisson system under a critical nonlinearity*, preprint, arXiv:1707.05353. [3](#), [4](#)
- [27] J. Frenkel, *4/3 problem in classical electrodynamics*, Phys. Rev. E **54** (1996), 5859–5862. [2](#)
- [28] D. Gilbarg, N.S. Trudinger, *Elliptic Partial Differential Equations of Second Order*, Springer, 1997. [21](#)
- [29] L. Jeanjean, *On the existence of bounded Palais-Smale sequences and application to a Landesman-Lazer-type problem set on \mathbb{R}^N* , Proc. Roy. Soc. Edinburgh Sect. A **129** (1999), 787–809. [4](#), [17](#)
- [30] L. Jeanjean, S. Le Coz, *An existence and stability result for standing waves of nonlinear Schrödinger equations*, Adv. Differential Equations **11** (2006), 813–840. [3](#)
- [31] H. Kikuchi, *Existence and stability of standing waves for Schrödinger-Poisson-Slater equation*, Adv. Nonlinear Stud. **7** (2007), 403–437. [3](#)
- [32] E.H. Lieb, M. Loss, *Analysis*, AMS, 2001. [9](#), [10](#)
- [33] G. Mie, *Grundlagen einer Theorie der Materie*, Ann. Phys. **345** (1913), 1–66. [2](#)
- [34] B. Podolsky, *A Generalized Electrodynamics*, Phys. Rev. **62** (1942), 68–71. [2](#), [5](#), [8](#)
- [35] D. Ruiz, *The Schrödinger-Poisson equation under the effect of a nonlinear local term*, J. Funct. Anal. **237** (2006), 655–674. [3](#), [13](#)
- [36] W.A. Strauss, *Existence of solitary waves in higher dimensions*, Comm. Math. Phys. **55** (1977), 149–162. [16](#)
- [37] M. Struwe, *On the evolution of harmonic mappings of Riemannian surfaces*, Comment. Math. Helvetici **60** (1985), 558–581. [4](#)
- [38] M. Willem, *Minimax theorems*, Birkhäuser, 1996. [14](#)

(P. d'Avenia)

DIPARTIMENTO DI MECCANICA, MATEMATICA E MANAGEMENT
POLITECNICO DI BARI
VIA ORABONA 4, 70125 BARI, ITALY
Email address: pietro.davenia@poliba.it

(G. Siciliano)

DEPARTAMENTO DE MATEMÁTICA - INSTITUTO DE MATEMÁTICA E ESTATÍSTICA
UNIVERSIDADE DE SÃO PAULO
RUA DO MATÃO 1010, 05508-090 SÃO PAULO, BRAZIL
Email address: sicilian@ime.usp.br