

Repository Istituzionale dei Prodotti della Ricerca del Politecnico di Bari

A Variational Analysis of a Gauged Nonlinear Schrödinger Equation.

This is a post print of the following article *Original Citation:* A Variational Analysis of a Gauged Nonlinear Schrödinger Equation / Pomponio, Alessio; Ruiz, D., - In: JOURNAL OF THE EUROPEAN MATHEMATICAL SOCIETY. - ISSN 1435-9855. - 17:6(2015), pp. 1463-1486. [10.4171/JEMS/535]

Availability: This version is available at http://hdl.handle.net/11589/6703 since: 2022-06-06

Published version DOI:10.4171/JEMS/535

Terms of use:

(Article begins on next page)

03 May 2024

A VARIATIONAL ANALYSIS OF A GAUGED NONLINEAR SCHRÖDINGER EQUATION

ALESSIO POMPONIO¹ AND DAVID RUIZ²

ABSTRACT. This paper is motivated by a gauged Schrödinger equation in dimension 2 including the so-called Chern-Simons term. The study of radial stationary states leads to the nonlocal problem:

$$-\Delta u(x) + \left(\omega + \frac{h^2(|x|)}{|x|^2} + \int_{|x|}^{+\infty} \frac{h(s)}{s} u^2(s) \, ds\right) u(x) = |u(x)|^{p-1} u(x),$$

where

$$h(r) = \frac{1}{2} \int_0^r s u^2(s) \, ds.$$

This problem is the Euler-Lagrange equation of a certain energy functional. In this paper the study of the global behavior of such functional is completed. We show that for $p \in (1,3)$, the functional may be bounded from below or not, depending on ω . Quite surprisingly, the threshold value for ω is explicit. From this study we prove existence and non-existence of positive solutions.

1. INTRODUCTION

In this paper we are concerned with a planar gauged Nonlinear Schrödinger Equation:

(1)
$$iD_0\phi + (D_1D_1 + D_2D_2)\phi + |\phi|^{p-1}\phi = 0.$$

Here $t \in \mathbb{R}$, $x = (x_1, x_2) \in \mathbb{R}^2$, $\phi : \mathbb{R} \times \mathbb{R}^2 \to \mathbb{C}$ is the scalar field, $A_\mu : \mathbb{R} \times \mathbb{R}^2 \to \mathbb{R}$ are the components of the gauge potential and $D_\mu = \partial_\mu + iA_\mu$ is the covariant derivative ($\mu = 0, 1, 2$).

The classical equation for the gauge potential A_{μ} is the Maxwell equation. However, the modified gauge field equation proposes to include the so-called Chern-Simons term into that equation (see for instance [23, Chapter 1]):

(2)
$$\partial_{\mu}F^{\mu\nu} + \frac{1}{2}\kappa\epsilon^{\nu\alpha\beta}F_{\alpha\beta} = j^{\nu}, \text{ with } F_{\mu\nu} = \partial_{\mu}A_{\nu} - \partial_{\nu}A_{\mu}$$

In the above equation, κ is a parameter that measures the strength of the Chern-Simons term. As usual, $\epsilon^{\nu\alpha\beta}$ is the Levi-Civita tensor, and super-indices are related to the Minkowski metric with signature (1, -1, -1). Finally, j^{μ} is the conserved matter current,

$$j^0 = |\phi|^2, \ j^i = 2 \text{Im} \left(\bar{\phi} D_i \phi \right).$$

²⁰¹⁰ Mathematics Subject Classification. 35J20, 35Q55.

Key words and phrases. Gauged Schrödinger Equations, Chern-Simons theory, Variational methods, concentration compactness.

A.P. is supported by M.I.U.R. - P.R.I.N. "Metodi variazionali e topologici nello studio di fenomeni non lineari", by GNAMPA Project "Metodi Variazionali e Problemi Ellittici Non Lineari" and by FRA2011 "Equazioni ellittiche di tipo Born-Infeld". D.R. is supported by the Spanish Ministry of Science and Innovation under Grant MTM2011-26717 and by J. Andalucia (FQM 116).

At low energies, the Maxwell term becomes negligible and can be dropped, giving rise to:

(3)
$$\frac{1}{2}\kappa\epsilon^{\nu\alpha\beta}F_{\alpha\beta} = j^{\nu}.$$

See [7,8,12–14] for the discussion above.

For the sake of simplicity, let us fix $\kappa = 2$. Equations (1) and (3) lead us to the problem:

(4)
$$iD_{0}\phi + (D_{1}D_{1} + D_{2}D_{2})\phi + |\phi|^{p-1}\phi = 0, \\ \partial_{0}A_{1} - \partial_{1}A_{0} = \operatorname{Im}(\bar{\phi}D_{2}\phi), \\ \partial_{0}A_{2} - \partial_{2}A_{0} = -\operatorname{Im}(\bar{\phi}D_{1}\phi), \\ \partial_{1}A_{2} - \partial_{2}A_{1} = \frac{1}{2}|\phi|^{2}.$$

As usual in Chern-Simons theory, problem (4) is invariant under gauge transformation,

(5)
$$\phi \to \phi e^{i\chi}, \quad A_{\mu} \to A_{\mu} - \partial_{\mu}\chi,$$

for any arbitrary C^{∞} function χ .

This model was first proposed and studied in [12–14], and sometimes has received the name of Chern-Simons-Schrödinger equation. The initial value problem, wellposedness, global existence and blow-up, scattering, etc. has been addressed in [2, 9, 11, 18, 19] for the case p = 3. See also [17] for a global existence result in the defocusing case.

The existence of stationary states for (4) and general p > 1 has been studied recently in [4] (with respect to that paper, our notation interchanges the indices 1 and 2). By using the ansatz:

$$\begin{aligned} \phi(t,x) &= u(|x|)e^{i\omega t}, & A_0(x) &= A_0(|x|), \\ A_1(t,x) &= -\frac{x_2}{|x|^2}h(|x|), & A_2(t,x) &= \frac{x_1}{|x|^2}h(|x|), \end{aligned}$$

in [4] it is found that *u* solves the equation: (6)

$$-\Delta u(x) + \left(\omega + \xi + \frac{h^2(|x|)}{|x|^2} + \int_{|x|}^{+\infty} \frac{h(s)}{s} u^2(s) \, ds\right) u(x) = |u(x)|^{p-1} u(x), \quad x \in \mathbb{R}^2,$$

where

$$h(r) = \frac{1}{2} \int_0^r s u^2(s) \, ds.$$

Here ξ in \mathbb{R} is an integration constant of A_0 , which takes the form:

$$A_0(r) = \xi + \int_r^{+\infty} \frac{h(s)}{s} u^2(s) \, ds.$$

Observe that (6) is a nonlocal equation. Moreover, in [4] it is shown that (6) is indeed the Euler-Lagrange equation of the energy functional:

$$I_{\omega+\xi}: H^1_r(\mathbb{R}^2) \to \mathbb{R},$$

defined as

$$I_{\omega+\xi}(u) = \frac{1}{2} \int_{\mathbb{R}^2} \left(|\nabla u(x)|^2 + (\omega+\xi)u^2(x) \right) dx + \frac{1}{8} \int_{\mathbb{R}^2} \frac{u^2(x)}{|x|^2} \left(\int_0^{|x|} su^2(s) \, ds \right)^2 dx - \frac{1}{p+1} \int_{\mathbb{R}^2} |u(x)|^{p+1} \, dx.$$

Here $H_r^1(\mathbb{R}^2)$ denotes the Sobolev space of radially symmetric functions. It is important to observe that the energy functional $I_{\omega+\xi}$ presents a competition between the nonlocal term and the local nonlinearity. The study of the behavior of the functional under this competition is one of the main motivations of this paper.

Given a stationary solution, and taking $\chi = ct$ in the gauge invariance (5), we obtain another stationary solution; the functions u(x), $A_1(x)$, $A_2(x)$ are preserved, and

$$\omega \to \omega + c, \quad A_0(x) \to A_0(x) - c$$

Therefore, the constant $\omega + \xi$ is a gauge invariant of the stationary solutions of the problem. By the above discussion we can take $\xi = 0$ in what follows, that is,

$$\lim_{|x| \to +\infty} A_0(x) = 0,$$

which was indeed assumed in [2,14].

For p > 3, it is shown in [4] that I_{ω} is unbounded from below, so it exhibits a mountain-pass geometry. In a certain sense, in this case the local nonlinearity dominates the nonlocal term. However the existence of a solution is not so direct, since for $p \in (3,5)$ the (PS) property is not known to hold. This problem is bypassed in [4] by using a constrained minimization taking into account the Nehari and Pohozaev identities, in the spirit of [20]. Moreover, infinitely many solutions have been found in [10] for p > 5 (possibly sign-changing).

A special case in the above equation is p = 3: in this case, static solutions can be found by passing to a self-dual equation, which leads to a Liouville equation that can be solved explicitly. Those are the unique positive solutions, as proved in [4]. For more information on the self-dual equations, see [5, 14, 23].

In case $p \in (1,3)$, solutions are found in [4] as minimizers on a L^2 sphere. Therefore, the value ω comes out as a Lagrange multiplier, and it is not controlled. Moreover, the global behavior of the energy functional I_{ω} is not studied.

The main purpose of this paper is to study whether I_{ω} is bounded from below or not for $p \in (1,3)$. In this case, the nonlocal term prevails over the local nonlinearity, in a certain sense. As we shall see, the situation is quite rich and unexpected a priori, and very different from the usual Nonlinear Schrödinger Equation. This situation differs also from the Schrödinger-Poisson problem (see [20]), which is another problem presenting a competition between local and nonlocal nonlinearities.

We shall prove the existence of a threshold value ω_0 such that I_{ω} is bounded from below if $\omega \ge \omega_0$, and it is not for $\omega \in (0, \omega_0)$. But, in our opinion, what is most surprising is that ω_0 has an explicit expression, namely:

(7)
$$\omega_0 = \frac{3-p}{3+p} 3^{\frac{p-1}{2(3-p)}} 2^{\frac{2}{3-p}} \left(\frac{m^2(3+p)}{p-1}\right)^{-\frac{p-1}{2(3-p)}},$$

with

$$n = \int_{-\infty}^{+\infty} \left(\frac{2}{p+1}\cosh^2\left(\frac{p-1}{2}r\right)\right)^{\frac{2}{1-p}} dr.$$

Let us give an idea of the proofs. It is not difficult to show that I_{ω} is coercive when the problem is posed on a bounded domain. So, there exists a minimizer u_n on the ball B(0, n) with Dirichlet boundary conditions. To prove boundedness of u_n , the problem is the possible loss of mass at infinity as $n \to +\infty$. The core of our proofs is a detailed study of the behavior of those masses. We are able to show that, if unbounded, the sequence u_n behaves as a soliton, if u_n is interpreted as a function of a single real variable. The proof uses a careful study of the level sets of u_n , which take into account the effect of the nonlocal term. Then, the energy functional I_{ω} admits a natural approximation through a convenient limit functional. Finally, the solutions of that limit functional, and their energy, can be found explicitly, so we can find ω_0 . See Section 2 for an heuristic explanation of the proof and a derivation of the limit functional.

Regarding the existence of solutions, a priori, the global minimizer could correspond to the zero solution. And indeed this is the case for large ω . Instead, we show that $\inf I_{\omega} < 0$ if $\omega > \omega_0$ is close to the threshold value. Therefore, the global minimizer is not trivial, and corresponds to a positive solution. The mountain pass theorem will provide the existence of a second positive solution.

If $\omega < \omega_0$, I_{ω} is unbounded from below, and hence the geometric assumptions of the mountain-pass theorem are satisfied. However, the boundedness of (PS) sequences seems to be a hard question in this case. Solutions are found for almost all values of $\omega \in (0, \omega_0)$, by using the well-known monotonicity trick of Struwe [22] (see also [15]).

Our main results are the following:

Theorem 1.1. For ω_0 as given in (7), there holds:

- (i) if $\omega \in (0, \omega_0)$, then I_{ω} is unbounded from below;
- (ii) if $\omega = \omega_0$, then I_{ω_0} is bounded from below, not coercive and $\inf I_{\omega_0} < 0$;
- (iii) if $\omega > \omega_0$, then I_{ω} is bounded from below and coercive.

Regarding the existence of solutions, we obtain the following result:

Theorem 1.2. Consider (6) with $\xi = 0$. There exist $\bar{\omega} > \tilde{\omega} > \omega_0$ such that:

- (i) if $\omega > \overline{\omega}$, then (6) has no solutions different from zero;
- (ii) if $\omega \in (\omega_0, \tilde{\omega})$, then (6) admits at least two positive solutions: one of them is a global minimizer for I_{ω} and the other is a mountain-pass solution;
- (iii) for almost every $\omega \in (0, \omega_0)$ (6) admits a positive solution.

The rest of the paper is organized as follows. Section 2 is devoted to some preliminary results. Moreover, we give a heuristic presentation of our proofs, which motivates the definition of the limit functional. This limit functional is studied in detail in Section 3. Finally, in Section 4 we prove Theorems 1.1 and 1.2.

Acknowledgement. This work has been partially carried out during a stay of A.P. in Granada. He would like to express his deep gratitude to the Departamento de Análisis Matemático for the support and warm hospitality.

The authors thank the referee for some observations that have helped to improve the clarity of the exposition of our proofs.

2. Preliminaries

Let us first fix some notations. We denote by $H_r^1(\mathbb{R}^2)$ the Sobolev space of radially symmetric functions, and $\|\cdot\|$ its usual norm. Other norms, like Lebesgue norms, will be indicated with a subscript. In particular, $\|\cdot\|_{H^1(\mathbb{R})}$, $\|\cdot\|_{H^1(a,b)}$ are used to indicate the norms of the Sobolev spaces of dimension 1. If nothing is specified, strong and weak convergence of sequences of functions are assumed in the space $H^1(\mathbb{R}^2)$.

In our estimates, we will frequently denote by C > 0, c > 0 fixed constants, that may change from line to line, but are always independent of the variable under consideration. We also use the notations O(1), o(1), $O(\varepsilon)$, $o(\varepsilon)$ to describe the asymptotic behaviors of quantities in a standard way. Finally the letters x, y indicate two-dimensional variables and r, s denote one-dimensional variables.

Let us start with the following proposition, proved in [4]:

Proposition 2.1. I_{ω} is a C^1 functional, and its critical points correspond to classical solutions of (6).

Next result deals with the behavior of I_{ω} under weak limits in $H_r^1(\mathbb{R}^2)$. Even if it is not explicitly stated in this form, Proposition 2.2 follows easily from [4, Lemma 3.2] and the compactness of the embedding $H_r^1(\mathbb{R}^2) \hookrightarrow L^q(\mathbb{R}^2)$, $q \in (2, +\infty)$ (see [21]).

Proposition 2.2. If $u_n \rightharpoonup u$, then

$$\int_{\mathbb{R}^2} \frac{u_n^2(x)}{|x|^2} \left(\int_0^{|x|} s u_n^2(s) \, ds \right)^2 dx \to \int_{\mathbb{R}^2} \frac{u^2(x)}{|x|^2} \left(\int_0^{|x|} s u^2(s) \, ds \right)^2 dx.$$

In particular, I_{ω} is weak lower semicontinuous. Moreover, if $u_n \rightharpoonup u$ then $I'_{\omega}(u_n)(\varphi) \rightarrow I'_{\omega}(u)(\varphi)$ for all $\varphi \in H^1_r(\mathbb{R}^2)$.

To finish the account of preliminaries, we now state an inequality which will prove to be fundamental in our analysis. This inequality is proved in [4], where also the maximizers are found.

Proposition 2.3. For any $u \in H^1_r(\mathbb{R}^2)$, (8)

$$\int_{\mathbb{R}^2} |u(x)|^4 \, dx \leqslant 2 \left(\int_{\mathbb{R}^2} |\nabla u(x)|^2 \, dx \right)^{1/2} \left(\int_{\mathbb{R}^2} \frac{u^2}{|x|^2} \left(\int_0^{|x|} s u^2(s) \, ds \right)^2 dx \right)^{1/2}.$$

As commented in the introduction, this paper is concerned with boundedness from below of I_{ω} . Let us give a rough idea of the arguments of our proof. First of all, consider u(r) a fixed function, and define $u_{\rho}(r) = u(r-\rho)$. Let us now estimate $I_{\omega}(u_{\rho})$ as $\rho \to +\infty$.

$$(2\pi)^{-1}I_{\omega}(u_{\rho}) = \frac{1}{2} \int_{-\rho}^{+\infty} (|u'|^2 + \omega u^2)(r+\rho) dr + \frac{1}{8} \int_{-\rho}^{\infty} \frac{u^2(r)}{r+\rho} \left(\int_{-\rho}^{r} (s+\rho)u^2(s) ds \right)^2 dr - \frac{1}{p+1} \int_{-\rho}^{\infty} |u|^{p+1}(r+\rho) dr.$$

We estimate the above expression by simply replacing the expressions $(r + \rho)$, $(s + \rho)$ with the constant ρ :

$$\sim \rho \left[\frac{1}{2} \int_{-\infty}^{+\infty} (|u|'^2 + \omega u^2) \, dr + \frac{1}{8} \int_{-\infty}^{+\infty} u^2(r) \left(\int_{-\infty}^r u^2(s) \, ds \right)^2 \, dr - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr \right]$$
$$= \rho \left[\frac{1}{2} \int_{-\infty}^{+\infty} (|u|'^2 + \omega u^2) \, dr + \frac{1}{24} \left(\int_{-\infty}^{+\infty} u^2 dr \right)^3 - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr \right].$$

This estimate will be made rigorous in Lemma 4.1. Therefore, it is natural to define the limit functional $J_{\omega}: H^1(\mathbb{R}) \to \mathbb{R}$,

$$J_{\omega}(u) = \frac{1}{2} \int_{-\infty}^{+\infty} \left(|u'|^2 + \omega u^2 \right) dr + \frac{1}{24} \left(\int_{-\infty}^{+\infty} u^2 dr \right)^3 - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} dr.$$

As a consequence of the above argument, if J_{ω} attains negative values, then I_{ω} will be unbounded from below.

The reverse is also true, but the proof is more delicate. We will show that if u_n is unbounded in $H_r^1(\mathbb{R}^2)$ and $I_{\omega}(u_n)$ is bounded from above, then somehow u_n contains a certain mass spreading to infinity, as u_{ρ} does. This will be made explicit in Proposition 4.2. But this will lead us to a contradiction if J_{ω} is positive on that

mass. The proof of this argument is however far from trivial, and is the core of this paper.

Summing up, we are able to relate I_{ω} with the limit functional J_{ω} in the following way:

$$\inf I_{\omega} > -\infty \iff \inf J_{\omega} = 0.$$

Moreover this characterization will give us the threshold value for ω , since the critical points of J_{ω} can be found explicitly, as will be shown in next Section.

3. The limit problem

In this section we deal with the limit functional $J_{\omega}: H^1(\mathbb{R}) \to \mathbb{R}$,

(9)
$$J_{\omega}(u) = \frac{1}{2} \int_{-\infty}^{+\infty} \left(|u'|^2 + \omega u^2 \right) dr + \frac{1}{24} \left(\int_{-\infty}^{+\infty} u^2 \, dr \right)^3 - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr.$$

Clearly, the Euler-Lagrange equation of (9) is the following problem:

(10)
$$-u'' + \omega u + \frac{1}{4} \left(\int_{-\infty}^{+\infty} u^2(s) \, ds \right)^2 u = |u|^{p-1} u, \quad \text{in } \mathbb{R}.$$

As we shall see later, we will find the explicit solutions of (10) later. But, first, let us study it from a variational point of view: this study will give us some further information on the solutions.

Before going on, we need a technical result, which is stated in next lemma. We think that such result must be well-known, but we have not been able to find a explicit reference.

Lemma 3.1. Let $u_n \in H^1(\mathbb{R})$ a sequence of even non-negative functions which are decreasing in r > 0, and assume that $u_n \rightarrow u_0$ weakly in $H^1(\mathbb{R})$. Then u_0 is also even, non-negative and decreasing in r > 0, and $u_n \rightarrow u_0$ in $L^q(\mathbb{R})$ for any $q \in (2, +\infty)$.

Proof. Observe that the set $A = \{u \in H^1(\mathbb{R}) \text{ nonnegative, even and decreasing in } r > 0\}$ is a closed and convex subset of $H^1(\mathbb{R})$. As a consequence, $u_0 \in A$.

Then, for any $r \in \mathbb{R}$, $r \neq 0$,

$$C \geqslant \left| \int_0^r u_n^2(s) \, ds \right| \geqslant u_n^2(r) |r| \Rightarrow u_n(r) \leqslant \frac{C}{\sqrt{|r|}},$$

and the same estimate works for u_0 . With this inequality, we can estimate:

$$\int_{-\infty}^{+\infty} |u_n - u_0|^q \, dr \leqslant \int_{-R}^{R} |u_n - u_0|^q \, dr + 2C \int_{|r| > R} r^{-q/2} \, dr$$
$$= \int_{-R}^{R} |u_n - u_0|^q \, dr + 4C \frac{2}{2-q} R^{\frac{2-q}{2}}.$$

Take into account that, by Rellich-Kondrachov Theorem, $u_n \to u_0$ in $L^q(-R, R)$ for any R > 0 fixed. Then, the above inequality implies that $u_n \to u_0$ in $L^q(\mathbb{R})$.

Some of the properties of the functional J_{ω} are discussed below:

Proposition 3.2. Consider the functional J_{ω} with $p \in (1,3)$ and $\omega > 0$. The following properties hold:

- *a)* J_{ω} *is coercive and attains its infimum.*
- b) 0 is a local minimum of J_{ω} . Indeed, there exists $r_0 > 0$ with the following property:

for any $r \in (0, r_0)$, there exists $\alpha > 0$ satisfying that $J_{\omega}(u) > \alpha$, for any $u \in H^1(\mathbb{R})$ with $||u||_{H^1(\mathbb{R})} = r$.

c) There exists $\omega_0 > 0$ such that $\min J_{\omega} < 0$ if and only if $\omega \in [0, \omega_0)$.

Proof. **Proof of a)** To prove coercivity, we use Gagliardo-Nirenberg inequality:

$$||u||_{L^4} \leq C ||u'||_{L^2}^{1/4} ||u||_{L^2}^{3/4}.$$

Hence

$$\int_{-\infty}^{+\infty} u^4 dr \leqslant \frac{C}{2} \left[\int_{-\infty}^{+\infty} |u'|^2 dr + \left(\int_{-\infty}^{+\infty} u^2 dr \right)^3 \right].$$

Then, (11)

$$J_{\omega}(u) \ge \frac{1}{4} \int_{-\infty}^{+\infty} |u'|^2 \, dr + \frac{1}{48} \left(\int_{-\infty}^{+\infty} u^2 \, dr \right)^3 + c \int_{-\infty}^{+\infty} u^4 \, dr - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr.$$

Observe that for any C > 0 we can choose D > 0 so that $t^3 \ge Ct - D$ for every $t \ge 0$. Applying this with $t = \int_{-\infty}^{+\infty} u^2 dr$ into (11), and renaming C, we obtain:

$$J_{\omega}(u) \ge \frac{1}{4} \int_{-\infty}^{+\infty} |u'|^2 \, dr + \int_{-\infty}^{+\infty} \left(Cu^2 + cu^4 - \frac{1}{p+1} |u|^{p+1} \right) dr - D.$$

Now, it suffices to take *C* so that the function $Cu^2 + cu^4 - \frac{1}{p+1}|u|^{p+1} \ge 0$ for any $u \in \mathbb{R}$.

Take now u_n such that $J_{\omega}(u_n) \to \inf J_{\omega}$. From the coercivity, it follows that u_n is bounded. Consider now the sequence $v_n = |u_n|^*$ of non-negative symmetrized functions. Clearly, v_n is also bounded, and it is easy to observe that $\inf J_{\omega} \leq J_{\omega}(v_n) \leq J_{\omega}(u_n) \to \inf J_{\omega}$.

Assume, passing to a subsequence, that $v_n \rightharpoonup v$ weakly in $H^1(\mathbb{R})$. By Lemma 3.1, $v_n \rightarrow v$ in $L^{p+1}(\mathbb{R})$. The weak lower semicontinuity of the norm allows us to conclude that u is a minimizer of J_{ω} .

Proof of b) This is quite standard. Indeed, by using Sobolev inequality,

$$J_{\omega}(u) \ge \frac{1}{2} \min\{1, \omega\} \|u\|_{H^{1}(\mathbb{R})}^{2} - C \|u\|_{H^{1}(\mathbb{R})}^{p+1}.$$

Proof of c) Let us define the map $\phi : [0, +\infty) \to \mathbb{R}$, $\phi(\omega) = \min J_{\omega}$. It is easy to check that ϕ is increasing and continuous. Moreover, $\phi(\omega) \leq 0$ for all ω (observe that $J_{\omega}(0) = 0$).

We claim that $\phi(\omega) = 0$ for large ω . Indeed, by the same arguments of the proof of **a**):

$$J_{\omega}(u) \ge \int_{-\infty}^{+\infty} \left(\frac{\omega}{2}u^2 + cu^4 - \frac{1}{p+1}|u|^{p+1}\right) dr.$$

For ω sufficiently large, $\frac{\omega}{2}u^2 + cu^4 - \frac{1}{p+1}|u|^{p+1} \ge 0$ for any $u \in \mathbb{R}$. Then $J_{\omega}(u) \ge 0$ for any $u \in H^1(\mathbb{R})$, proving the claim.

We now show that $\phi(0) < 0$. On that purpose, fix $u \in H^1(\mathbb{R})$ and define $u_{\lambda}(r) = \lambda^{\frac{2}{p-1}} u(\lambda r)$. There holds:

$$J_0(u_\lambda) = \frac{1}{2}\lambda^{\frac{p+3}{p-1}} \int_{-\infty}^{+\infty} |u'|^2 \, dr + \frac{1}{24}\lambda^{\frac{3(5-p)}{p-1}} \left(\int_{-\infty}^{+\infty} u^2 \, dr\right)^3 - \frac{1}{p+1}\lambda^{\frac{p+3}{p-1}} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr$$

Therefore, for λ sufficiently small, $J_0(u_\lambda)$ has the sign of the term

$$\frac{1}{2} \int_{-\infty}^{+\infty} |u'|^2 \, dr - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} \, dr$$

It suffices to take *u* such that this quantity is negative to conclude.

So, we can define $\omega_0 = \min\{\omega \ge 0 : \phi(\omega) = 0\} > 0$.

As a consequence of the previous result, for $\omega \in [0, \omega_0)$ there exists a nontrivial solution for (10), which corresponds to a global minimum of J_{ω} . As announced in the introduction, the expression for ω_0 will found later on.

We now pass to finding the explicit solutions of problem (10). For any k > 0 we denote by $w_k \in H^1(\mathbb{R})$ the unique positive radial solution of:

(12)
$$-w_k'' + kw_k = w_k^p, \quad \text{in } \mathbb{R}$$

Let us state some well-known properties of this equation. First, the Hamiltonian of w_k is equal to 0, that is,

(13)
$$-\frac{1}{2}|w_k'(r)|^2 + \frac{k}{2}w_k^2(r) - \frac{1}{p+1}w_k^{p+1}(r) = 0, \text{ for all } r \in \mathbb{R}.$$

It is also known that any solution of (12) is of the form $u(x) = \pm w_k(x - y)$, for some $y \in \mathbb{R}$. Moreover,

(14)
$$w_k(r) = k^{\frac{1}{p-1}} w_1(\sqrt{k}r), \text{ where } w_1(r) = \left(\frac{2}{p+1}\cosh^2\left(\frac{p-1}{2}r\right)\right)^{\frac{1}{1-p}}.$$

In what follows we define

$$m = \int_{-\infty}^{+\infty} w_1^2 \, dr.$$

The following relations are also well known, and can be deduced from (13):

(15)
$$\int_{-\infty}^{+\infty} |w_1'|^2 dr = \frac{p-1}{p+3}m, \qquad \int_{-\infty}^{+\infty} w_1^{p+1} dr = \frac{2(p+1)}{p+3}m.$$

Proposition 3.3. *Let us consider the equation:*

(16)
$$k = \omega + \frac{1}{4}m^2 k^{\frac{5-p}{p-1}}, \ k > 0.$$

Then, u is a nontrivial solution of (10) if and only if $u(r) = w_k(r - \xi)$ for some $\xi \in \mathbb{R}$ and k a root of (16).

Define:

(17)
$$\omega_1 = \left(\frac{(5-p)m^2}{4(p-1)}\right)^{-\frac{p-1}{2(3-p)}} - \frac{m^2}{4} \left(\frac{(5-p)m^2}{4(p-1)}\right)^{-\frac{(5-p)}{2(3-p)}}$$

The following holds:

- (1) if $\omega > \omega_1$, equation (16) has no solution and there is no nontrivial solution of (10);
- (2) if $\omega = \omega_1$, equation (16) has only one solution k_0 and $w_{k_0}(r)$ is the only nontrivial solution of (10) (apart from translations);
- (3) if $\omega \in (0, \omega_1)$, equation (16) has two solutions $k_1(\omega) < k_2(\omega)$ and $w_{k_1}(r), w_{k_2}(r)$ are the only two non-trivial solutions of (10) (apart from translations).

Proof. Let u be a nontrivial solution of (10), and define $k = \omega + \frac{1}{4} \left(\int_{-\infty}^{+\infty} u^2 dr \right)^2$. Then, u is a solution of $-u'' + ku = u^p$, so $u(r) = w_k(r - \xi)$ for some $\xi \in \mathbb{R}$. By using (14), we obtain:

$$k = \omega + \frac{1}{4} \left(\int_{-\infty}^{+\infty} w_k^2(r) \, dr \right)^2 = \omega + \frac{1}{4} k^{\frac{4}{p-1}} \left(\int_{-\infty}^{+\infty} w_1^2(\sqrt{k}r) \, dr \right)^2.$$

A change of variables leads us to equation (16).

Moreover,

$$1 1.$$

Therefore, the function $(0, +\infty) \ni k \mapsto k^{\frac{5-p}{p-1}}$ is convex. Therefore, there exists $\omega_1 > 0$ with the properties indicated.

In order to get the exact value of ω_1 , observe that the function $k \mapsto \omega_1 + \frac{1}{4}m^2k^{\frac{5-p}{p-1}} - k$ has a degenerate 0. Then, ω_1 solves the system:

$$\begin{cases} \omega + \frac{m^2}{4}k^{\frac{5-p}{p-1}} = k, \\ \frac{5-p}{4(p-1)}m^2k^{\frac{5-p}{p-1}-1} = 1 \end{cases}$$

From this one obtains formula (17).

In our next result, we obtain information from Proposition 3.3.

Proposition 3.4. Let ω_0 , ω_1 be the values defined in Propositions 3.2 and 3.3. Then:

(1) $\omega_0 < \omega_1$, and ω_0 has the expression:

(18)
$$\omega_0 = \frac{3-p}{3+p} \, 3^{\frac{p-1}{2(3-p)}} \, 2^{\frac{2}{3-p}} \left(\frac{m^2(3+p)}{p-1}\right)^{-\frac{p-1}{2(3-p)}},$$

where m is as in (3).

(2) For any $\omega \in (0, \omega_1)$, $J_{\omega}(w_{k_1}) > J_{\omega}(w_{k_2})$. In particular, for any $\omega \in (0, \omega_0)$, w_{k_2} is a global minimizer of J_{ω} .

Proof. We consider the energy functional J_{ω} evaluated on the curve $k \mapsto w_k$. In the computations that follow we use (14) and change of variables. We have

$$\psi(k) := J_{\omega}(w_k) = \frac{k^{\frac{3+p}{2(p-1)}}}{2} \int_{-\infty}^{+\infty} |w_1'(r)|^2 dr + \omega \frac{k^{\frac{5-p}{2(p-1)}}}{2} \int_{-\infty}^{+\infty} w_1^2(r) dr + \frac{k^{\frac{3(5-p)}{2(p-1)}}}{24} \left(\int_{-\infty}^{+\infty} w_1^2(r) dr \right)^3 - \frac{k^{\frac{3+p}{2(p-1)}}}{p+1} \int_{-\infty}^{+\infty} |w_1(r)|^{p+1} dr.$$

Plugging (15) into that expression,

$$\psi(k) = m \left[\frac{p-5}{2(3+p)} k^{\frac{3+p}{2(p-1)}} + \frac{\omega}{2} k^{\frac{5-p}{2(p-1)}} + \frac{m^2}{24} k^{\frac{3(5-p)}{2(p-1)}} \right].$$

Then:

$$\frac{d}{dk}\psi(k) = m k^{\frac{7-3p}{2(p-1)}} \frac{5-p}{4(p-1)} \left[-k + \omega + \frac{1}{4}m^2 k^{\frac{5-p}{p-1}} \right]$$

In particular, the roots of (16) are exactly the critical points of ψ . Observe that:

$$\frac{5-p}{2(p-1)} < \frac{3+p}{2(p-1)} < \frac{3(5-p)}{2(p-1)}.$$

Then ψ is increasing near 0 (for $\omega > 0$) and near infinity. Therefore, for $\omega \in (0, \omega_1)$, its first root corresponds to a local maximum of ψ and the second one to a local minimum, so $J(w_{k_1}) > J(w_{k_2})$. Take now $\omega \in (0, \omega_0)$. Since in this case the minimizer is nontrivial, it must correspond to w_{k_2} . Moreover, $\omega_0 < \omega_1$.

In order to get the value of ω_0 , observe that $J_{\omega_0}(w_{k_2}) = 0$. Therefore, $\omega_0 > 0$ solves:

$$\begin{cases} \omega + \frac{1}{4}m^2k^{\frac{5-p}{p-1}} = k, \\ \frac{p-5}{2(3+p)}k^{\frac{3+p}{2(p-1)}} + \frac{\omega}{2}k^{\frac{5-p}{2(p-1)}} + \frac{m^2}{24}k^{\frac{3(5-p)}{2(p-1)}} = 0. \end{cases}$$

From there, expression (18) follows.

Remark 3.5. Observe that the map ψ defined in the proof of Proposition 3.4 gives us a quite clear interpretation of the functional J_{ω} . Indeed, k is a critical point of ψ if and only if w_k is a critical point of J_{ω} . Moreover, the following holds.

- (1) If $\omega > \omega_1$, ψ is positive and increasing without critical points.
- If ω = ω₁, ψ is still positive and increasing, but it has an inflection point at k = k₀.
- (3) If $\omega \in (0, \omega_1)$, ψ has a local maximum and minimum attained at k_1 and k_2 , respectively.
- (4) If $\omega = \omega_0$, $\psi(k_2) = 0$. Observe then, in this case, the minimum of J_{ω_0} is 0, and is attained at 0 and w_{k_2} .
- (5) If $\omega \in [0, \omega_0)$, $\psi(k_2) < 0$ and then w_{k_2} is the unique global minimizer, with $J_{\omega}(w_{k_2}) < 0$.

Remark 3.6. In general, we cannot obtain a more explicit expression of m depending on p, but it can be easily approximated by using some software. In Figure 1 the maps $\omega_0(p)$ and $\omega_1(p)$ have been plotted.



For some specific values of p, m can be explicitly computed, and hence ω_0 and ω_1 . For instance, if p = 2, m = 6, $\omega_1 = \frac{2}{9\sqrt{3}}$ and $\omega_0 = \frac{2}{5\sqrt{15}}$.

We finish this section with a technical result that will be of use later in the proof of Theorem 1.1.

Proposition 3.7. Assume $\omega \ge \omega_0$, and $u_n \in H^1(\mathbb{R})$ such that $J_{\omega}(u_n) \to 0$. There holds

- (1) if $\omega > \omega_0$, then $u_n \to 0$ in $H^1(\mathbb{R})$;
- (2) if $\omega = \omega_0$, then, up to a subsequence, either $u_n \to 0$ or $u_n(\cdot x_n) \to w_{k_2}$ in $H^1(\mathbb{R})$, for some sequence $x_n \in \mathbb{R}$.

Proof. Since J_{ω} is coercive, we have that u_n is bounded. If $u_n \to 0$ in $H^1(\mathbb{R})$, we are done. Otherwise, we have that:

$$o_n(1) = J_{\omega}(u_n) \ge \frac{1}{2} \int_{-\infty}^{+\infty} \left(|u'_n(r)|^2 + \omega u_n^2(r) \right) dr - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u_n(r)|^{p+1} dr.$$

Then, $u_n \not\rightarrow 0$ in $L^{p+1}(\mathbb{R})$. We can apply concentration-compactness lemma (see [16, Lemma I.1]), and there exists $\xi_n \in \mathbb{R}$ such that $\int_{\xi_n-1}^{\xi_n+1} u_n^2 \ge \varepsilon > 0$. Therefore,

 $\tilde{u}_n(r) = u_n(r - \xi_n) \rightharpoonup u \neq 0$ weakly in $H^1(\mathbb{R})$. Define $v_n = \tilde{u}_n - u$, which clearly converges weakly to 0 in $H^1(\mathbb{R})$.

Step 1: $v_n \to 0$ in $L^2(\mathbb{R})$. We just compute

$$\begin{split} \rho_n(1) &= J_{\omega}(u_n) = J_{\omega}(\tilde{u_n}) = J_{\omega}(v_n + u) \\ &= \frac{1}{2} \int_{-\infty}^{+\infty} \left(|v_n'|^2 + |u'|^2 + 2v_n'u' \right) dr + \frac{\omega}{2} \int_{-\infty}^{+\infty} \left(v_n^2 + u^2 + 2v_n u \right) dr \\ &+ \frac{1}{8} \left[\left(\int_{-\infty}^{+\infty} v_n^2 dr \right)^3 + \left(\int_{-\infty}^{+\infty} u^2 dr \right)^3 + 3 \left(\int_{-\infty}^{+\infty} v_n^2 dr \right)^2 \left(\int_{-\infty}^{+\infty} u^2 dr \right) \right. \\ &+ 3 \left(\int_{-\infty}^{+\infty} v_n^2 dr \right) \left(\int_{-\infty}^{+\infty} u^2 dr \right)^2 \right] - \frac{1}{p+1} \int_{-\infty}^{+\infty} |v_n + u|^{p+1} dr + o_n(1). \end{split}$$

Here the mixed products converge to zero, since $v_n \rightarrow 0$. Passing to a subsequence, we can assume that $v_n \rightarrow 0$ almost everywhere. Then, the well-known Brezis-Lieb lemma ([3]) implies that

$$\int_{-\infty}^{+\infty} |v_n + u|^{p+1} \, dr - \int_{-\infty}^{+\infty} (|v_n|^{p+1} + |u|^{p+1}) \, dr \to 0.$$

Then,

$$o_n(1) = J_{\omega}(u_n) = J_{\omega}(v_n) + J_{\omega}(u) + \frac{3}{8} \left[\left(\int_{-\infty}^{+\infty} v_n^2 \, dr \right)^2 \left(\int_{-\infty}^{+\infty} u^2 \, dr \right) + \left(\int_{-\infty}^{+\infty} v_n^2 \, dr \right) \left(\int_{-\infty}^{+\infty} u^2 \, dr \right)^2 \right] + o_n(1).$$

It is here that the assumption $\omega \ge \omega_0$ is crucial. Indeed, it implies that $J_{\omega}(v_n) \ge 0$, $J_{\omega}(u) \ge 0$. Recall that $u \ne 0$, to conclude the proof of Step 1.

Step 2: Conclusion.

By interpolation,

$$\|v_n\|_{L^{p+1}} \leqslant \|v_n\|_{L^2}^{\alpha} \|v_n\|_{L^{p+2}}^{1-\alpha},$$

for some $\alpha \in (0,1)$. Since v_n is bounded in $H^1(\mathbb{R})$, then all norms above are bounded. Then, by Step 1, $||v_n||_{L^{p+1}} \to 0$. In other words, $\tilde{u}_n \to u$ in $L^{p+1}(\mathbb{R})$.

From this it is easy to conclude. Indeed,

$$o_n(1) = J_{\omega}(\tilde{u}_n) = \frac{1}{2} \int_{-\infty}^{+\infty} \left(|\tilde{u}'_n|^2 + \omega \tilde{u}_n^2 \right) dr + \frac{1}{8} \left(\int_{-\infty}^{+\infty} \tilde{u}_n^2 dr \right)^3 - \frac{1}{p+1} \int_{-\infty}^{+\infty} |\tilde{u}_n|^{p+1} dr$$
$$0 \leqslant J_{\omega}(u) = \frac{1}{2} \int_{-\infty}^{+\infty} \left(|u'|^2 + \omega u^2 \right) dr + \frac{1}{8} \left(\int_{-\infty}^{+\infty} u^2 dr \right)^3 - \frac{1}{p+1} \int_{-\infty}^{+\infty} |u|^{p+1} dr.$$

Then, $\|\tilde{u}_n\|_{H^1(\mathbb{R})} \to \|u\|_{H^1(\mathbb{R})}$. And this implies that $\tilde{u}_n \to u$ in $H^1(\mathbb{R})$, finishing the proof.

4. PROOF OF THEOREMS 1.1, 1.2

Lemma 4.1. Let $U \in H^1(\mathbb{R})$ be an even function which decays to zero exponentially at infinity, and define $U_{\rho}(r) = U(r - \rho)$. Then there exists C > 0 such that:

$$I_{\omega}(U_{\rho}) = 2\pi\rho J_{\omega}(U) - C + o_{\rho}(1).$$

Proof. We have

$$(2\pi)^{-1}I_{\omega}(U_{\rho}) = \frac{1}{2} \int_{0}^{+\infty} \left(|U_{\rho}'|^{2} + \omega U_{\rho}^{2} \right) r \, dr + \frac{1}{8} \int_{0}^{+\infty} \frac{U_{\rho}^{2}(r)}{r} \left(\int_{0}^{r} s U_{\rho}^{2}(s) \, ds \right)^{2} dr$$

$$(19) \qquad -\frac{1}{p+1} \int_{0}^{+\infty} |U_{\rho}|^{p+1} r \, dr.$$

Let us, first of all, evaluate the local terms. By the evenness and the exponential decay of U, we get

$$\int_{0}^{+\infty} |U_{\rho}'|^{2} r \, dr = \int_{-\infty}^{+\infty} |U'(r-\rho)|^{2} (r-\rho) \, dr + \rho \int_{-\infty}^{+\infty} |U'(r-\rho)|^{2} \, dr + o_{\rho}(1)$$
(20)
$$= \rho \int_{-\infty}^{+\infty} |U'|^{2} \, dr + o_{\rho}(1).$$

Analogously

(21)
$$\int_{0}^{+\infty} U_{\rho}^{2} r \, dr = \rho \int_{-\infty}^{+\infty} U^{2} \, dr + o_{\rho}(1),$$

(22)
$$\int_0^{+\infty} |U_{\rho}|^{p+1} r \, dr = \rho \int_{-\infty}^{+\infty} |U|^{p+1} \, dr + o_{\rho}(1).$$

For what concerns the nonlocal term, we have

$$\begin{split} \int_{0}^{+\infty} \frac{U_{\rho}^{2}(r)}{r} \left(\int_{0}^{r} sU_{\rho}^{2}(s) \, ds \right)^{2} dr &- \rho \int_{0}^{+\infty} U_{\rho}^{2}(r) \left(\int_{0}^{r} U_{\rho}^{2}(s) \, ds \right)^{2} dr \\ &= \underbrace{\int_{0}^{+\infty} U_{\rho}^{2}(r) \left(\frac{1}{r} - \frac{1}{\rho} \right) \left(\int_{0}^{r} sU_{\rho}^{2}(s) \, ds \right)^{2} dr}_{(I)} \\ &+ \underbrace{\frac{1}{\rho} \int_{0}^{+\infty} U_{\rho}^{2}(r) \left[\left(\int_{0}^{r} sU_{\rho}^{2}(s) \, ds \right)^{2} - \left(\int_{0}^{r} \rho U_{\rho}^{2}(s) \, ds \right)^{2} \right] dr.}_{(II)} \end{split}$$

Let us study the term (I):

$$\begin{split} (I) &= \int_{-\infty}^{+\infty} U_{\rho}^{2}(r) \frac{\rho - r}{r\rho} \left(\int_{-\infty}^{r} sU_{\rho}^{2}(s) \, ds \right)^{2} dr + o_{\rho}(1) \\ &= -\int_{-\infty}^{+\infty} U^{2}(r) \frac{r}{(\rho + r)\rho} \left(\int_{-\infty}^{r} (s + \rho)U^{2}(s) \, ds \right)^{2} dr + o_{\rho}(1) \\ &= \int_{0}^{+\infty} U^{2}(r) \frac{r}{(\rho - r)\rho} \left(\int_{-\infty}^{r} (s + \rho)U^{2}(s) \, ds \right)^{2} dr \\ &- \int_{0}^{+\infty} U^{2}(r) \frac{r}{(\rho + r)\rho} \left(\int_{-\infty}^{r} (s + \rho)U^{2}(s) \, ds \right)^{2} dr + o_{\rho}(1) \\ &= \int_{0}^{+\infty} U^{2}(r) \left(\frac{r}{(\rho - r)\rho} - \frac{r}{(\rho + r)\rho} \right) \left(\int_{-\infty}^{-r} (s + \rho)U^{2}(s) \, ds \right)^{2} dr \\ &+ \int_{0}^{+\infty} U^{2}(r) \frac{r}{(\rho + r)\rho} \left[\left(\int_{-\infty}^{-r} (s + \rho)U^{2}(s) \, ds \right)^{2} - \left(\int_{-\infty}^{r} (s + \rho)U^{2}(s) \, ds \right)^{2} \right] dr \\ &+ o_{\rho}(1) \\ &= \frac{1}{\rho} \int_{0}^{+\infty} U^{2}(r) \left(\frac{2r^{2}\rho^{2}}{(\rho - r)(\rho + r)} \right) \left(\int_{-\infty}^{-r} \frac{s + \rho}{\rho} U^{2}(s) \, ds \right)^{2} dr \\ &+ \int_{0}^{+\infty} U^{2}(r) \frac{r\rho}{(\rho + r)} \left[\left(\int_{-\infty}^{-r} \frac{s + \rho}{\rho} U^{2}(s) \, ds \right)^{2} - \left(\int_{-\infty}^{r} \frac{s + \rho}{\rho} U^{2}(s) \, ds \right)^{2} \right] dr \\ &+ o_{\rho}(1). \end{split}$$

We now pass to the limit by Lebesgue Theorem, and obtain:

$$(I) = \int_0^{+\infty} U^2(r) r \left[\left(\int_{-\infty}^{-r} U^2(s) \, ds \right)^2 - \left(\int_{-\infty}^{r} U^2(s) \, ds \right)^2 \right] dr + o_\rho(1)$$

= $-C_I + o_\rho(1).$

Let us study the term (II):

$$(II) = \frac{1}{\rho} \int_0^{+\infty} U_{\rho}^2(r) \left(\int_0^r (s+\rho) U_{\rho}^2(s) \, ds \right) \left(\int_0^r (s-\rho) U_{\rho}^2(s) \, ds \right) dr$$

$$= \frac{1}{\rho} \int_{-\infty}^{+\infty} U_{\rho}^2(r) \left(\int_{-\infty}^r (s+\rho) U_{\rho}^2(s) \, ds \right) \left(\int_{-\infty}^r (s-\rho) U_{\rho}^2(s) \, ds \right) dr + o_{\rho}(1)$$

$$= \int_{-\infty}^{+\infty} U^2(r) \left(\int_{-\infty}^r \frac{s+2\rho}{\rho} U^2(s) \, ds \right) \left(\int_{-\infty}^r s U^2(s) \, ds \right) dr + o_{\rho}(1).$$

Again by Lebesgue Theorem,

$$(II) = 2 \int_{-\infty}^{+\infty} U^2(r) \left(\int_{-\infty}^{r} U^2(s) \, ds \right) \left(\int_{-\infty}^{r} s U^2(s) \, ds \right) dr + o_{\rho}(1)$$

= $-C_{II} + o_{\rho}(1).$

Observe that the above expression is negative since the function $r \mapsto \int_{-\infty}^{r} sU^{2}(s) ds$ is negative. Therefore, denoting by $C = C_{I} + C_{II} > 0$, we have (23)

$$\int_{0}^{+\infty} \frac{U_{\rho}^{2}(r)}{r} \left(\int_{0}^{r} s U_{\rho}^{2}(s) \, ds \right)^{2} dr = \rho \int_{0}^{+\infty} U_{\rho}^{2}(r) \left(\int_{0}^{r} U_{\rho}^{2}(s) \, ds \right)^{2} dr - C + o_{\rho}(1).$$

Hence the conclusion follows by (19), (20), (21), (22) and (23).

Hence the conclusion follows by (19), (20), (21), (22) and (23).

In our next result we study the behavior of unbounded sequences with energy bounded from above. This will be essential for the proof of Theorems 1.1, 1.2.

Proposition 4.2. Assume $\omega > 0$, and $u_n \in H^1_r(\mathbb{R}^2)$ such that $||u_n||$ is unbounded but $I_{\omega}(u_n)$ is bounded from above. Then, there exists a subsequence (still denoted by u_n) such that:

i) for all
$$\varepsilon > 0$$
, $\int_{\varepsilon \|u_n\|^2}^{+\infty} (|u'_n|^2 + u_n^2) dr \le C$;
ii) there exists $\delta \in (0, 1)$ such that $\int_{\delta \|u_n\|^2}^{\delta^{-1} \|u_n\|^2} (|u'_n|^2 + u_n^2) dr \ge c > 0$;
iii) $\|u_n\|_{L^2(\mathbb{R}^2)} \to +\infty$.

Proof. The beginning of the proof follows the ideas of [20, Theorem 4.3]. The main difference is that here we cannot conclude directly that I_{ω} is bounded from below, and indeed this fact depends on ω . The proof of Theorem 1.1 will require much more work.

We start using inequality (8) and Cauchy-Schwartz inequality to estimate I_{ω} :

(24)
$$I_{\omega}(u) \ge \frac{\pi}{2} \int_{0}^{+\infty} \left(|u'|^{2} + \omega u^{2} \right) r \, dr + \frac{\pi}{8} \int_{0}^{+\infty} \frac{u^{2}(r)}{r} \left(\int_{0}^{r} su^{2}(s) \, ds \right)^{2} dr + 2\pi \int_{0}^{+\infty} \left(\frac{\omega}{4} u^{2} + \frac{1}{8} u^{4} - \frac{1}{p+1} |u|^{p+1} \right) r \, dr.$$

Define

$$f: \mathbb{R}_+ \to \mathbb{R}, \qquad f(t) = \frac{\omega}{4}t^2 + \frac{1}{8}t^4 - \frac{1}{p+1}t^{p+1}.$$

Then, the set $\{t > 0 : f(t) < 0\}$ is of the form (α, β) , where α, β are positive constants depending only on p, ω . Moreover, we denote by $-c_0 = \min f < 0$.

For each function u_n , we define:

$$A_n = \{ x \in \mathbb{R}^2 : u_n(x) \in (\alpha, \beta) \}, \ \rho_n = \sup\{ |x| : x \in A_n \}.$$

With these definitions, we can rewrite (24) in the form (25)

$$I_{\omega}(u_n) \ge \frac{\pi}{2} \int_0^{+\infty} \left(|u_n'|^2 + \omega u_n^2 \right) r \, dr + \frac{\pi}{8} \int_0^{+\infty} \frac{u_n^2(r)}{r} \left(\int_0^r s u_n^2(s) \, ds \right)^2 dr - c_0 |A_n|.$$

In particular this implies that $|A_n|$ must diverge, and hence ρ_n . This already proves (iii).

By Strauss Lemma [21], we have

(26)
$$\alpha \leqslant u_n(\rho_n) \leqslant \frac{\|u_n\|}{\sqrt{\rho_n}}, \Rightarrow \|u_n\|^2 \geqslant \alpha^2 \rho_n.$$

We now estimate the nonlocal term. For that, define

(27)
$$B_n = A_n \cap B(0, \gamma_n), \text{ for } \gamma_n \in (0, \rho_n) \text{ such that } |B_n| = \frac{1}{2} |A_n|.$$

Then,

$$\int_{0}^{+\infty} \frac{u_{n}^{2}(r)}{r} \left(\int_{0}^{r} s u_{n}^{2}(s) \, ds \right)^{2} dr \geq \frac{1}{4\pi^{2}} \int_{\gamma_{n}}^{+\infty} \frac{u_{n}^{2}(r)}{r} \left(\int_{B_{n}} u_{n}^{2}(x) \, dx \right)^{2} dr$$
$$\geq c |A_{n}|^{2} \int_{\gamma_{n}}^{+\infty} \frac{u_{n}^{2}(r)}{r} \, dr$$
$$\geq c |A_{n}|^{2} \int_{A_{n} \setminus B_{n}} \frac{u_{n}^{2}(x)}{|x|^{2}} \, dx$$
$$\geq c \frac{|A_{n}|^{2}}{\rho_{n}^{2}} \int_{A_{n} \setminus B_{n}} u_{n}^{2}(x) \, dx$$
$$\geq c \frac{|A_{n}|^{2}}{\rho_{n}^{2}}.$$

$$(28)$$

Hence, by (24), (26) and (28), we get

$$I_{\omega}(u_n) \ge c\rho_n + c\frac{|A_n|^3}{\rho_n^2} - c_0|A_n| = \rho_n \left(c + c\frac{|A_n|^3}{\rho_n^3} - c_0\frac{|A_n|}{\rho_n}\right).$$

Observe that $t \mapsto c + ct^3 - c_0 t$ is strictly positive near zero and goes to $+\infty$, as $t \to +\infty$. Then we can assume, passing to a subsequence, that $|A_n| \sim \rho_n$. In other words, there exists m > 0 such that $\rho_n |A_n|^{-1} \to m$ as $n \to +\infty$. Taking into account (25) we conclude that up to a subsequence, $||u_n||^2 \sim \rho_n$. More-

Taking into account (25) we conclude that up to a subsequence, $||u_n||^2 \sim \rho_n$. Moreover, for any fixed $\varepsilon > 0$, we have:

$$C\rho_n \ge \|u_n\|_{L^2}^2 \ge \int_{\varepsilon\rho_n}^{+\infty} u_n^2 r \, dr \ge \varepsilon\rho_n \int_{\varepsilon\rho_n}^{+\infty} u_n^2 \, dr.$$

An analogous estimate works also for $\int_{\varepsilon\rho_n}^{+\infty} |u'_n|^2 dr$. This proves (i).

We now show that for some $\delta > 0$, $||u_n||_{H^1(\delta\rho_n,\rho_n)} \not\rightarrow 0$, which implies assertion (ii).

First, recall the definition of B_n and γ_n in (27). Then,

$$\int_{\gamma_n}^{\rho_n} u_n^2(r) \, dr \ge \rho_n^{-1} \int_{\gamma_n}^{\rho_n} u_n^2(r) r \, dr \ge \rho_n^{-1} \int_{A_n \setminus B_n} u_n^2(x) dx \ge \rho_n^{-1} |A_n \setminus B_n| \alpha^2 > c > 0.$$

To conclude it suffices to show that $\gamma_n \sim \rho_n$. Indeed, define

(29)
$$C_n = B_n \cap B(0, \tau_n), \text{ for } \tau_n \in (0, \gamma_n) \text{ such that } |C_n| = \frac{1}{2}|B_n|.$$

We can repeat the estimate (28) with A_n , B_n replaced with B_n , C_n respectively, to obtain that

$$\int_0^{+\infty} \frac{u_n^2(r)}{r} \left(\int_0^r s u_n^2(s) \, ds \right)^2 dr \ge c \frac{|B_n|^3}{\gamma_n^2}.$$

Hence,

$$I_{\omega}(u_{n}) \ge c\rho_{n} + c\frac{|A_{n}|^{3}}{\gamma_{n}^{2}} - c_{0}|A_{n}| = \gamma_{n} \left(c\frac{\rho_{n}}{\gamma_{n}} + c\frac{|A_{n}|^{3}}{\gamma_{n}^{3}} - c_{0}\frac{|A_{n}|}{\gamma_{n}} \right).$$

And we are done since $I_{\omega}(u_n)$ is bounded from above.

Proof of Theorem 1.1. If $\omega \in (0, \omega_0)$, then $J_{\omega}(w_{k_2}) < 0$ (see Proposition 3.2): applying Lemma 4.1 to $U = w_{k_2}$ we conclude assertion (i).

15

We now prove (*ii*) and (*iii*). Let us denote by $H^1_{0,r}(B(0,R))$ the Sobolev space of radial functions with zero boundary value. Given any $n \in \mathbb{N}$, Proposition 4.2 implies that $I_{\omega}|_{H^1_{0,r}(B(0,n))}$ is coercive (indeed, this is an immediate consequence of (24)). So, there exists u_n a minimizer for $I_{\omega}|_{H^1_{0,r}(B(0,n))}$. Moreover,

$$I_{\omega}(u_n) \to \inf I_{\omega}, \text{ as } n \to +\infty.$$

If u_n is bounded, then $I_{\omega}(u_n)$ is also bounded and therefore $\inf I_{\omega}$ is finite. In what follows we assume that u_n is an unbounded sequence. Then, the sequence u_n satisfies the hypotheses of Proposition 4.2. Let $\delta > 0$ be given by that proposition.

The proof will be divided in several steps.

Step 1:
$$\int_{\frac{\delta}{2}||u_n||^2}^{\frac{2}{\delta}||u_n||^2} |u_n|^{p+1} dr \not\rightarrow 0.$$

By Proposition 4.2, i), we have the

Proposition 4.2, i), we have that:

$$\sum_{k=1}^{\left[\frac{\delta}{2}\|u_n\|^2\right]} \int_{\frac{\delta}{2}\|u_n\|^2 + k-1}^{\frac{\delta}{2}\|u_n\|^2 + k} \left(|u_n'|^2 + u_n^2\right) dr \leqslant \int_{\frac{\delta}{2}\|u_n\|^2}^{\delta\|u_n\|^2} \left(|u_n'|^2 + u_n^2\right) dr \leqslant C.$$

Taking the smaller summand in the left hand side we find x_n ,

$$\frac{\delta}{2} \|u_n\|^2 \leqslant x_n \leqslant \delta \|u_n\|^2 - 1,$$

such that

$$||u_n||^2_{H^1(x_n,x_n+1)} \leq \frac{C}{||u_n||^2}$$

Reasoning in an analogous way, we can choose y_n ,

$$\delta^{-1} \|u_n\|^2 + 1 \leqslant y_n \leqslant 2\delta^{-1} \|u_n\|^2$$

such that

$$||u_n||^2_{H^1(y_n, y_n+1)} \leq \frac{C}{||u_n||^2}.$$

Observe that if $\delta^{-1} ||u_n||^2 \ge n$, the choice of y_n can be arbitrary, but it is unnecessary. Take $\phi_n: [0, +\infty] \to [0, 1]$ be a C^∞ -function such that

$$\phi_n(r) = \begin{cases} 0, & \text{if } r \leqslant x_n, \\ 1, & \text{if } x_n + 1 \leqslant r \leqslant y_n, \quad |\phi'_n(r)| \leqslant 2. \\ 0, & \text{if } r \geqslant y_n + 1. \end{cases}$$

We have

$$0 = I'_{\omega}(u_n)[\phi_n u_n] \ge 2\pi \int_{x_n}^{y_n} \left(|u'_n|^2 + \omega u_n^2 \right) r \, dr - 2\pi \int_{x_n}^{y_n} |u_n|^{p+1} r \, dr + O(1)$$
$$\ge \|u_n\|^2 \left(\frac{\delta}{2} \int_{x_n}^{y_n} \left(|u'_n|^2 + \omega u_n^2 \right) dr - \frac{2}{\delta} \int_{x_n}^{y_n} |u_n|^{p+1} \, dr \right) + O(1).$$

This, together with the fact that $||u_n||_{H^1(x_n,y_n)}$ does not tend to zero, allows us to conclude the proof of Step 1.

Step 2: Exponential decay.

At this point we can apply the concentration-compactness principle (see [16, Lemma 1.1]); there exists $\sigma > 0$ such that

$$\sup_{\xi\in[x_n,\ y_n]}\int_{\xi-1}^{\xi+1}u_n^2\,dr\geqslant 2\sigma>0.$$

Let us define:

(30)
$$D_n = \left\{ \xi > 0 : \int_{\xi - 1}^{\xi + 1} \left(|u'_n|^2 + u_n^2 \right) dr \ge \sigma \right\} \neq \emptyset, \text{ and } \xi_n = \max D_n \in [x_n, n + 1).$$

Let us observe that $\xi_n \sim ||u_n||^2$; indeed $\xi_n \ge x_n \ge c ||u_n||^2$ and, moreover,

$$||u_n||^2 \ge c \int_{\xi_n - 1}^{\xi_n + 1} \left(|u'_n|^2 + u_n^2 \right) r \, dr \ge c(\xi_n - 1) \int_{\xi_n - 1}^{\xi_n + 1} \left(|u'_n|^2 + u_n^2 \right) \, dr \ge c(\xi_n - 1).$$

By definition, $\int_{\zeta-1}^{\zeta+1} (|u'_n|^2 + u_n^2) dr < \sigma$ for all $\zeta > \xi_n$. By embedding of $H^1(\zeta - 1, \zeta + 1)$ in L^{∞} , $0 < u_n(\zeta) < C\sigma$ for any $\zeta > \xi_n$. From this we will get exponential decay of u_n . Indeed, u_n is a solution of

$$-u_n''(r) - \frac{u'(r)}{r} + \omega u_n(r) + f_n(r)u_n(r) = |u_n(r)|^{p-1}u_n(r),$$

with

$$f_n(r) = \frac{h_n^2(r)}{r^2} + \int_r^n \frac{h_n(s)}{s} u_n^2(s) \, ds, \quad h_n(r) = \frac{1}{2} \int_0^r u_n^2(s) s \, ds.$$

It is important to observe that $0 \leq f_n(r) \leq C$ for all $r > \delta ||u_n||^2$. Then, by taking smaller σ , if necessary, we can conclude that there exists C > 0 such that

$$|u_n(r)| < C \exp\left(-\sqrt{\omega}(r-\xi_n)\right), \quad \text{ for all } r > \xi_n.$$

The local C^1 regularity theory for the Laplace operator (see [6, Section 3.4]) implies a similar estimate for $u'_n(r)$. In other words,

(31)
$$|u_n(r)| + |u'_n(r)| < C \exp\left(-\sqrt{\omega}(r-\xi_n)\right), \quad \text{for all } r > \xi_n.$$

Step 3: Splitting of $I_{\omega}(u_n)$.

Reasoning as in the beginning of Step 1, we can take z_n :

$$\xi_n - 3 \|u_n\| \leqslant z_n \leqslant \xi_n - 2 \|u_n\|,$$

such that

$$||u_n||^2_{H^1(z_n, z_n+1)} \leq \frac{C}{||u_n||}.$$

Define $\psi_n: [0, +\infty] \to [0, 1]$ be a smooth function such that

(32)
$$\psi_n(r) = \begin{cases} 0, & \text{if } r \leq z_n, \\ 1, & \text{if } r \geq z_n + 1, \end{cases} \quad |\psi'_n(r)| \leq 2.$$

In what follows we want to estimate $I_{\omega}(u_n)$ with $I_{\omega}(\psi_n u_n)$ and $I_{\omega}((1 - \psi_n)u_n)$. Let us start evaluating the local terms.

$$\int_{0}^{n} |u_{n}'|^{2} r \, dr = \int_{0}^{n} |(u_{n}\psi_{n})'|^{2} r \, dr + \int_{0}^{n} \left| \left(u_{n}(1-\psi_{n}) \right)' \right|^{2} r \, dr + O(||u_{n}||),$$
$$\int_{0}^{n} u_{n}^{2} r \, dr = \int_{0}^{n} |u_{n}\psi_{n}|^{2} r \, dr + \int_{0}^{n} |u_{n}(1-\psi_{n})|^{2} r \, dr + O(||u_{n}||),$$
$$\int_{0}^{n} |u_{n}|^{p+1} r \, dr = \int_{0}^{n} |u_{n}\psi_{n}|^{p+1} r \, dr + \int_{0}^{n} |u_{n}(1-\psi_{n})|^{p+1} r \, dr + O(||u_{n}||).$$

Let us study now the nonlocal term.

$$\begin{split} &\int_{0}^{n} \frac{u_{n}^{2}(r)}{r} \left(\int_{0}^{r} su_{n}^{2}(s) \, ds \right)^{2} dr = \int_{0}^{n} \frac{u_{n}^{2}(r)\psi_{n}^{2}(r)}{r} \left(\int_{0}^{r} su_{n}^{2}(s)\psi_{n}^{2}(s) \, ds \right)^{2} dr \\ &+ \int_{0}^{n} \frac{u_{n}^{2}(r)(1-\psi_{n}(r))^{2}}{r} \left(\int_{0}^{r} su_{n}^{2}(s)(1-\psi_{n}(s))^{2} \, ds \right)^{2} dr \\ &+ \underbrace{\int_{0}^{n} \frac{u_{n}^{2}(r)\psi_{n}^{2}(r)}{r} \left(\int_{0}^{r} su_{n}^{2}(s)(1-\psi_{n}(s))^{2} \, ds \right)^{2} dr}_{(I)} \\ &+ 2\underbrace{\int_{0}^{n} \frac{u_{n}^{2}(r)\psi_{n}^{2}(r)}{r} \left(\int_{0}^{r} su_{n}^{2}(s)\psi_{n}^{2}(s) \, ds \right) \left(\int_{0}^{r} su_{n}^{2}(s)(1-\psi_{n}(s))^{2} \, ds \right) dr}_{(II)} \\ &+ O(||u_{n}||). \end{split}$$

We now estimate:

 $(I) \geqslant 0,$

$$(II) = \int_{z_n}^n \frac{u_n^2(r)\psi_n^2(r)}{r} \left(\int_{z_n}^r su_n^2(s)\psi_n^2(s)\,ds\right) \left(\int_0^{z_n+1} su_n^2(s)(1-\psi_n(s))^2\,ds\right)dr$$
$$+ O(\|u_n\|)$$
$$\geqslant c_n\|u_n(1-\psi_n)\|_{L^2(\mathbb{R}^2)}^2 + O(\|u_n\|),$$

where

$$c_n = \int_{z_n}^n \frac{u_n^2(r)\psi_n^2(r)}{r} \left(\int_{z_n}^r s u_n^2(s)\psi_n^2(s) \, ds\right) dr \ge c > 0.$$

Therefore, we get

(33)
$$I_{\omega}(u_n) \ge I_{\omega}(u_n\psi_n) + I_{\omega}(u_n(1-\psi_n)) + c \|u_n(1-\psi_n)\|_{L^2(\mathbb{R}^2)}^2 + O(\|u_n\|).$$

Step 4: The following estimate holds:

(34)
$$I_{\omega}(u_n\psi_n) = 2\pi\xi_n J_{\omega}(u_n\psi_n) + O(||u_n||).$$

Indeed, by taking into account Proposition 4.2, (31) and the definition of ψ_n (32), we have

$$\begin{aligned} \left| \int_{0}^{n} (u_{n}\psi_{n})^{2}r \, dr - \xi_{n} \int_{0}^{n} (u_{n}\psi_{n})^{2} \, dr \right| &\leq \int_{0}^{n} (u_{n}\psi_{n})^{2} |r - \xi_{n}| \, dr \\ &\leq \int_{\xi_{n} - 3\|u_{n}\|}^{\xi_{n} + \|u_{n}\|} u_{n}^{2} |r - \xi_{n}| \, dr + o(1) \\ &\leq O(\|u_{n}\|) \int_{\xi_{n} - 3\|u_{n}\|}^{\xi_{n} + \|u_{n}\|} u_{n}^{2} \, dr + o(1) = O(\|u_{n}\|). \end{aligned}$$

19

The estimates are similar for the other local terms of I_{ω} . For what concerns the nonlocal term, we get

$$\int_{0}^{n} \frac{(u_{n}\psi_{n})^{2}(r)}{r} \left(\int_{0}^{r} s(u_{n}\psi_{n})^{2}(s) \, ds\right)^{2} dr - \xi_{n} \int_{0}^{n} (u_{n}\psi_{n})^{2}(r) \left(\int_{0}^{r} (u_{n}\psi_{n})^{2}(s) \, ds\right)^{2} dr$$

$$= \underbrace{\int_{0}^{n} (u_{n}\psi_{n})^{2}(r) \left(\frac{1}{r} - \frac{1}{\xi_{n}}\right) \left(\int_{0}^{r} s(u_{n}\psi_{n})^{2}(s) \, ds\right)^{2} dr}_{(I)}$$

$$+ \underbrace{\frac{1}{\xi_{n}} \int_{0}^{n} (u_{n}\psi_{n})^{2}(r) \left[\left(\int_{0}^{r} s(u_{n}\psi_{n})^{2}(s) \, ds\right)^{2} - \left(\int_{0}^{r} \xi_{n}(u_{n}\psi_{n})^{2}(s) \, ds\right)^{2}\right] dr}_{(II)}$$

where

(35)
$$(I) \leq \int_{\xi_n - 3\|u_n\|}^{\xi_n + \|u_n\|} u_n^2(r) \frac{|\xi_n - r|}{r\xi_n} \left(\int_{\xi_n - 3\|u_n\|}^{\xi_n + \|u_n\|} su_n^2(s) \, ds \right)^2 dr + o(1) = O(\|u_n\|)$$

and

(36)

$$(II) \leqslant \frac{1}{\xi_n} \int_{\xi_n - 3\|u_n\|}^{\xi_n + \|u_n\|} u_n^2(r) \left| \int_{\xi_n - 3\|u_n\|}^{\xi_n + \|u_n\|} (s + \xi_n) u_n^2(s) \, ds \right| \left| \int_{\xi_n - 3\|u_n\|}^{\xi_n + \|u_n\|} (s - \xi_n) u_n^2(s) \, ds \right| dr$$

+ $o(1)$
= $O(\|u_n\|).$

Step 5: Conclusion for $\omega > \omega_0$.

By (33) and (34), we have

(37)
$$I_{\omega}(u_n) \ge 2\pi \xi_n J_{\omega}(u_n\psi_n) + I_{\omega}(u_n(1-\psi_n)) + c \|u_n(1-\psi_n)\|_{L^2(\mathbb{R}^2)}^2 + O(\|u_n\|).$$

Recall that $||u_n\psi_n||^2_{H^1(\mathbb{R})} \ge \sigma > 0$. By Proposition 3.7, we have that $J_{\omega}(u_n\psi_n) \rightarrow c > 0$, up to a subsequence. Since $\xi_n \sim ||u_n||^2$, it turns out from (37) that $I_{\omega}(u_n) > I_{\omega}(u_n(1-\psi_n))$. But this is a contradiction with the definition of u_n , which proves that $\inf I_{\omega} > -\infty$.

Let us now show that I_{ω} is coercive. Indeed, take $u_n \in H^1(\mathbb{R}^2)$ an unbounded sequence, and assume that $I_{\omega}(u_n)$ is bounded from above. By Proposition 4.2, (iii), we would obtain that $I_{\hat{\omega}}(u_n) \to -\infty$ for any $\omega_0 < \hat{\omega} < \omega$, a contradiction.

Step 6: Conclusion for $\omega = \omega_0$.

As above, (37) gives a contradiction unless $J_{\omega}(u_n\psi_n) \to 0$. Proposition 3.7 now implies that $\psi_n u_n(\cdot - t_n) \to w_{k_2}$ up to a subsequence, for some $t_n \in (0, +\infty)$. Since $\xi_n \in D_n$ (see their definition in (30)), we have that $|t_n - \xi_n|$ is bounded. With this extra information, we have a better estimate of the decay of the solutions: indeed,

(38)
$$|u_n(r)| + |u'_n(r)| < C \exp\left(-\sqrt{\omega}|r - \xi_n|\right), \text{ for all } r > \xi_n - 2||u_n||.$$

This allows us to do the cut-off procedure in a much more accurate way. Indeed, take

$$\tilde{z}_n = \xi_n - \|u_n\|.$$

Then, (38) implies that

(39)

$$||u_n||^2_{H^1(\tilde{z}_n, \tilde{z}_n+1)} \leq C \exp(-\sqrt{\omega}||u_n||).$$

Define $\tilde{\psi}_n : [0, +\infty] \to [0, 1]$ accordingly:

$$\tilde{\psi}_n(r) = \begin{cases} 0, & \text{if } r \leqslant \tilde{z}_n, \\ 1, & \text{if } r \geqslant \tilde{z}_n + 1, \end{cases} \quad |\tilde{\psi}'_n(r)| \leqslant 2.$$

The advantage is that, in the estimate of $I_{\omega}(u_n)$, now the errors are exponentially small. Indeed, by repeating the estimates of Step 3 with the new information (39), we obtain:

$$I_{\omega}(u_n) \ge I_{\omega}(u_n\tilde{\psi}_n) + I_{\omega}(u_n(1-\tilde{\psi}_n)) + c \|u_n(1-\tilde{\psi}_n)\|_{L^2(\mathbb{R}^2)}^2 + o(1).$$

Let us show that in this case (34) becomes

$$I_{\omega}(u_n\bar{\psi}_n) = 2\pi\xi_n J_{\omega}(u_n\bar{\psi}_n) + O(1).$$

Indeed, by (38) and (39), we have

$$\left|\int_0^n (u_n\tilde{\psi}_n)^2 r \, dr - \xi_n \int_0^n (u_n\tilde{\psi}_n)^2 \, dr\right| \leqslant \int_{-\infty}^{+\infty} (u_n\tilde{\psi}_n)^2 |r - \xi_n| \, dr \leqslant C;$$

the other local terms can be estimated similarly. For what concerns the nonlocal term, we repeat the arguments of the previous case using in (35) and (36) the informations contained in (38) and (39).

$$I_{\omega}(u_n) \ge I_{\omega}(u_n\psi_n) + I_{\omega}(u_n(1-\psi_n)) + c \|u_n(1-\psi_n)\|_{L^2(\mathbb{R}^2)}^2 + O(1)$$

= $2\pi\xi_n J_{\omega}(u_n\tilde{\psi}_n) + I_{\omega}(u_n(1-\tilde{\psi}_n)) + c \|u_n(1-\tilde{\psi}_n)\|_{L^2(\mathbb{R}^2)}^2 + O(1)$
 $\ge I_{(\omega+2c)}(u_n(1-\tilde{\psi}_n)) + O(1).$

But, by Case 1, we already know that $I_{(\omega+2c)}$ is bounded from below, and hence $\inf I_{\omega_0} > -\infty$.

Finally, applying Lemma 4.1 to $U = w_{k_2}$, we readily get that I_{ω_0} is not coercive.

Proof of Theorem **1.2***.* We shall prove each assessment separately.

Proof of (i). Let u be a solution of (6). We multiply (6) by u and integrate: taking into account the inequality (8), we get

0

$$0 = \int_{\mathbb{R}^2} \left(|\nabla u|^2 + \omega u^2 \right) dx + \frac{3}{4} \int_{\mathbb{R}^2} \frac{u^2(x)}{|x|^2} \left(\int_0^{|x|} s u^2(s) \, ds \right)^2 dx - \int_{\mathbb{R}^2} |u|^{p+1} dx$$
$$\geqslant \frac{1}{4} \int_{\mathbb{R}^2} |\nabla u|^2 dx + \int_{\mathbb{R}^2} \left(\omega u^2 + \frac{3}{4} u^4 - |u|^{p+1} \right) dx.$$

Observe that there exists $\bar{\omega} > 0$ such that, for $\omega > \bar{\omega}$, the function $t \mapsto \omega t^2 + \frac{3}{4}t^4 - |t|^{p+1}$ is non-negative. Therefore u must be identically zero.

Proof of (ii). First, we observe that since $\inf I_{\omega_0} < 0$, there exists $\tilde{\omega} > \omega_0$ such that $\inf I_{\omega} < 0$ if and only if $\omega \in (\omega_0, \tilde{\omega})$. Since, by Theorem 1.1 and Proposition 2.2, I_{ω} is coercive and weakly lower semicontinuous, we infer that the infimum is attained.

Clearly, 0 is a local minimum for I_{ω} . Then, if $\omega \in (\omega_0, \tilde{\omega})$, so the functional satisfies the geometrical assumptions of the Mountain Pass Theorem, see [1]. Since I_{ω} is coercive, (PS) sequences are bounded. By the compact embedding of $H_r^1(\mathbb{R}^2)$ into $L^{p+1}(\mathbb{R}^2)$ and Proposition 2.2, standard arguments show that I_{ω} satisfies the

Palais-Smale condition and so we find a second solution which is at a positive energy level.

Proof of (iii). Let now consider $\omega \in (0, \omega_0)$. Performing the rescaling $u \mapsto u_\omega = \sqrt{\omega} u(\sqrt{\omega} \cdot)$, we get

$$I_{\omega}(u_{\omega}) = \omega \left[\frac{1}{2} \int_{\mathbb{R}^2} \left(|\nabla u|^2 + u^2 \right) dx + \frac{1}{8} \int_{\mathbb{R}^2} \frac{u^2(x)}{|x|^2} \left(\int_0^{|x|} s u^2(s) \, ds \right)^2 dx - \frac{\omega^{\frac{p-3}{2}}}{p+1} \int_{\mathbb{R}^2} |u|^{p+1} \, dx \right].$$

Define $\lambda = \omega^{\frac{p-3}{2}}$ and $\mathcal{I}_{\lambda} : H^1_r(\mathbb{R}^2) \to \mathbb{R}$ as

$$\begin{aligned} \mathcal{I}_{\lambda}(u) &= \frac{1}{2} \int_{\mathbb{R}^2} \left(|\nabla u|^2 + u^2 \right) dx + \frac{1}{8} \int_{\mathbb{R}^2} \frac{u^2(x)}{|x|^2} \left(\int_0^{|x|} s u^2(s) \, ds \right)^2 dx \\ &- \frac{\lambda}{p+1} \int_{\mathbb{R}^2} |u|^{p+1} \, dx. \end{aligned}$$

Since \mathcal{I}_{λ} satisfies the geometrical assmptions of the Mountain Pass Theorem, by [15, Theorem 1.1], we infer that, for almost every λ , the functional \mathcal{I}_{λ} possesses a bounded Palais-Smale sequence u_n . Assume $u_n \rightarrow u$; Proposition 2.2 and standard arguments imply that u is a critical point of \mathcal{I}_{λ} . Making the change of variables back we obtain a solution of (6) for almost every $\omega \in (0, \omega_0)$.

Finally, in order to find positive solutions of (6), we simply observe that the whole argument applies to the functional $I^+_{\omega}: H^1_r(\mathbb{R}^2) \to \mathbb{R}$

$$I_{\omega}^{+}(u) = \frac{1}{2} \int_{\mathbb{R}^{2}} \left(|\nabla u|^{2} + \omega u^{2} \right) dx + \frac{1}{8} \int_{\mathbb{R}^{2}} \frac{u^{2}(x)}{|x|^{2}} \left(\int_{0}^{|x|} su^{2}(s) \, ds \right)^{2} dx - \frac{1}{p+1} \int_{\mathbb{R}^{2}} (u^{+})^{p+1} \, dx.$$

Due to the maximum principle, the critical points of I_{ω}^+ are positive solutions of (6).

REFERENCES

- A. Ambrosetti and P. H. Rabinowitz, Dual variational methods in critical point theory and applications, J. Functional Analysis 14 (1973), 349-381.
- [2] L. Bergé, A. de Bouard and J. C. Saut, Blowing up time-dependent solutions of the planar Chern-Simons gauged nonlinear Schrödinger equation, Nonlinearity 8 (1995), 235-253.
- [3] H. Brézis, and E. Lieb, A relation between pointwise convergence of functions and convergence of functionals, Proc. Amer. Math. Soc. 88 (1983), 486-490.
- [4] J. Byeon, H. Huh, and J. Seok, Standing waves of nonlinear Schrödinger equations with the gauge field, J. Funct. Anal. 263 (2012), no. 6, 1575-1608.
- [5] V. Dunne, Self-dual Chern-Simons Theories, Springer, 1995.
- [6] D. Gilbarg and N. S. Trudinger, Elliptic Partial Differential Equations of Second Order, Springer, 3rd edition, 1998.
- [7] C. Hagen, A new gauge theory without an elementary photon, Ann. of Phys. 157 (1984), 342-359.
- [8] C. Hagen, Rotational anomalies without anyons, Phys. Review D 31 (1985), 2135-2136.
- [9] H. Huh, Blow-up solutions of the Chern-Simons-Schrödinger equations, Nonlinearity 22 (2009) 967-974.
- [10] H. Huh, Standing waves of the Schrödinger equation coupled with the Chern-Simons gauge field. J. Math. Phys. 53 (2012), no. 6, 063702, 8 pp.

- [11] H. Huh, Energy Solution to the Chern-Simons-Schrödinger Equations, Abstract and Applied Analysis Volume 2013, Article ID 590653, 7 pages.
- [12] R. Jackiw and S.-Y. Pi, Soliton solutions to the gauged nonlinear Schrödinger Equations, Phys. Review Letters 64 (1990), 2969-2972.
- [13] R. Jackiw and S.-Y. Pi, Classical and quantal nonrelativistic Chern-Simons theory, Phys. Rev. D 42 (1990) 3500-3513.
- [14] R. Jackiw and S.-Y. Pi, Self-dual Chern-Simons solitons, Progr. Theoret. Phys. Suppl. 107 (1992) 1-40.
- [15] L. Jeanjean, On the existence of bounded Palais-Smale sequences and applications to a Landesman-Lazer type problem set on \mathbb{R}^N , Proc. Roy. Soc. Edinburgh A 129 (1999) 787-809.
- [16] P.-L. Lions, The concentration-compactness principle in the calculus of variations. The locally compact case, part 1 and 2, Ann. Inst. Henri. Poincaré Anal. Non linéaire 1 (1984), 109-145 and 223-283.
- [17] B. Liu and P. Smith, Global wellposedness of the equivariant Chern-Simons-Schrödinger equation, preprint arXiv:1312.5567.
- [18] B. Liu, P. Smith, and D. Tataru, Local wellposedness of Chern-Simons-Schrödinger, Int. Math. Res. Notices, doi: 10.1093/imrn/rnt161.
- [19] S.-J. Oh, F. Pusateri, Decay and Scattering for the Chern-Simons-Schrödinger Equations, preprint arXiv:1311.2088.
- [20] D. Ruiz, The Schrödinger-Poisson equation under the effect of a nonlinear local term, J. Funct. Anal. 237 (2006), 655-674.
- [21] W.-A. Strauss, Existence of solitary waves in higher dimensions, Comm. Math. Phys. 55 (1977), 149-162.
- [22] M. Struwe, On the evolution of harmonic mappings of Riemannian surfaces, Comment. Math. Helv. 60 (1985) 558-581.
- [23] G. Tarantello, Self-Dual Gauge Field Vortices: An Analytical Approach, PNLDE 72, Birkhäuser Boston, Inc., Boston, MA, 2007.

¹DIPARTIMENTO DI MECCANICA, MATEMATICA E MANAGEMENT, POLITECNICO DI BARI, VIA E. Orabona 4, 70125 Bari, Italy.

²DPTO. ANÁLISIS MATEMÁTICO, GRANADA, 18071 SPAIN. *E-mail address*: a.pomponio@poliba.it, daruiz@ugr.es